

World Journal of *Gastroenterology*

World J Gastroenterol 2019 September 7; 25(33): 4796-5016



**EDITORIAL**

- 4796** Role of NLRP3 inflammasome in inflammatory bowel diseases
Tourkochristou E, Aggeletopoulou I, Konstantakis C, Triantos C

OPINION REVIEW

- 4805** Gastroesophageal reflux disease, obesity and laparoscopic sleeve gastrectomy: The burning questions
Bou Daher H, Sharara AI

REVIEW

- 4814** Intestinal permeability in the pathogenesis of liver damage: From non-alcoholic fatty liver disease to liver transplantation
Nicoletti A, Ponziani FR, Biolato M, Valenza V, Marrone G, Sganga G, Gasbarrini A, Miele L, Grieco A
- 4835** Crosstalk network among multiple inflammatory mediators in liver fibrosis
Zhangdi HJ, Su SB, Wang F, Liang ZY, Yan YD, Qin SY, Jiang HX

MINIREVIEWS

- 4850** Neoadjuvant radiotherapy for rectal cancer management
Feeney G, Sehgal R, Sheehan M, Hogan A, Regan M, Joyce M, Kerin M
- 4870** *Helicobacter pylori* virulence genes
Šterbenc A, Jarc E, Poljak M, Homan M
- 4885** Occupational exposure to vinyl chloride and liver diseases
Fedeli U, Girardi P, Mastrangelo G

ORIGINAL ARTICLE**Basic Study**

- 4892** *Ex vivo* effect of vascular wall stromal cells secretome on enteric ganglia
Dothel G, Bernardini C, Zannoni A, Spirito MR, Salaroli R, Bacci ML, Forni M, Ponti FD
- 4904** Towards a standard diet-induced and biopsy-confirmed mouse model of non-alcoholic steatohepatitis: Impact of dietary fat source
Boland ML, Oró D, Tølbøl KS, Thrane ST, Nielsen JC, Cohen TS, Tabor DE, Fernandes F, Tovchigrechko A, Veidal SS, Warrenner P, Sellman BR, Jelsing J, Feigh M, Vrang N, Trevaskis JL, Hansen HH
- 4921** Identification of hepatitis B virus and liver cancer bridge molecules based on functional module network
Huang XB, He YG, Zheng L, Feng H, Li YM, Li HY, Yang FX, Li J

Retrospective Cohort Study

- 4933** Proton pump inhibitor use increases mortality and hepatic decompensation in liver cirrhosis
De Roza MA, Kai L, Kam JW, Chan YH, Kwek A, Ang TL, Hsiang JC
- 4945** Prognostic value of preoperative carcinoembryonic antigen/tumor size in rectal cancer
Cai D, Huang ZH, Yu HC, Wang XL, Bai LL, Tang GN, Peng SY, Li YJ, Huang MJ, Cao GW, Wang JP, Luo YX

Retrospective Study

- 4959** Value of controlled attenuation parameter in fibrosis prediction in nonalcoholic steatohepatitis
Lee JJ, Lee HW, Lee KS
- 4970** Lymphocyte-to-monocyte ratio effectively predicts survival outcome of patients with obstructive colorectal cancer
Chen XQ, Xue CR, Hou P, Lin BQ, Zhang JR

Observational Study

- 4985** Tenofovir is a more suitable treatment than entecavir for chronic hepatitis B patients carrying naturally occurring rtM204I mutations
Choe WH, Kim K, Lee SY, Choi YM, Kwon SY, Kim JH, Kim BJ

META-ANALYSIS

- 4999** Efficacy of *Lactobacillus rhamnosus* GG in treatment of acute pediatric diarrhea: A systematic review with meta-analysis
Li YT, Xu H, Ye JZ, Wu WR, Shi D, Fang DQ, Liu Y, Li LJ

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Amedeo Lonardo, MD, Doctor, Senior Lecturer, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena 41126, Italy

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. The *WJG* Editorial Board consists of 701 experts in gastroenterology and hepatology from 58 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, etc. The *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2019 edition of Journal Citation Report® cites the 2018 impact factor for *WJG* as 3.411 (5-year impact factor: 3.579), ranking *WJG* as 35th among 84 journals in gastroenterology and hepatology (quartile in category Q2). CiteScore (2018): 3.43.

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Yu-Jie Ma*

Proofing Production Department Director: *Yun-Xiaojuan Wu*

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Subrata Ghosh, Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE

Ze-Mao Gong, Director

PUBLICATION DATE

September 7, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Efficacy of *Lactobacillus rhamnosus* GG in treatment of acute pediatric diarrhea: A systematic review with meta-analysis

Ya-Ting Li, Hong Xu, Jian-Zhong Ye, Wen-Rui Wu, Ding Shi, Dai-Qiong Fang, Yang Liu, Lan-Juan Li

ORCID number: Ya-Ting Li (0000-0002-0761-4967); Hong Xu (0000-0001-6453-4683); Jian-Zhong Ye (0000-0001-8174-2580); Wenrui Wu (0000-0002-8457-9675); Ding Shi (0000-0003-3988-9321); Dai-Qiong Fang (0000-0002-0758-3988); Yang Liu (0000-0003-0246-4418); Lan-Juan Li (0000-0001-6945-0593).

Author contributions: Li YT, Xu H, and Ye JZ contributed equally to this work; Li YT and Xu H identified eligible articles and extracted applicable data; Li YT, Xu H, and Ye JZ contributed to the design of the study; Wu WR contributed to the analysis and interpretation of the outcomes; Liu Y, Fang DQ, and Shi D participated in writing and editing the article; all authors approved the final draft of the manuscript.

Supported by the National Natural Science Foundation of China, No. 81330011.

Conflict-of-interest statement: None.

PRISMA 2009 Checklist statement: The guidelines of the PRISMA 2009 Statement have been adopted in this manuscript.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works

Ya-Ting Li, Jian-Zhong Ye, Wen-Rui Wu, Ding Shi, Dai-Qiong Fang, Lan-Juan Li, State Key Laboratory for the Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou 310003, Zhejiang Province, China

Ya-Ting Li, Jian-Zhong Ye, Wen-Rui Wu, Ding Shi, Dai-Qiong Fang, Lan-Juan Li, Collaborative Innovation Center for the Diagnosis and Treatment of Infectious Diseases, School of Medicine, Zhejiang University, Hangzhou 310003, Zhejiang Province, China

Hong Xu, Department of Orthopedics, Xiaoshan Traditional Chinese Medical Hospital, Hangzhou 310003, Zhejiang Province, China

Yang Liu, Department of Orthopedics, Clinical Sciences, Lund, Lund University, Lund 22185, Sweden

Corresponding author: Lan-Juan Li, MD, PhD, Academic Research, Doctor, Professor, Senior Researcher, State Key Laboratory for the Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, No. 79, Qingchun Road, Hangzhou 310003, Zhejiang Province, China. ljli@zju.edu.cn

Telephone: +86-571-87236759

Fax: +86-571-87236459

Abstract

BACKGROUND

Diarrhea is a major infectious cause of childhood morbidity and mortality worldwide. In clinical trials, *Lactobacillus rhamnosus* GG ATCC 53013 (LGG) has been used to treat diarrhea. However, recent randomized controlled trials (RCTs) found no evidence of a beneficial effect of LGG treatment.

AIM

To evaluate the efficacy of LGG in treating acute diarrhea in children.

METHODS

The EMBASE, MEDLINE, PubMed, Web of Science databases, and the Cochrane Central Register of Controlled Trials were searched up to April 2019 for meta-analyses and RCTs. The Cochrane Review Manager was used to analyze the relevant data.

RESULTS

Nineteen RCTs met the inclusion criteria and showed that compared with the control group, LGG administration notably reduced the diarrhea duration [mean difference (MD) -24.02 h, 95% confidence interval (CI) (-36.58, -11.45)]. More

on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: April 28, 2019

Peer-review started: April 28, 2019

First decision: May 30, 2019

Revised: July 4, 2019

Accepted: July 19, 2019

Article in press: July 19, 2019

Published online: September 7, 2019

P-Reviewer: Tuna Kirsaciloglu CT, Reyes VEE, Rhoads JM

S-Editor: Yan JP

L-Editor: Wang TQ

E-Editor: Wu YXJ



effective results were detected at a high dose $\geq 10^{10}$ CFU per day [MD -22.56 h, 95%CI (-36.41, -8.72)] *vs* a lower dose. A similar reduction was found in Asian and European patients [MD -24.42 h, 95%CI (-47.01, -1.82); MD -32.02 h, 95%CI (-49.26, -14.79), respectively]. A reduced duration of diarrhea was confirmed in LGG participants with diarrhea for less than 3 d at enrollment [MD -15.83 h, 95%CI (-20.68, -10.98)]. High-dose LGG effectively reduced the duration of rotavirus-induced diarrhea [MD -31.05 h, 95%CI (-50.31, -11.80)] and the stool number per day [MD -1.08, 95%CI (-1.87, -0.28)].

CONCLUSION

High-dose LGG therapy reduces the duration of diarrhea and the stool number per day. Intervention at the early stage is recommended. Future trials are expected to verify the effectiveness of LGG treatment.

Key words: *Lactobacillus rhamnosus* GG; Acute diarrhea; Children; Rotavirus; Probiotics; Systematic review; Meta-analysis

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The treatment effectiveness of *Lactobacillus rhamnosus* GG (LGG) for acute diarrhea in children was assessed in our study. LGG was confirmed to effectively reduce the duration of diarrhea and the stool number per day. LGG was particularly efficacious in patients treated at a dose $> 10^{10}$ CFU/day, those treated at an early stage of illness, and those diagnosed with rotavirus-positive diarrhea.

Citation: Li YT, Xu H, Ye JZ, Wu WR, Shi D, Fang DQ, Liu Y, Li LJ. Efficacy of *Lactobacillus rhamnosus* GG in treatment of acute pediatric diarrhea: A systematic review with meta-analysis. *World J Gastroenterol* 2019; 25(33): 4999-5016

URL: <https://www.wjgnet.com/1007-9327/full/v25/i33/4999.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v25.i33.4999>

INTRODUCTION

The World Health Organization and United Nations International Children's Emergency Fund define diarrhea as more than three loose or watery stools during a 24-h period. A duration of 14 days is the proposed criterion for acute diarrhea or persistent diarrhea. Diarrhea is a major infectious cause of childhood morbidity and mortality worldwide, especially in developing countries^[1]. As the second most common cause of death among children under 5 years of age^[2], the frequency of acute diarrhea in one year is approximately two to three episodes per child^[1]. Previous data showed that the incidence of diarrhea was 6 to 12 episodes in 12 months per child in developing countries^[3].

The goals of treatment are prevention or resolution of dehydration and reduction of the diarrhea duration and infectious period^[4]. Oral rehydration, gut motility inhibitors, and antibiotics are used to treat acute gastroenteritis^[4]. Oral rehydration contributes to a reduced likelihood of dehydration but has no appreciable effects on bowel movements or the duration of diarrhea and is not utilized to its full extent^[5]. Antibiotics should be considered if pathogenic bacteria are detected. Smectite and zinc remain under-utilized as adjuvant therapies^[6,7].

Probiotic supplements have gained considerable popularity in the global market and are predicted to generate 64 billion United States dollars in revenue by 2023^[8]. Probiotics have health benefits for hosts^[9] and have been evaluated in the treatment of diarrhea, and multiple mechanisms of diarrhea improvement have been identified. Probiotics modulate the host immune response^[10]. Furthermore, colonic bacterial metabolites such as short-chain fatty acids increase colonic Na and fluid absorption through a cyclic adenosine monophosphate-independent mechanism^[5]. In clinical trials, the well-known probiotics *Saccharomyces boulardii*, *Lactobacillus reuteri* DSM 17938, and *Lactobacillus rhamnosus* GG ATCC 53013 (LGG) have been used to treat diarrhea^[2,4]. Previously, rotavirus-induced diarrhea was considered an adaptation disease associated with LGG treatment^[11]. Wolvers D revealed that the probiotic dose mediated the effectiveness of treatment, and 10^{10} - 10^{11} CFU per day was

recommended^[12]. In addition, a greater effect was observed in the early stage of illness, and a poorer effect on invasive bacterial diarrhea versus watery diarrhea was observed. LGG treatment has been endorsed by leading experts^[13-15]. However, most recent randomized controlled trials (RCTs) conducted by Schnadower *et al*^[9] yielded no evidence of a beneficial effect of LGG treatment. Therefore, we conducted a meta-analysis to evaluate the available validated data and update existing knowledge and thus provide guidance to patients.

MATERIALS AND METHODS

Literature search

Relevant studies published before April 2019 were retrieved from the EMBASE, MEDLINE, PubMed, Web of Science databases, and the Cochrane Central Register of Controlled Trials (CENTRAL, the Cochrane Library). The search strategy was conducted with medical subject headings and the search terms “diarrhoea, diarrhea, diarrh*, gastroenteritis, probiotic*, *Lactobacillus rhamnosus* GG, *Lactobacillus* GG, and LGG”. No language restrictions were applied. Additional studies were identified by manually searching review articles.

Study selection

Nineteen RCTs describing LGG interventions for acute diarrhea were included. The PRISMA statement and the guidelines from the Cochrane Collaboration were followed for this evidence-based medicine study^[16,17]. The participants were children aged less than 18 years. The dose of LGG was provided in various forms at different times. Antibiotic-associated diarrhea and persistent diarrhea were excluded. Other applications of LGG, such as preventive strategies, were not included. Some particular article types without complete data were excluded, such as abstracts and letters. We also excluded studies using mixtures of more than one probiotic strain. The primary outcomes were directly related to the development of persistent diarrhea, including the duration of diarrhea and diarrhea lasting ≥ 3 and ≥ 4 d. Secondary outcomes included the hospital stay duration, stool frequency, and improvement in stool consistency and vomiting.

Data extraction

Two investigators (Li YT and Xu H) independently identified eligible articles and extracted applicable data following the inclusion criteria. Quality control was assessed by another reviewer (Wu WR). The data set included the baseline characteristics of the participants, the duration of diarrhea, the hospital stay duration, the time to improvement in stool consistency, the mean number of stools per day during diarrhea episodes, the proportion of patients with vomiting, the duration of vomiting, stool frequency on days 2 and 3 after treatment, and the number of patients with diarrhea lasting ≥ 3 or 4 d. Cochrane Review Manager (Version 5.1. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2011) and STATA version 12.0 (StataCorp LP, College Station, TX, United States) were used for data analyses. Any discrepancies were resolved by discussion.

Risk of bias

All included trials were evaluated following the Cochrane Collaboration’s risk of bias tool. Seven domains were examined to identify the bias risk: selection bias, including random sequence generation and allocation concealment, performance bias, including blinding of participants and personnel, detection bias, including blinding of outcome assessments, attrition bias, including incomplete outcome data, reporting bias, including selective reporting, and other bias. Adequate allocation concealment was implemented to ensure blinding of the participants and investigators to avoid influences on the measures. Randomization was performed based on confirmed allocation concealment. Unclear allocation concealment was noted when no method was mentioned. The integrity of the data was evaluated, including the proportion of excluded participants (<http://www.cochrane-handbook.org>).

Statistical analysis

The Cochrane Review Manager was used to analyze the relevant data. The mean differences (MDs) in continuous data under LGG or placebo treatment were measured. Dichotomous results are pooled and presented as risk ratios. Additionally, 95% confidence intervals (CIs) are reported for all types of outcomes. I^2 and χ^2 values were calculated to quantify and reflect heterogeneity. A P -value < 0.05 indicates that heterogeneity should not be ignored; thus, a random-effects model was used. A fixed-effects model was employed when no statistically significant inconsistency was

detected. Publication bias was assessed by funnel plot asymmetry^[18]. Sensitivity analyses were conducted to detect the robustness of results by assessing randomization, missing data, blinding, and allocation concealment. Each individual study was systematically removed from the meta-analysis, and the effect was recalculated and estimated from the remaining studies (Supporting information Figure S1). Regression analysis was conducted, and the relationships between the duration of diarrhea and other covariates, including publication year, participant age, the duration of diarrhea before study enrollment, and the LGG dosage, were examined. Subgroup analyses were performed to diminish significant inconsistency. Preplanned subgroup analyses were performed according to the following clinical characteristics and results from sensitivity or regression analysis: (1) The dosage of LGG per day. A dosage of 10^{10} CFU/day was observed to be a critical element of effective treatment in the study by Szajewska *et al*^[13]. In addition, a larger dose was suggested in other studies^[19,20]; (2) The etiology of diarrhea. Diarrhea mortality and severe diarrhea were most frequently caused by rotavirus in children^[21]. Compared to control children, several rotavirus-positive children with watery stools in a probiotic group were reported to exhibit a marked reduction in diarrhea symptoms after 24 h^[22]. A meta-analysis performed by Szajewska *et al*^[23] in 2007 concluded that the duration of rotavirus-induced diarrhea was significantly attenuated by LGG supplementation; (3) The site of treatment (inpatient *vs* outpatient); (4) Vaccination status; (5) Geography of the clinical trials. The location of the study affected the sanitary habits, exposure to various pathogens, and nutrient status of the participants. All studied environmental factors contribute to various outcomes; (6) Early probiotic administration. A beneficial effect of probiotics was reported in the course of disease when initiated early^[12]; and (7) Publication date.

RESULTS

Study selection

A total of 349 potentially relevant studies were identified. The process of screening was carried out according to the flow diagram shown in Figure S2 (Supporting information). The characteristics of each included study are summarized in Table 1. With 988 participants in a 2007 meta-analysis and 2683 participants in a 2013 meta-analysis, a total of 4073 participants in 19 RCTs were identified in the literature. Two experimental arms in the study of Basu *et al*^[24] were listed separately to exhibit different doses of probiotics, which were marked as Basu 2009a and Basu 2009b. Therefore, the figures, tables, and full texts of 18 articles were reviewed^[8,24-40]. A large number of trials were conducted in Europe and Asia. Patients were recruited from outpatient, inpatient, and emergency departments. Inconsistency existed in the daily doses and routes of LGG supplementation during the treatment period. Different criteria were used to define diarrhea in the included studies. Diarrhea resolution was commonly defined as passage of the first normal stool or the last watery stool.

Antibiotic treatment before recruitment was assessed, and different studies varied regarding the use of antibiotics. Similarly, the duration of treatment varied. Studies of moderate to high quality were adequately assessed and are summarized in Figure S3 (Supporting information).

Evaluation before enrollment (days)

Before enrollment, age was assessed in 16 studies, and the duration of diarrhea was reported in nine studies (Supporting information Figures S4 and S5). No obvious difference in age was found. The statistical differences and high heterogeneity resulting from the duration of diarrhea [MD -6.21 h, 95%CI (-9.04, -3.38)] could be reduced by subgrouping according to the outcomes of the sensitivity analysis (Supporting information Figure S1). The subgroup excluding the study of Ritchie *et al*^[37] performed in 2010 showed acceptable heterogeneity, and no statistical significance was observed for the duration of diarrhea before study enrollment [MD -0.9 h, 95%CI (-4.02, 2.22)] ($I^2 = 10\%$). Sensitivity analysis revealed differences in the duration of diarrhea before study enrollment between the two groups in the study of Ritchie *et al*^[37], which recruited aboriginal children in the Northern Territory of Australia. Social disadvantages and poverty contributed to malnutrition in these children^[4]. However, no significant differences in the primary and secondary outcomes were found by sensitivity analysis, which is inconsistent with the findings reported in previous meta-analyses^[4,13] (Supporting information Figure S1).

Duration of diarrhea

A reduced duration of diarrhea was found in the LGG group compared to that in the

Table 1 Characteristics of the included trials

Article	Type of article	Age group	Country	Patient source	n (exp/control)	Inclusion criteria	Exclusion criteria	LGG (dosage)	Control group	Duration of intervention	Etiology
Basu <i>et al</i> ^[23] , 2007	RCT; 1 center; Duration: 1 yr	Children	India	Inpatients	323/323	≥ 3 watery stools/day without visible blood or mucus; <10 white blood cells/high-power field and no red cells, mucus flakes, or bacteria on stool microscopy; negative hanging drop preparation; negative bacterial stool culture	Systemic illness other than diarrhea on admission; systemic complications of diarrhea during hospitalization; failure to provide informed consent	120 × 10 ⁶ ; CFU/day	ORF	7 d	Bacterial diarrhea excluded; Rotavirus-induced diarrhea 75.8%
Basu <i>et al</i> ^[24] , 2009a	RCT; 1 center; Duration: 1 yr	Children	India	Inpatients	188/185	≥ 3 watery stools/day without macroscopic blood or mucus, <10 white blood cells/high-power field, and no red blood cells, mucus flakes, or bacteria on stool microscopy; negative hanging drop preparation; negative bacterial stool culture	Symptoms of illness other than diarrhea; development of any systemic complication of diarrhea during hospitalization; failure to provide informed consent	2 × 10 ¹⁰ ; CFU/day	ORF	7 d or until diarrhea stopped	Bacterial diarrhea excluded; Rotavirus diarrhea 57.1%
Basu <i>et al</i> ^[24] , 2009b	RCT; 1 center; Duration: 1 yr	Children	India	Inpatients	186/185	≥ 3 watery stools/day without macroscopic blood or mucus, <10 white blood cells/high-power field, and no red blood cells, mucus flakes, or bacteria on stool microscopy; negative hanging drop preparation; negative bacterial stool culture	Symptoms of illness other than diarrhea; development of any systemic complication of diarrhea during hospitalization; failure to provide informed consent	2 × 10 ¹² ; CFU/day	ORF	7 d or until diarrhea stopped	Bacterial diarrhea excluded; Rotavirus-induced diarrhea 56.06%
Canani <i>et al</i> ^[25] , 2007	RCT; 6 centers; Duration: 12 mo	3-36 mo	Italy	Outpatients	100/92	> 2 loose or liquid stools/day for <48 h	Malnutrition; severe dehydration; coexisting acute systemic illness; immunodeficiency; underlying severe chronic disease; cystic fibrosis; food allergy or other chronic GI diseases; use of probiotics in the previous 3 wk; antibiotics or any other antidiarrheal medication in the previous 3 wk; poor compliance	12 × 10 ⁹ ; CFU/day	No details given	5 d	Stool culture in only a few participants and no data presented
Costa <i>et al</i> ^[27] , 2003	RCT; 1 center	Boys, 1-24 mo	Brazil	Inpatients	61/63	Acute diarrhea (3 or more watery or loose stools per 24 h during at least one 24-h period in the 72 h before admission) with moderate or severe dehydration after correction with rapid IV fluids	Systemic infections requiring antibiotics; severe malnutrition (weight for age < 65% of NCHS standards; bloody diarrhea	10 ¹⁰ ; CFU/day	Inulin 320 mg/day	Unclear	Rotavirus-induced diarrhea 50%; Bloody diarrhea excluded
Czerwionka-Szaflarska <i>et al</i> ^[28] , 2009	RCT; 1 center	Unclear	Poland	Inpatients	50/50	Infants and children with acute infectious diarrhea and failed oral rehydration	Bloody stools; coexisting disease that may influence the course of diarrhea	50 ml/kg/day	Unclear	Unclear	Bloody diarrhea excluded; Rotavirus-induced diarrhea 58%

Schnadower <i>et al</i> ^[8] , 2018	RCT	3-48 mo	United States	University-affiliated PED	483/488	≥ 3 watery stools per day, with or without vomiting, for fewer than 7 d	Pancreatitis, biliious emesis, or hematochezia; a known allergy to <i>L. rhamnosus</i> CG or to microcrystalline cellulose or a known allergy to erythromycin, clindamycin, and beta-lactam antibiotic agents; caregiver did not speak English or Spanish; children receiving antibiotics	1 × 10 ¹⁰ ; CFU twice daily	Matching placebo	5 d	Norovirus GI or GII 19.6%; Rotavirus 17.7%; Adenovirus 9.1%; Clostridium difficile 7.4%; Shigella 5.0%
Guandalini <i>et al</i> ^[29] , 2000	RCT; multicenter; Duration: 1 yr	1-36 mo	Listed as follows	Inpatients and outpatients	147/140	Infants and children with > 4 liquid or	Previous probiotic usage; untreated underlying chronic small bowel disease; inflammatory bowel disease; any underlying chronic disease or immunosuppressive disease or treatment	≥ 10 × 10 ⁹ ; CFU/250 mL/day with ORF	ORF	As tolerated for 4-6 h, then ad libitum	Rotavirus 35%; Bacteria 24%; Parasites 4.5%; No pathogens 34.5%; Bloody diarrhea 8.7%
Guarino <i>et al</i> ^[30] , 1997	RCT; 1 center Duration: 3 mo	3-36 mo	Italy	Outpatients	52/48	semiliquid stools/day for 1 to 5 d Infants and children with ≥ 3 watery stools/day for < 48 h	Antibiotic treatment in the last 3 wk, breastfeeding: a weight: height ratio < the 5th percentile	6 × 10 ⁹ ; CFU/day with ORF	ORF	≤ 5 d	Rotavirus-induced diarrhea 61%
Isolauri <i>et al</i> ^[31] , 1994	RCT; 1 center Duration: not stated	≤ 36 mo	Finland	Inpatients	21/21	Infants and children with > 3 watery stools/day for < 7 d and stools positive for rotavirus; average dehydration of approximately 5% in both groups	Not stated	2 × 10 ¹⁰ ; CFU/day	No probiotic	5 d	Rotavirus-induced diarrhea 100%
Jasinski <i>et al</i> ^[32] , 2002	RCT; 12 centers; Duration: not stated	1-36 mo	Africa	Inpatients and outpatients	45/52	> 3 watery stools in 12 h or 1 liquid or semiliquid stool with mucus, pus, or blood; < 5 d	Antibiotic or probiotic use in the last 5 d; chronic diseases of the small or large intestine; immunosuppression; phenylketonuria	ORS + LGG 10 ¹⁰ CFU/day	ORS with no LGG	Unclear	Bacterial pathogens 68%; Rotavirus 40.0%; parasites 6%; No pathogens identified: probiotic group 25%
Misra <i>et al</i> ^[33] , 2009	RCT; 1 center; Duration: not stated	≤ 36 mo	Egypt Europe America India	Inpatients	105/105	> 3 stools per day (watery or assuming the shape of the container)	Parents refused consent; children living outside the municipal area; bloody diarrhea; severe dehydration; shock, inability to take and retain oral foods; suspected systemic infection	1 × 10 ⁶⁻⁹ CFU/day	Crystalline micro cellulose	Unclear	Rotavirus 25.6%; Bloody diarrhea excluded; White blood cells in stools 14.3%; Bacterial diarrhea 4.7%
Nixon <i>et al</i> ^[34] , 2012	RCT	6-72 mo	United States	PED	77/78	More than 2 loose stools in the last 24 h	Risk factors for non-viral diarrhea (prolonged diarrhea lasting more than 7 d, gross blood, antibiotic exposure, or inflammatory bowel disease); immune compromise; risk factors for probiotic-associated systemic illness or an allergy to milk products	LGG powder twice daily	Inulin	5 d	Unclear
Pant <i>et al</i> ^[35] , 1996	RCT; 1 center; Duration: 6 wk	1-24 mo	Thailand	Inpatients	20/19	Infants and children with > 3 watery stools in last 24 h and diarrhea for < 14 d	Exclusive breastfeeding; septicemia	10 ⁹⁻¹⁰ CFU twice daily	Placebo	2 d	Bloody stools 33.3%;

Raza <i>et al</i> ^[34] , 1995	RCT; 1 center Duration: 2 mo	1-24 mo	Pakistan	Inpatients	21/19	Undernourished infants and children with > 3 watery stools in the last 24 h for < 14 d and at least moderate dehydration	Severe malnutrition; septicemia	$2 \times 10^{11.12}$ CFU/day	Placebo	2 d	Rotavirus 17.9%; Astrovirus 2.5% Bloody diarrhea
Ritchie <i>et al</i> ^[37] , 2010	RCT; 1 center; Duration: 21 mo	4-24 mo	Australia	Unclear	33/31	Aboriginal children with acute diarrhea defined as ≥ 3 loose stools during 24 h before presentation for < 7 d and able to tolerate ORF	Oxygen required during the study period; chronic cardiac, renal, or respiratory disease; previous gastrointestinal surgery; proven sucrose intolerance; suspected on known immunodeficiency; probiotic use before enrollment; younger than 4 mo of age	$> 15 \times 10^9$ CFU/day	Identical placebo	3 d	Bacterial pathogens 12.5%; Rotavirus 8.5%; Parasites 6%
Shornikova <i>et al</i> ^[38] , 1997	RCT; 1 center; Duration: 1 yr	1-36 mo	Russia	Inpatients	59/64	≥ 1 watery stool in the last 24 h and diarrhea for < 5 d	Not stated	10^{10} CFU/day	Placebo	5 d	Rotavirus 27.4%; Bacterial diarrhea 21%
Sindhu <i>et al</i> ^[39] , 2014	RCT	6-60 mo	India	Unclear	65/59	Diarrhea was defined as ≥ 3 loose watery stools within a 24-h period	Coinfections (the presence of both rotavirus and Cryptosporidium); severe malnutrition; probiotic consumption in the preceding month; allergy to probiotics; acute abdomen or colitis	10^{10} CFU and 170 mg of microcrystalline /day cellulose	170 mg of cellulose	4 wk	Rotavirus 52.4%; Cryptosporidium species 47.6%
Sunny <i>et al</i> ^[40] , 2014	Open-label; RCT	6-60 mo	India	OPD or PED	100/100	Passage of three or more loose stools in the last 24 h	Severe malnutrition; dysentery; clinical evidence of coexisting acute systemic illnesses; clinical evidence of chronic disease; probiotic use in the preceding three weeks; antibiotic use	1×10^{10} CFU per day	ORS and zinc 20 mg/d	5 d	Rotavirus 24.1%

The study of Guandalini *et al*^[29] was conducted in Poland, Pakistan, Egypt, Croatia, Italy, Slovenia, Netherlands, Greece, Israel, the United Kingdom, and Portugal. RCT: Randomized controlled trial; PED: Pediatric emergency department; OPD: Outpatient department; LGG: Lactobacillus rhamnosus GG.

matched group according to 15 RCTs submitted to meta-analysis, which included 3721 participants [MD -24.02 h, 95%CI (-36.58, -11.45)] (Figure 1A). Significantly heterogeneous results were detected among the included trials ($I^2 = 98\%$). Our data support the results of the prior meta-analyses^[4] indicating that LGG treatment reduced participants' duration of diarrhea.

Subgroup analyses were conducted based on clinical features such as age, geographical location, treatment time, outpatient or inpatient settings, the time of enrollment, and literature quality scores. Differences in methodological quality could not explain the statistically significant heterogeneity (Supporting information Figure S6). Regression analysis between the duration of diarrhea and LGG dose revealed that different doses of LGG contributed to the heterogeneity ($P = 0.009$, adjusted R-squared = 40.21%), suggesting that subgroups according to a high or low dose of LGG should be assessed. A reduced duration of diarrhea was noted in the studies applying $> 10^{10}$ CFU/day of LGG [MD -22.56 h, 95%CI (-36.41, -8.72)] (Figure 1A). In contrast, although only three studies used lower dosages, no statistically significant differences were detected in the groups receiving lower dosages [MD -30.95 h, 95%CI (-83.28, -21.39)] (Figure 1A). A reduced duration of diarrhea was supported in the studies with participants who received LGG treatment before the second day of diarrhea symptoms [MD -1.58 h, 95%CI (-3.05, -0.11)] and during the second to third days of diarrhea symptoms [MD -15.83 h, 95%CI (-20.06, -10.98)] (Figure 1B). However, Ritchie *et al.*^[37] enrolled participants with diarrhea for more than 3 d, and no statistically significant differences were found in the duration of diarrhea [MD 1.2 h, 95%CI (-21.42, 23.82)] (Figure 1B). A reduced diarrhea duration was reported in studies performed in both Asia and Europe [MD -24.42 h, 95%CI (-47.10, -1.82); MD -32.02 h, 95%CI (-49.26, -14.79), respectively]. Paradoxically, the reduction in the diarrhea duration in other regions was not statistically significant [MD -9.35 h, 95%CI (-20.77, 2.07)] (Figure 1C). In the etiological analysis, the effectiveness of LGG was clearly demonstrated in rotavirus-induced diarrhea cases [seven RCTs; MD -31.05 h, 95%CI (-50.31, -11.80)] (Figure 2). Analysis with the studies carried out in the 1990s and 2000s revealed a clear reduction in the diarrhea duration [MD -36.32 h, 95%CI (-62.20, -10.45); MD -29.40 h, 95%CI (-50.56, -8.25), respectively] (Supporting information Figure S7). In contrast, no reduction in the diarrhea duration was observed in the analysis with studies carried out in the 2010s [MD -3.43 h, 95%CI (-13.25, 6.39)] (Supporting information Figure S7). No studies evaluated the effectiveness of LGG in children vaccinated against rotavirus.

Diarrhea ≥ 3 d

A meta-analysis of four RCTs was performed using a fixed-effects model. The risk of experiencing diarrhea for 3 or more days was reduced when patients received LGG [odds ratio (OR) 0.54, 95%CI (0.38, 0.77)] (Figure 3A).

Diarrhea ≥ 4 d



Three studies were pooled ($n = 479$) and showed a reduction in the risk of diarrhea lasting for 4 or more days for participants treated with LGG [OR 0.58, 95%CI (0.4, 0.84)] (Figure 3B).

Stool number and consistency

Stool number and consistency were evaluated in most trials. Eight trials reported the mean number of stools in one day during diarrhea episodes. A notable decrease in the stool number per day was noted in the LGG group [MD -0.9, 95%CI (-1.56, -0.23)] (Figure 4A). However, a significantly reduced stool number was observed in the high-dose LGG groups receiving no less than 10^{10} CFU/day [MD -1.08, 95%CI (-1.87, -0.28)], while the lower-dose groups showed no significant reduction [MD -0.25 d, 95%CI (-1.43, 0.93)] (Figure 4A). After the intervention, stool frequency was evaluated on days 2 and 3. Seven trials provided data on day 2, and the overall effect did not differ between the two groups [MD -0.46, 95%CI (-1.06, 0.15)] (Figure 4B). In addition, similar frequencies were observed in the two groups on day 3, with no differences between them [MD 0.34, 95%CI (-0.29, 0.97)] (Figure 4C). Three trials calculated the mean time to improvement in stool consistency, and an obvious reduction was reported [MD -5.65, 95%CI (-7.49, -3.80)] (Figure 4D).

Hospital stay duration

A total of 1823 participants from six RCTs were analyzed. Due to statistically significant heterogeneity, a random-effects model was used, which revealed a significant reduction in the hospital stay duration in the two groups [MD -39.16 h, 95%CI (-72.24, -6.07)] (Figure 5A). A reduction in the hospital stay duration was found in rotavirus-positive children [MD -21.12 h, 95%CI (-26.96, -15.28)] (Figure 5B).

A**B**




Figure 1 *Lactobacillus* GG vs control with regard to the duration of diarrhea (hours). A: High dose and low dose; B: The duration of diarrhea before *Lactobacillus rhamnosus* GG participants' enrollment: ≤ 2 d (>1 d), ≤ 3 d (>2 d), and ≤ 4 d (>3 d); C: Geography of the clinical trials: Asia, Europe, and other continents. LGG: *Lactobacillus rhamnosus* GG; CI: Confidence interval; SD: Standard deviation.

Vomiting

Vomiting in different trials was reported as the number of participants with vomiting [number (%)] or as the duration of vomiting (hours). Compared with the placebo group, no difference in the risk of vomiting was reported in the experimental group [OR 1.11, 95%CI (0.59, 2.12)] (Figure 6A). Furthermore, no reduction in the duration of vomiting was noted with LGG treatment [MD -2.02 h, 95%CI (-4.24, 0.21)] (Figure 6B).

Adverse effects

Probiotics have been proposed to be well-tolerated and safe therapeutic agents. Most authors did not report adverse effects. Raza *et al*^[36] reported one case of myoclonic jerks in their trial. Lower rates of respiratory infection, wheezing, and even vulvar abscess were noted in Schnadower's trial^[8,39], but these effects were not thought to be correlated with LGG use^[40]. Aggarwal *et al*^[40] reported no adverse effects, and a meta-analysis performed in 2013 showed comparable rates of adverse effects among study groups^[13]. In our study, eight studies effectively evaluated the safety of LGG. Adverse effects were reported on a coded reporting form or during daily telephone calls^[26,34]. In Schnadower's study, the caregivers completed a daily diary that was collected by telephone or through email^[8]. However, the reporting methods were unclear in five articles^[24,37,39-41]. In general, no adverse effects or similar rates of side effects were documented in the LGG and placebo groups.

Risk of bias in the included studies

The risk of bias in 18 articles was assessed according to the Cochrane Handbook for Systematic Reviews of Interventions. One trial employed alternating group allocation, and the random sequence generation method was not reported in five trials. Other RCTs provided detailed randomization methods, which mainly included computer-generated strategies, resulting in a low risk of selection bias. Allocation concealment was not applied in two trials and was not mentioned in seven. Nine trials used the sealed envelope technique for allocation concealment. Double blinding was strictly executed in 12 trials, while four trials allowed openness to patients or doctors, and




Figure 2 *Lactobacillus* GG vs control with regard to mean duration of diarrhea (hours) in children with rotavirus diarrhea. LGG: *Lactobacillus rhamnosus* GG; CI: Confidence interval; SD: Standard deviation.

two trials did not report a detailed blinding method. For detection bias, investigators were blinded to the group assignments in ten trials, while blinding assessments were not performed in three trials. Most trials provided complete data with a loss to follow-up rate less than 10%, although one trial had an unknown risk of incomplete outcome data, reflecting a low risk of attrition bias (Supporting information Figure S3).

Publication bias

According to Egger's^[18] regression asymmetry test, no small sample or publication bias was found in a funnel plot [$P = 0.10$, 95%CI (-11.33, -1.14)] (Supporting information Figure S8).

DISCUSSION

Findings and agreement or disagreement with other studies

Nineteen trials comparing a control group with an experimental group treated with LGG were identified in this meta-analysis. In summary, the analysis revealed that treatment with LGG reduced both the duration of diarrhea and the hospital stay duration, especially in specific patient subsets. A striking finding was the time to improvement in stool consistency, which more investigators have confirmed since 2010^[8,34,40]. In the whole range of diarrhea cases, the management of stools with this probiotic strain showed a modest beneficial effect on the number of stools per day and the time to improvement in stool consistency. However, no reduction in stool frequency was observed on days 2 and 3. Compared with the placebo group, the risk of diarrhea lasting more than 3 and 4 d was reduced by LGG administration. In both groups, similar rates of vomiting and adverse effects were observed.

Evidence from RCTs confirmed the beneficial effect of LGG on rotavirus-induced diarrhea^[42]. In addition to interference with viral replication, most recent studies have shown that LGG prevented injuries to the epithelium and ameliorated rotavirus-induced diarrhea by modulating immune cells, such as dendritic cells and inflammatory cytokines^[43,44]. The marked statistical difference in the diarrhea duration with a higher dosage of probiotics reflected greater effectiveness, which confirmed the dose dependence of dendritic cell activation. Treatment efficacy was related to the dose of LGG^[45]. As confirmed in the study of Szajewska *et al*^[13] in 2013, the importance of a daily LGG dose is high, and a dosage of 10^{10} CFU/day is needed for a positive effect. The statistical heterogeneity between studies can be explained by the timing of the LGG intervention, which was directly correlated with indicators such as the duration of diarrhea before study enrollment. Although the heterogeneity persisted in the subgroup with the shortest duration of diarrhea before study enrollment, probiotics should be applied early in the course of disease. Moreover, symptoms are usually mild at the early stage. Differences in prominent pathogens, sanitation conditions, and common comorbidities lead to dissimilarities between various study locations. Due to differences in the treatment effect among regions, the implications for clinical practice should be evaluated. The nutrient intake and dietary structure of humans have continuously changed in recent decades, which may have caused the reduced effectiveness of LGG reflected in the results of the trials conducted in the 2010s.

Probiotics manipulate and restore the gut microbiota, thus benefitting the




Figure 3 *Lactobacillus* GG vs control with regard to the presence of diarrhea. A: Diarrhea lasting > 3 d; B: Diarrhea lasting > 4 d. LGG: *Lactobacillus rhamnosus* GG; CI: Confidence interval.

prevention of diarrhea. Various therapeutic interventions designed to alter the microbiota range from probiotic administration to fecal microbiota transplantation^[46,47]. However, due to the limited number of included studies and the self-limiting nature of disease, strategies should also be discussed in detail. Vomiting was reported as an adverse event in numerous studies^[48,49], and it is one of the most common symptoms associated with diarrhea^[50,51]. Additionally, less frequent clinical symptoms were observed in the probiotic groups^[4], although our meta-analysis showed no improvement in the risk or duration of vomiting.

Safety

The safety of probiotic supplementation is generally certain. Nevertheless, pathologies correlated with the use of probiotic products to treat gastrointestinal disorders have been identified, such as endocarditis, sepsis, and bacteremia^[52-54]. Unfortunately, the most prevalent strain implicated in the adverse effects was *Lactobacillus rhamnosus*. Conversely, most authors in our analysis did not report adverse effects or the adverse effects were not thought to be correlated with LGG treatment. In addition to the interventions, the primary illness contributed the most to the participant drop-out rate. A higher frequency of negative effects attributed to probiotics was found in catheterized (82.5%) and immunosuppressed (66%) participants^[55]. Further safety evaluations of probiotics are necessary in the clinical setting, especially for susceptible individuals, such as those with immunodeficiency, immunosuppression, or malnourishment.

Application prospects

Preventing or correcting dehydration through treatment with zinc or 0.9% saline solution is the main approach used for diarrhea management^[56]. However, during diarrhea episodes, infectious symptoms are not fully alleviated and the gut microbiota is not restored by rehydration measures^[57]. Probiotics were investigated as therapeutic agents for diarrhea. The mechanisms by which probiotics alleviate diarrhea are described below. Host defenses are reinforced by enhanced antimicrobial peptide secretion. Probiotics prevent disruption of gut barrier integrity and stimulate the expression of junctional adhesion and tight junction molecules^[58-61]. They produce short-chain fatty acids and induce the production of IgA to resist infections^[62-64]. In epithelial cells and mucin, probiotics compete for binding sites to arrest pathogen colonization^[65]. Probiotics can specifically and nonspecifically interfere with the viral cycle, thus impeding the progression of rotavirus-induced diarrhea^[66-68]. The prevalence of diarrhea is seasonal, and almost all cases of rotavirus-induced diarrhea




Figure 4 *Lactobacillus* GG vs control with regard to stool number and consistency. A: The average stool number per day (high dose and low dose); B: Stool frequency on day 2; C: Stool frequency on day 3; D: The mean time to improvement in stool consistency. LGG: *Lactobacillus rhamnosus* GG; CI: Confidence interval; SD: Standard deviation.




Figure 5 *Lactobacillus* GG vs control. A: The duration of hospital stay (hours); B: The hospital stay duration of rotavirus-positive children (hours). LGG: *Lactobacillus rhamnosus* GG; CI: Confidence interval; SD: Standard deviation.

occur from January to May in Russia^[38]. By contrast, in regions where rotavirus is not prevalent, bacterial diarrhea commonly occurs from June to October^[38]. Influenza seasons, dietary habits, and antibiotic use must be considered when evaluating heterogeneity in further studies. The efficacy of probiotic treatment was altered based on host and environmental factors^[12]. Overall, our study supported the previous systematic reviews which concluded that LGG is an effective treatment for children with acute diarrhea.

Conclusions and limitations

Although most studies have suggested that LGG is efficacious, limited identification of pathogens, small sample sizes, varying therapeutic strategies, and methodological limitations such as articles without a strictly blinded design, including a lack of a standard clinical parameter format, weakened the conclusions and precluded further analyses across studies^[69]. For example, Czerwionka-Szaflarska *et al*^[28] did not specifically define the treatment applied, although a significantly reduced duration of diarrhea was detected. Salazar-Lindo *et al*^[41] partially depicted the duration of diarrhea in children with or without LGG treatment. Although factors varied in the trials, according to the same criterion for both groups, no evidence suggests that a poor study design leads to overestimation of probiotic efficacy^[4]. Appropriate subgroups, such as those stratified by etiology and nutritional status, are indispensable. In 2016, approximately 8.4% of children (480000) presenting with diarrhea ultimately died due to the condition worldwide (<https://data.unicef.org/topic/child-health/diarrhoeal-disease/>). Assessments of the availability of vaccines, the applicability of probiotics, and the effectiveness of current treatments under severe conditions and cost-effect analyses must be performed to optimize therapeutic strategies for acute diarrhea management in children.

In summary, the following conclusions were cautiously established: LGG reduces the duration of diarrhea, particularly in patients with rotavirus-positive diarrhea receiving a dosage no less than 10^{10} CFU per day and in patients treated at the early stage. In addition, studies conducted in Asia and Europe showed greater treatment efficacy. The therapeutic effect of LGG supplementation on the stool number per day and hospital stay duration associated with rotavirus-induced diarrhea is high.




Figure 6 *Lactobacillus* GG vs control with regard to vomiting. A: The number of participants with vomiting [number (%)] B: The duration of vomiting (hours). LGG: *Lactobacillus rhamnosus* GG; CI: Confidence interval; SD: Standard deviation.

ARTICLE HIGHLIGHTS

Research background

Diarrhea is a major infectious cause of childhood morbidity and mortality worldwide. Preventing or correcting dehydration through treatment with zinc or 0.9% saline solution is the main approach for diarrhea management; however, during diarrhea episodes, infectious symptoms are not fully alleviated by rehydration measures. Probiotics restore the gut microbiota and have been reported to reduce the duration of diarrhea.

Research motivation

Although previous studies have reported that *Lactobacillus rhamnosus* GG (LGG) is an effective therapeutic agent for acute diarrhea in children, a recent large, high-quality RCT found no adequate evidence of a beneficial effect of LGG treatment.

Research objectives

To evaluate the efficacy of LGG in treating acute diarrhea in children and provide some reference for future trials of treatments for diarrhea.

Research methods

The EMBASE, MEDLINE, PubMed, Web of Science databases, and the Cochrane Central Register of Controlled Trials were searched up to April 2019 for meta-analyses and randomized controlled trials (RCTs). Cochrane Review Manager was used to analyze the relevant data and primary outcomes, including the duration of diarrhea and diarrhea lasting ≥ 3 and ≥ 4 d. Secondary outcomes included the hospital stay duration, stool frequency, and improvement in stool consistency and vomiting.

Research results

The systematic review identified 19 RCTs that met the inclusion criteria and indicated that compared with the control group, LGG administration notably reduced the diarrhea duration [mean difference (MD) -24.02 h, 95% confidence interval (CI) (-36.58, -11.45)]. Greater reductions were detected at a high dose of $\geq 10^{10}$ CFU per day [MD -22.56 h, 95%CI (-36.41, -8.72)] and in LGG participants with diarrhea for less than 3 days at study enrollment [MD -15.83 h, 95%CI (-20.68, -10.98)]. The study locations contributed to differences in the reduction in the diarrhea duration in Asia and Europe [MD -24.42 h, 95%CI (-47.01, -1.82); MD -32.02 h, 95%CI (-49.26, -14.79), respectively]. High-dose LGG treatment was confirmed to effectively reduce the duration of rotavirus-induced diarrhea [MD -31.05 h, 95%CI (-50.31, -11.80)] and stool number [MD -1.08, 95%CI (-1.87, -0.28)].

Research conclusions

The following conclusions were cautiously established: compared to control children, children

who received a course of LGG had better outcomes, including a markedly reduced duration of diarrhea, especially those with rotavirus-positive diarrhea, those who received no less than 10^{10} CFU per day, and those treated at the early stage. Furthermore, studies conducted in Asia and Europe reported greater treatment efficacy. The therapeutic effect of LGG supplementation on the stool number per day and hospital stay duration associated with rotavirus-induced diarrhea was high.

Research perspectives

Our study found better outcomes among children with acute diarrhea who were treated by LGG supplementation. Limited identification of pathogens, small sample sizes, and a lack of a standard clinical parameter format precluded further analyses across studies, thus weakening the evidence required to guide clinical practice. Investigations are required to assess the cost-effectiveness of treating diarrhea with probiotics.

REFERENCES

- 1 Walker CLF, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, O'Brien KL, Campbell H, Black RE. Global burden of childhood pneumonia and diarrhoea. *Lancet* 2013; **381**: 1405-1416 [PMID: 23582727 DOI: 10.1016/S0140-6736(13)60222-6]
- 2 do Carmo MS, Santos CID, Araújo MC, Girón JA, Fernandes ES, Monteiro-Neto V. Probiotics, mechanisms of action, and clinical perspectives for diarrhea management in children. *Food Funct* 2018; **9**: 5074-5095 [PMID: 30183037 DOI: 10.1039/c8fo00376a]
- 3 Savarino SJ, Bourgeois AL. Diarrhoeal disease: Current concepts and future challenges. *Epidemiology of diarrhoeal diseases in developed countries. Trans R Soc Trop Med Hyg* 1993; **87** Suppl 3: 7-11 [PMID: 8108853 DOI: 10.1016/0035-9203(93)90529-y]
- 4 Allen SJ, Martinez EG, Gregorio GV, Dans LF. Probiotics for treating acute infectious diarrhoea. *Cochrane Database Syst Rev* 2010; CD003048 [PMID: 21069673 DOI: 10.1002/14651858.CD003048.pub3]
- 5 Binder HJ, Brown I, Ramakrishna BS, Young GP. Oral rehydration therapy in the second decade of the twenty-first century. *Curr Gastroenterol Rep* 2014; **16**: 376 [PMID: 24562469 DOI: 10.1007/s11894-014-0376-2]
- 6 Bryce J, Tetteri N, Victora CG, Mason E, Daelmans B, Bhutta ZA, Bustreo F, Songane F, Salama P, Wardlaw T. Countdown to 2015: Tracking intervention coverage for child survival. *Lancet* 2006; **368**: 1067-1076 [PMID: 16997661 DOI: 10.1016/S0140-6736(06)69339-2]
- 7 Pérez-Gaxiola G, Cuello-García CA, Florez ID, Pérez-Pico VM. Smectite for acute infectious diarrhoea in children. *Cochrane Database Syst Rev* 2018; **4**: CD011526 [PMID: 29693719 DOI: 10.1002/14651858.CD011526.pub2]
- 8 Schnadower D, Tarr PI, Casper TC, Gorelick MH, Dean JM, O'Connell KJ, Mahajan P, Levine AC, Bhatt SR, Roskind CG, Powell EC, Rogers AJ, Vance C, Sapient RE, Olsen CS, Metheny M, Dickey VP, Hall-Moore C, Freedman SB. Lactobacillus rhamnosus GG versus Placebo for Acute Gastroenteritis in Children. *N Engl J Med* 2018; **379**: 2002-2014 [PMID: 30462938 DOI: 10.1056/NEJMoa1802598]
- 9 Guarner F, Schaafsma GJ. Probiotics. *Int J Food Microbiol* 1998; **39**: 237-238 [PMID: 9553803 DOI: 10.1016/S0168-1605(97)00136-0]
- 10 Lomax AR, Calder PC. Probiotics, immune function, infection and inflammation: A review of the evidence from studies conducted in humans. *Curr Pharm Des* 2009; **15**: 1428-1518 [PMID: 19442167 DOI: 10.2174/138161209788168155]
- 11 Szajewska H, Mrukowicz JZ. Probiotics in the treatment and prevention of acute infectious diarrhea in infants and children: A systematic review of published randomized, double-blind, placebo-controlled trials. *J Pediatr Gastroenterol Nutr* 2001; **33** Suppl 2: S17-S25 [PMID: 11698781 DOI: 10.1097/00005176-200110002-00004]
- 12 Wolvers D, Antoine JM, Myllyluoma E, Schrezenmeir J, Szajewska H, Rijkers GT. Guidance for substantiating the evidence for beneficial effects of probiotics: Prevention and management of infections by probiotics. *J Nutr* 2010; **140**: 698S-712S [PMID: 20107143 DOI: 10.3945/jn.109.113753]
- 13 Szajewska H, Skórka A, Ruszczyński M, Gieruszczak-Białek D. Meta-analysis: Lactobacillus GG for treating acute gastroenteritis in children—updated analysis of randomised controlled trials. *Aliment Pharmacol Ther* 2013; **38**: 467-476 [PMID: 23841880 DOI: 10.1111/apt.12403]
- 14 Szajewska H, Guarino A, Hojsak I, Indrio F, Kolacek S, Shamir R, Vandeplass Y, Weizman Z; European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. Use of probiotics for management of acute gastroenteritis: A position paper by the ESPGHAN Working Group for Probiotics and Prebiotics. *J Pediatr Gastroenterol Nutr* 2014; **58**: 531-539 [PMID: 24614141 DOI: 10.1097/MPG.0000000000000320]
- 15 Guarino A, Guandalini S, Lo Vecchio A. Probiotics for Prevention and Treatment of Diarrhea. *J Clin Gastroenterol* 2015; **49** Suppl 1: S37-S45 [PMID: 26447963 DOI: 10.1097/MCG.0000000000000349]
- 16 Cipriani A, Furukawa TA, Salanti G, Chaimani A, Atkinson LZ, Ogawa Y, Leucht S, Ruhe HG, Turner EH, Higgins JPT, Egger M, Takeshima N, Hayasaka Y, Imai H, Shinohara K, Tajika A, Ioannidis JPA, Geddes JR. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *Lancet* 2018; **391**: 1357-1366 [PMID: 29477251 DOI: 10.1016/S0140-6736(17)32802-7]
- 17 Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: Explanation and elaboration. *BMJ* 2009; **339**: b2700 [PMID: 19622552 DOI: 10.1136/bmj.b2700]
- 18 Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629-634 [PMID: 9310563 DOI: 10.1136/bmj.315.7109.629]
- 19 Johnston BC, Goldenberg JZ, Vandvik PO, Sun X, Guyatt GH. Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database Syst Rev* 2011; CD004827 [PMID: 22071814 DOI: 10.1002/14651858.CD004827.pub3]
- 20 Johnston BC, Ma SS, Goldenberg JZ, Thorlund K, Vandvik PO, Loeb M, Guyatt GH. Probiotics for the

- prevention of Clostridium difficile-associated diarrhea: A systematic review and meta-analysis. *Ann Intern Med* 2012; **157**: 878-888 [PMID: 23362517 DOI: 10.7326/0003-4819-157-12-201212180-00563]
- 21 **Cunliffe NA**, Kilgore PE, Bresee JS, Steele AD, Luo N, Hart CA, Glass RI. Epidemiology of rotavirus diarrhoea in Africa: A review to assess the need for rotavirus immunization. *Bull World Health Organ* 1998; **76**: 525-537 [PMID: 9868844 DOI: 10.1146/annurev.publhealth.19.1.527]
 - 22 **Simakachorn N**, Pichaiat V, Rithipornpaisarn P, Kongkaew C, Tongpradit P, Varavithya W. Clinical evaluation of the addition of lyophilized, heat-killed Lactobacillus acidophilus LB to oral rehydration therapy in the treatment of acute diarrhea in children. *J Pediatr Gastroenterol Nutr* 2000; **30**: 68-72 [PMID: 10630442 DOI: 10.1097/00005176-200001000-00020]
 - 23 **Szajewska H**, Skórka A, Ruszczyński M, Gieruszczak-Bialek D. Meta-analysis: Lactobacillus GG for treating acute diarrhoea in children. *Aliment Pharmacol Ther* 2007; **25**: 871-881 [PMID: 17402990 DOI: 10.1111/j.1365-2036.2007.03282.x]
 - 24 **Basu S**, Paul DK, Ganguly S, Chatterjee M, Chandra PK. Efficacy of high-dose Lactobacillus rhamnosus GG in controlling acute watery diarrhea in Indian children: A randomized controlled trial. *J Clin Gastroenterol* 2009; **43**: 208-213 [PMID: 18813028 DOI: 10.1097/MCG.0b013e31815a5780]
 - 25 **Basu S**, Chatterjee M, Ganguly S, Chandra PK. Efficacy of Lactobacillus rhamnosus GG in acute watery diarrhoea of Indian children: A randomised controlled trial. *J Paediatr Child Health* 2007; **43**: 837-842 [PMID: 17803667 DOI: 10.1111/j.1440-1754.2007.01201.x]
 - 26 **Canani RB**, Cirillo P, Terrin G, Cesarano L, Spagnuolo MI, De Vincenzo A, Albano F, Passariello A, De Marco G, Manguso F, Guarino A. Probiotics for treatment of acute diarrhoea in children: Randomised clinical trial of five different preparations. *BMJ* 2007; **335**: 340 [PMID: 17690340 DOI: 10.1136/bmj.39272.581736.55]
 - 27 **Costa-Ribeiro H**, Ribeiro TC, Mattos AP, Valois SS, Neri DA, Almeida P, Cerqueira CM, Ramos E, Young RJ, Vanderhoof JA. Limitations of probiotic therapy in acute, severe dehydrating diarrhea. *J Pediatr Gastroenterol Nutr* 2003; **36**: 112-115 [PMID: 12500005 DOI: 10.1097/00005176-200301000-00021]
 - 28 **Czerwionka-Szaflarska M**, Murawska S, Swincow G. Evaluation of influence of oral treatment with probiotic and/or oral rehydration solution on course of acute diarrhoea in children. *Przegl Gastroenterol* 2009; **4**: 166-172
 - 29 **Guandalini S**, Pensabene L, Zikri MA, Dias JA, Casali LG, Hoekstra H, Kolacek S, Massar K, Micetic-Turk D, Papadopoulou A, de Sousa JS, Sandhu B, Szajewska H, Weizman Z. Lactobacillus GG administered in oral rehydration solution to children with acute diarrhea: A multicenter European trial. *J Pediatr Gastroenterol Nutr* 2000; **30**: 54-60 [PMID: 10630440 DOI: 10.1097/00005176-200001000-00018]
 - 30 **Guarino A**, Canani RB, Spagnuolo MI, Albano F, Di Benedetto L. Oral bacterial therapy reduces the duration of symptoms and of viral excretion in children with mild diarrhea. *J Pediatr Gastroenterol Nutr* 1997; **25**: 516-519 [PMID: 9360205 DOI: 10.1097/00005176-199711000-00005]
 - 31 **Isolauri E**, Kaila M, Mykkänen H, Ling WH, Salminen S. Oral bacteriotherapy for viral gastroenteritis. *Dig Dis Sci* 1994; **39**: 2595-2600 [PMID: 7995184 DOI: 10.1007/BF02087695]
 - 32 **Jasinski C TM**, Tanzi MN, Schelotto F, Varela G, Zanetta E, Acuña A, and Arenas C, del Pilar Gadea M, Sirok A, Betancor L, Grotiuz G, Sandin D, Combol A, Xavier B, Vignoli R, Nairac A. Efficacy of Lactobacillus GG in oral rehydration solution. *Pediatr* 2002; **22**: 231-243
 - 33 **Misra S**, Sabui TK, Pal NK. A randomized controlled trial to evaluate the efficacy of lactobacillus GG in infantile diarrhea. *J Pediatr* 2009; **155**: 129-132 [PMID: 19559297 DOI: 10.1016/j.jpeds.2009.01.060]
 - 34 **Nixon AF**, Cunningham SJ, Cohen HW, Crain EF. The effect of Lactobacillus GG on acute diarrheal illness in the pediatric emergency department. *Pediatr Emerg Care* 2012; **28**: 1048-1051 [PMID: 23023475 DOI: 10.1097/PEC.0b013e31826cad9f]
 - 35 **Pant AR**, Graham SM, Allen SJ, Harikul S, Sabchareon A, Cuevas L, Hart CA. Lactobacillus GG and acute diarrhoea in young children in the tropics. *J Trop Pediatr* 1996; **42**: 162-165 [PMID: 8699584 DOI: 10.1093/tropej/42.3.162]
 - 36 **Raza S**, Graham SM, Allen SJ, Sultana S, Cuevas L, Hart CA. Lactobacillus GG promotes recovery from acute nonbloody diarrhea in Pakistan. *Pediatr Infect Dis J* 1995; **14**: 107-111 [PMID: 7746691 DOI: 10.1097/00006454-199502000-00005]
 - 37 **Ritchie BK**, Brewster DR, Tran CD, Davidson GP, McNeil Y, Butler RN. Efficacy of Lactobacillus GG in aboriginal children with acute diarrhoeal disease: A randomised clinical trial. *J Pediatr Gastroenterol Nutr* 2010; **50**: 619-624 [PMID: 20400916 DOI: 10.1097/MPG.0b013e3181bbf53d]
 - 38 **Shornikova AV**, Isolauri E, Burkanova L, Lukovnikova S, Vesikari T. A trial in the Karelian Republic of oral rehydration and Lactobacillus GG for treatment of acute diarrhoea. *Acta Paediatr* 1997; **86**: 460-465 [PMID: 9183482 DOI: 10.1111/j.1651-2227.1997.tb08913.x]
 - 39 **Sindhu KN**, Sowmyanarayanan TV, Paul A, Babji S, Ajampur SS, Priyadarshini S, Sarkar R, Balasubramanian KA, Wanke CA, Ward HD, Kang G. Immune response and intestinal permeability in children with acute gastroenteritis treated with Lactobacillus rhamnosus GG: A randomized, double-blind, placebo-controlled trial. *Clin Infect Dis* 2014; **58**: 1107-1115 [PMID: 24501384 DOI: 10.1093/cid/ciu065]
 - 40 **Aggarwal S**, Upadhyay A, Shah D, Teotia N, Agarwal A, Jaiswal V. Lactobacillus GG for treatment of acute childhood diarrhoea: An open labelled, randomized controlled trial. *Indian J Med Res* 2014; **139**: 379-385 [PMID: 24820831]
 - 41 **Salazar-Lindo E**, Miranda-Langschwager P, Campos-Sanchez M, Chea-Woo E, Sack RB. Lactobacillus casei strain GG in the treatment of infants with acute watery diarrhea: A randomized, double-blind, placebo controlled clinical trial [ISRCTN67363048]. *BMC Pediatr* 2004; **4**: 18 [PMID: 15345099 DOI: 10.1186/1471-2431-4-18]
 - 42 **Ahmadi E**, Alizadeh-Navaei R, Rezai MS. Efficacy of probiotic use in acute rotavirus diarrhea in children: A systematic review and meta-analysis. *Caspian J Intern Med* 2015; **6**: 187-195 [PMID: 26644891]
 - 43 **Liu F**, Li G, Wen K, Wu S, Zhang Y, Bui T, Yang X, Kocher J, Sun J, Jortner B, Yuan L. Lactobacillus rhamnosus GG on rotavirus-induced injury of ileal epithelium in gnotobiotic pigs. *J Pediatr Gastroenterol Nutr* 2013; **57**: 750-758 [PMID: 24280990 DOI: 10.1097/MPG.0b013e3182a356e1]
 - 44 **Jiang Y**, Ye L, Cui Y, Yang G, Yang W, Wang J, Hu J, Gu W, Shi C, Huang H, Wang C. Effects of Lactobacillus rhamnosus GG on the maturation and differentiation of dendritic cells in rotavirus-infected mice. *Benef Microbes* 2017; **8**: 645-656 [PMID: 28670908 DOI: 10.3920/BM2016.0157]
 - 45 **Cai S**, Kandasamy M, Rahmat JN, Tham SM, Bay BH, Lee YK, Mahendran R. Lactobacillus rhamnosus GG Activation of Dendritic Cells and Neutrophils Depends on the Dose and Time of Exposure. *J Immunol*

- Res* 2016; **2016**: 7402760 [PMID: 27525288 DOI: 10.1155/2016/7402760]
- 46 **Khoruts A.** Targeting the microbiome: From probiotics to fecal microbiota transplantation. *Genome Med* 2018; **10**: 80 [PMID: 30376869 DOI: 10.1186/s13073-018-0592-8]
 - 47 **Vemuri RC,** Gundamaraju R, Shinde T, Eri R. Therapeutic interventions for gut dysbiosis and related disorders in the elderly: Antibiotics, probiotics or faecal microbiota transplantation? *Benef Microbes* 2017; **8**: 179-192 [PMID: 28008784 DOI: 10.3920/BM2016.0115]
 - 48 **Boudraa G,** Benbouabdellah M, Hachelaf W, Boisset M, Desjeux JF, Touhami M. Effect of feeding yogurt versus milk in children with acute diarrhea and carbohydrate malabsorption. *J Pediatr Gastroenterol Nutr* 2001; **33**: 307-313 [PMID: 11593127 DOI: 10.1097/00005176-200109000-00015]
 - 49 **Kurugöl Z,** Koturoğlu G. Effects of *Saccharomyces boulardii* in children with acute diarrhoea. *Acta Paediatr* 2005; **94**: 44-47 [PMID: 15858959 DOI: 10.1111/j.1651-2227.2005.tb01786.x]
 - 50 **Henriksson R,** Bergström P, Franzén L, Lewin F, Wagenius G. Aspects on reducing gastrointestinal adverse effects associated with radiotherapy. *Acta Oncol* 1999; **38**: 159-164 [PMID: 10227436 DOI: 10.1080/028418699431564]
 - 51 **Uhnoo I,** Svensson L, Wadell G. Enteric adenoviruses. *Baillieres Clin Gastroenterol* 1990; **4**: 627-642 [PMID: 1962727 DOI: 10.1016/0950-3528(90)90053-J]
 - 52 **Cannon JP,** Lee TA, Bolanos JT, Danziger LH. Pathogenic relevance of *Lactobacillus*: A retrospective review of over 200 cases. *Eur J Clin Microbiol Infect Dis* 2005; **24**: 31-40 [PMID: 15599646 DOI: 10.1007/s10096-004-1253-y]
 - 53 **De Groote MA,** Frank DN, Dowell E, Glode MP, Pace NR. *Lactobacillus rhamnosus* GG bacteremia associated with probiotic use in a child with short gut syndrome. *Pediatr Infect Dis J* 2005; **24**: 278-280 [PMID: 15750472 DOI: 10.1097/01.inf.0000154588.79356.e6]
 - 54 **Molinaro M,** Aiazzi M, La Torre A, Cini E, Banfi R. [*Lactobacillus Rhamnosus* sepsis in a preterm infant associated with probiotic integrator use: A case report.]. *Recenti Prog Med* 2016; **107**: 485-486 [PMID: 27727257 DOI: 10.1701/2354.25230]
 - 55 **Gouriet F,** Million M, Henri M, Fournier PE, Raoult D. *Lactobacillus rhamnosus* bacteremia: An emerging clinical entity. *Eur J Clin Microbiol Infect Dis* 2012; **31**: 2469-2480 [PMID: 22544343 DOI: 10.1007/s10096-012-1599-5]
 - 56 **World Health Organization.** *The Treatment of Diarrhea: A Manual for Physicians and other Senior Health Workers.* Geneva: World Health Organization 2005;
 - 57 **Das RR.** Zinc in acute childhood diarrhea: Is it universally effective? *Indian J Pharmacol* 2012; **44**: 140; author reply 140-140; author reply 141 [PMID: 22345893 DOI: 10.4103/0253-7613.91891]
 - 58 **Karczewski J,** Troost FJ, Konings I, Dekker J, Kleerebezem M, Brummer RJ, Wells JM. Regulation of human epithelial tight junction proteins by *Lactobacillus plantarum* in vivo and protective effects on the epithelial barrier. *Am J Physiol Gastrointest Liver Physiol* 2010; **298**: G851-G859 [PMID: 20224007 DOI: 10.1152/ajpgi.00327.2009]
 - 59 **Mennigen R,** Nolte K, Rijcken E, Utech M, Loeffler B, Senninger N, Bruewer M. Probiotic mixture VSL#3 protects the epithelial barrier by maintaining tight junction protein expression and preventing apoptosis in a murine model of colitis. *Am J Physiol Gastrointest Liver Physiol* 2009; **296**: G1140-G1149 [PMID: 19221015 DOI: 10.1152/ajpgi.90534.2008]
 - 60 **Yang F,** Wang A, Zeng X, Hou C, Liu H, Qiao S. *Lactobacillus reuteri* I5007 modulates tight junction protein expression in IPEC-J2 cells with LPS stimulation and in newborn piglets under normal conditions. *BMC Microbiol* 2015; **15**: 32 [PMID: 25888437 DOI: 10.1186/s12866-015-0372-1]
 - 61 **Zyrek AA,** Cichon C, Helms S, Enders C, Sonnenborn U, Schmidt MA. Molecular mechanisms underlying the probiotic effects of *Escherichia coli* Nissle 1917 involve ZO-2 and PKC ζ redistribution resulting in tight junction and epithelial barrier repair. *Cell Microbiol* 2007; **9**: 804-816 [PMID: 17087734 DOI: 10.1111/j.1462-5822.2006.00836.x]
 - 62 **Fukushima Y,** Kawata Y, Hara H, Terada A, Mitsuoka T. Effect of a probiotic formula on intestinal immunoglobulin A production in healthy children. *Int J Food Microbiol* 1998; **42**: 39-44 [PMID: 9706796 DOI: 10.1016/S0168-1605(98)00056-7]
 - 63 **Rautava S,** Arvilommi H, Isolauri E. Specific probiotics in enhancing maturation of IgA responses in formula-fed infants. *Pediatr Res* 2006; **60**: 221-224 [PMID: 16864708 DOI: 10.1203/01.pdr.0000228317.72933.db]
 - 64 **Naidu AS,** Bidlack WR, Clemens RA. Probiotic spectra of lactic acid bacteria (LAB). *Crit Rev Food Sci Nutr* 1999; **39**: 13-126 [PMID: 10028126 DOI: 10.1080/10408699991279187]
 - 65 **Bujňáková D,** Kmet' V. Inhibitory potential of lactobacilli against *Escherichia coli* internalization by HT 29 cells. *Folia Microbiol (Praha)* 2012; **57**: 269-272 [PMID: 22528301 DOI: 10.1007/s12223-012-0122-9]
 - 66 **Gonzalez-Ochoa G,** Flores-Mendoza LK, Icedo-Garcia R, Gomez-Flores R, Tamez-Guerra P. Modulation of rotavirus severe gastroenteritis by the combination of probiotics and prebiotics. *Arch Microbiol* 2017; **199**: 953-961 [PMID: 28634691 DOI: 10.1007/s00203-017-1400-3]
 - 67 **Kang JY,** Lee DK, Ha NJ, Shin HS. Antiviral effects of *Lactobacillus ruminis* SPM0211 and *Bifidobacterium longum* SPM1205 and SPM1206 on rotavirus-infected Caco-2 cells and a neonatal mouse model. *J Microbiol* 2015; **53**: 796-803 [PMID: 26502964 DOI: 10.1007/s12275-015-5302-2]
 - 68 **Paim FC,** Langel SN, Fischer DD, Kandasamy S, Shao L, Alhamo MA, Huang HC, Kumar A, Rajashekara G, Saif LJ, Vlasova AN. Effects of *Escherichia coli* Nissle 1917 and Ciprofloxacin on small intestinal epithelial cell mRNA expression in the neonatal piglet model of human rotavirus infection. *Gut Pathog* 2016; **8**: 66 [PMID: 27999620 DOI: 10.1186/s13099-016-0148-7]
 - 69 **Kolaček S,** Hojsak I, Berni Canani R, Guarino A, Indrio F, Orel R, Pot B, Shamir R, Szajewska H, Vandenplas Y, van Goudoever J, Weizman Z; ESPGHAN Working Group for Probiotics and Prebiotics. Commercial Probiotic Products: A Call for Improved Quality Control. A Position Paper by the ESPGHAN Working Group for Probiotics and Prebiotics. *J Pediatr Gastroenterol Nutr* 2017; **65**: 117-124 [PMID: 28644359 DOI: 10.1097/MPG.0000000000001603]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

