



All

Images

Videos

翻译成中文

关闭取词

29,400 Results

Any time ▾

What are the risk factors for post-ERCP pancreatitis?

<https://www.medscape.com/answers/181364-14343> ▾

This patient with acute gallstone pancreatitis underwent endoscopic retrograde cholangiopancreatography. Cholangiogram shows no stones in common bile duct and multiple small stones in gallbladder.

Risk factors for post-ERCP pancreatitis: a systematic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4035895>

May 15, 2014 · Patients with pancreatic diseases, such as obstruction of the pancreatic duct and pancreatolithiasis, needed drainage of pancreatic juice via a pancreatic stent. A similar study was reported by Masci et al . in 2003 [7], which showed that the risk factors for PEP may change according to the development of instruments and technologies.

Cited by: 24

Author: Jian-Jun Chen, Xi-Mo Wang, Xing-Qian...

Publish Year: 2014

Risk factors for post-ERCP pancreatitis: a systematic ...

<https://eurjmedres.biomedcentral.com/articles/10.1186/2047-783X-19-26> ▾

May 15, 2014 · Conclusions. It appears that female gender, previous PEP, previous pancreatitis, endoscopic sphincterotomy, precut sphincterotomy, Sphincter of Oddi dysfunction, and non-prophylactic pancreatic duct stent are the risk factors for post-ERCP pancreatitis.

Cited by: 24

Author: Jian-Jun Chen, Xi-Mo Wang, Xing-Qian...

Publish Year: 2014

Prevention and Management of Post-ERCP Pancreatitis ...

pancreas.imedpub.com/prevention-and-management-of-postercp-pancreatitis.php?aid=162 ▾



INTRODUCTION

RECOGNITION

RISK FACTORS FOR POST-ER...

P



Endoscopic retrograde cholangiopancreatography (ERCP) is a fundamental tool in the management of in magnetic resonance cholangiopancreatography (MRCP) and endoscopic ultrasound (EUS), ERCP has biliary and pancreatic diseases. With technological advances evolved into a therapeutic rather than diagnostic procedure. Nevertheless, these procedures carry significant risk, with post-ERCP pancreatitis (PEP) being the most frequent and dreaded of these (Figure 1). The incidence of pancreatitis af...

See more on imedpub.com

Cited by: 1

Author: Ara B Sahakian, James L Buxbaum, Ja...

Publish Year: 2014

Name of Journal: *World Journal of Gastrointestinal Endoscopy*

Manuscript NO: 48811

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Risk factors for the development of **post-endoscopic retrograde cholangiopancreatography pancreatitis in patients with asymptomatic common bile duct stones**

Saito H *et al.* Risk factors for PEP in asymptomatic CBD stones

Match Overview

7 matches

1	Crossref 88 words Hirokazu Saito, Takehiko Koga, Masafumi Sakaguchi, Yoshihiro Kadono et al. "Post-endoscopic retrograde cholangio	4%
2	Crossref 29 words Hirokazu Saito, Takehiko Koga, Masafumi Sakaguchi, Yoshihiro Kadono et al. "Post-endoscopic retrograde cholangio	1%
3	Crossref 12 words Dumonceau, Jean-Marc, Angelo Andriulli, B. Elmunzer, Alberto Mariani, Tobias Meister, Jacques Deviere, Tomasz Mar	1%
4	Internet 10 words crawled on 29-Aug-2019 f6publishing.blob.core.windows.net	<1%



All

Images

Videos

翻译成中文

关闭取词

27,200 Results

Any time ▾

Post-endoscopic retrograde cholangiopancreatography ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4324870>

Nov 17, 2014 · Acute **pancreatitis** remains the most common complication of endoscopic retrograde cholangiopancreatography (ERCP). It is reported to occur in 2–10% of unselected patient samples and up to 40% of high-risk patients. The purpose of this article is to review the evidence behind the known risk factors ...

Cited by: 14

Author: Adarsh M. Thaker, Jeffrey D. Mosko, Tyle...

Publish Year: 2015

Post-endoscopic retrograde cholangiopancreatography ...

<https://academic.oup.com/gastro/article/3/1/32/589252> ▾

**Introduction**

Definition and Classification ...

Pathogenesis of Post-Ercp



Endoscopic retrograde cholangiopancreatography (ERCP) plays an ever-expanding role in the management of diseases involving the bile duct and pancreatic duct but, as an invasive procedure, it carries significant risks to the patient. The most common complication is acute pancreatitis, which is reported to occur in 2–10% of patients overall (ranging from 2–4% in low-risk patients up to 8–40% in high-risk patients) [1, 2]. A recent meta-analysis of 108 randomized, controlled trials (RCTs) report...

See more on [oup.com](https://academic.oup.com)

Cited by: 14

Author: Adarsh M. Thaker, Jeffrey D. Mosko, Tyle...

Publish Year: 2015

Published: Nov 17, 2014

Risk of pancreatitis after endoscopic retrograde ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2697892>

Aug 06, 2008 · **Pancreatitis** is the most common and serious complication of endoscopic retrograde cholangiopancreatography (ERCP), occurring in 2% 1 to 15% 2 of cases according to criteria defined by Cotton and others. 3 Several technical and patient-related **risk factors for post-ERCP pancreatitis**, which act independently or together, have been identified. 3 – 5 To date, clinicians have attempted to minimize the incidence and severity of post-ERCP ...

Cited by: 20

Author: Hiroyuki Matsubayashi, Akira Fukutomi, ...

Publish Year: 2009

Analysis of the risk factors for severity in post ...



전체

이미지

뉴스

쇼핑

지도

더보기

설정

도구

검색결과 약 97,100개 (0.64초)

<https://www.ncbi.nlm.nih.gov/pmc/PMC6278246> 이 페이지 번역하기

Risk Factors for Post-Endoscopic Retrograde ... - NCBI

G Li 저술 - 2018 - 3회 인용 - 관련 학술자료

2018. 11. 26. - The main indication was cholelithiasis (56.6%). The most **common** corresponding techniques were **endoscopic** nasobiliary drainage (65.8%), followed by **bile duct stone** removal with lithotomy balloon (58.3%). Based on the recorded **ERCP**, complications occurred in 80 (4.48%) **patients**.

관련 질문

How common are bile duct stones? ▼

What happens if you have a stone in your bile duct? ▼

What causes stones to form in the bile duct? ▼

What are the complications of ERCP? ▼

사용자 의견

<https://www.ncbi.nlm.nih.gov/pmc/PMC6124257> 이 페이지 번역하기

Risk factors for recurrence of common bile duct stones after ...

S Li 저술 - 2018 - 2회 인용 - 관련 학술자료

2018. 6. 4. - EST for **common bile duct stones** is safe as indicated by **patients'** ... Keywords: