

Match Overview

Rank	Source	Words	Similarity
1	Internet crawled on 04-Nov-2017 repository.kulib.kyoto-u.ac.jp	91 words	3%
2	Internet crawled on 03-Mar-2017 bmccancer.biomedcentral.com	24 words	1%
3	Crossref "Abstracts of the 48th EASD Annual Meeting of the European Association for the Study of Diabetes", Diabetologia, 2012	18 words	1%
4	Internet crawled on 05-Jul-2019 f6publishing.blob.core.windows.net	14 words	<1%
5	Internet crawled on 02-Aug-2019 www.dovepress.com	12 words	<1%
6	Internet crawled on 31-Jul-2019 www.tandfonline.com	12 words	<1%

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 48925

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Pathological response measured using virtual microscopic slides for gastric cancer patients who underwent neoadjuvant chemotherapy

Kawai S *et al.* Predictivity of pathological response

Sadayuki Kawai, Tadakazu Shimoda, Takashi Nakajima, Masanori Terashima, Katsuhiko Omae, Nozomu Machida, Hirofumi Yasui

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 70,700 条结果 (用时 0.56 秒)

Advanced Gastric Cancer - Learn About a Treatment Option

 [Prescription treatment website](#) ▼

Find Information About a Treatment Option for Advanced Gastric Cancer. 24/7 Nurse Support. Online Tools & Resources. Patient-Friendly Language. Patient Support Program. Caregiver Information. Doctor Discussion Guide. Product Information. Financial Assistance.

Google 学术: Pathological response measured using virtual microscopic slides for gastric cancer patients who underwent neoadjuvant chemotherapy

... carcinoma treated with neoadjuvant chemotherapy - Becker - 被引用次数: 450

... define the pathological response rate for gastric cancer ... - Nakamura - 被引用次数: 10

... treatment response and outcome of patients with ... - Ribeiro Jr - 被引用次数: 152

phase II trial of neoadjuvant cisplatin–fluorouracil followed by ...

<https://academic.oup.com/annonc/article/17/9/1404/214867> - 翻译此页

作者: B Brenner - 2006 - 被引用次数: 65 - 相关文章

Background: The aim of the study was to evaluate the efficacy and toxicity of neoadjuvant chemotherapy with intravenous (i.v.) cisplatin and fluorouracil (5-FU), surgery and postoperative intraperitoneal (i.p.) floxuridine (FUdR) and leucovorin (LV) in patients with locally advanced gastric cancer.

Correlation of pathological complete response with survival after ...

<https://www.ncbi.nlm.nih.gov/pubmed/29370182> - 翻译此页

作者: Z Li - 2018 - 被引用次数: 5 - 相关文章

Tumor blood supply may predict neoadjuvant chemotherapy ...

<https://journals.sagepub.com/doi/full/10.1177/0300060519845491>

Gastric cancer (GC) is the second most common cause of cancer-related deaths worldwide, with almost 50% of all cases occurring in China. 1 Most patients are already at an advanced stage when a diagnosis of GC is made. 2,3 Although surgical resection is the main treatment for localized/regional GC, many patients with advanced GC fail to achieve satisfactory R0 resection rates after surgery alone.

Author: Xiang Ji, Qiaoyun Yang, Hui Qin, Jie ... Publish Year: 2019

Evaluating the response of gastric carcinomas to ...

<https://www.sciencedirect.com/science/article/pii/S0009926015002767>

Introduction. Preoperative neoadjuvant chemotherapy (NC) has been successfully applied to the treatment of locally advanced gastric carcinomas. 1, 2, 3 For patients who responded well to NC, prolonged survival times were observed compared with non-responsive patients. 1 In general, the potential benefits of NC include a reduction in tumour volume and a decrease in tumour staging for ...

Cited by: 12 Author: L. Tang, Z.-Y. Li, Z.-W. Li, X.-P. Zhang, Y...

Publish Year: 2015

Assessment of Histopathological Response in Gastric and ...

<https://www.hindawi.com/journals/ism/2012/519351>

Both microscopic and macroscopic features of a tumour are used to assess response and the absence of cancer cells in the resected specimen represents a complete response [8]. Less than 30% of patients receiving neoadjuvant therapy achieve a complete pathologic response [9].

Cited by: 11 Author: A. Mirza, A. Naveed, S. Hayes, L. Formel...

Publish Year: 2012

[PDF] Determination of the optimal cutoff percentage of ...

<https://link.springer.com/content/pdf/10.1007/s10120-014-0401-z.pdf>

The International Gastric Cancer Association and The Japanese Gastric Cancer Association 2014 Abstract Background Pathological response rate (pathRR) is a common endpoint used to assess the efficacy of preoperative therapy for gastric cancer. PathRR is estimated based on the percentage of the residual tumor area in the primary tumorous bed.

Cited by: 8 Author: Kenichi Nakamura, Takeshi Kuwata, Tada...

Publish Year: 2015

Determination of the optimal cutoff percentage of residual ...

11,600 Results Any time ▾

[Regional Arterial Infusion Chemotherapy improves the ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4664910>

Regional Arterial Infusion Chemotherapy improves the Pathological Response rate for advanced gastric cancer with Short-term Neoadjuvant Chemotherapy Zhen-Feng Wu , 1 Qin-Hong Cao , 1 Xiao-Yu Wu , a, 1, * Che Chen , 1 Zhe Xu , b, 1, * Wei-Su Li , 1 Xue-Quan Yao , 1 and Fu-Kun Liu 1

Cited by: 2 Author: Zhenfeng Wu, Qin-hong Cao, Xiaoyu Wu,...

Publish Year: 2015

[Tumor blood supply may predict neoadjuvant chemotherapy ...](#)

<https://journals.sagepub.com/doi/full/10.1177/0300060519845491>

Gastric cancer (GC) is the second most common cause of cancer-related deaths worldwide, with almost 50% of all cases occurring in China. 1 Most patients are already at an advanced stage when a diagnosis of GC is made. 2,3 Although surgical resection is the main treatment for localized/regional GC, many patients with advanced GC fail to achieve satisfactory R0 resection rates after surgery alone.

Author: Xiang Ji, Qiaoyun Yang, Hui Qin, Jie ... Publish Year: 2019

[Evaluating the response of gastric carcinomas to ...](#)

<https://www.sciencedirect.com/science/article/pii/S0009926015002767>

Introduction. Preoperative neoadjuvant chemotherapy (NC) has been successfully applied to the treatment of locally advanced gastric carcinomas.1, 2, 3 For patients who responded well to NC, prolonged survival times were observed compared with non-responsive patients. 1 In general, the potential benefits of NC include a reduction in tumour volume and a decrease in tumour staging for ...

Cited by: 12 Author: L. Tang, Z.-Y. Li, Z.-W. Li, X.-P. Zhang, Y....

Publish Year: 2015

[Assessment of Histopathological Response in Gastric and ...](#)

<https://www.hindawi.com/journals/isrn/2012/519351>

Furthermore, the results published by Ajani et al. for a series of gastric cancer patients treated with neoadjuvant chemoradiotherapy showed a complete response rate of 20% with a 100% five-year survival rate . The combination therapy appears to be superior to chemotherapy alone in achieving complete response.

Cited by: 11 Author: A. Mirza, A. Naveed, S. Hayes, L. Formel...

Publish Year: 2012

[Pathological Response after Neoadjuvant Treatment in ...](#)

https://www.researchgate.net/publication/296696196_Pathological_Response_after...

Pathological Response after Neoadjuvant Treatment in Advanced Gastric Cancer: Impact on Survival