



All

Images

Videos

翻译成中文

关闭取词

4,420 Results

Any time ▾

Intravenous infusion of lidocaine significantly reduces ...

[https://bjanaesthesia.org/article/S0007-0912\(18\)30530-0/fulltext](https://bjanaesthesia.org/article/S0007-0912(18)30530-0/fulltext)

Jul 01, 2018 · **Propofol** use during sedation for colonoscopy can result in cardiopulmonary complications. **Intravenous lidocaine** can alleviate visceral pain and **decrease propofol requirements** during surgery. We tested the hypothesis that i.v. **lidocaine reduces propofol requirements** during colonoscopy and **improves post-colonoscopy recovery**.

Author: C. Forster, Audrey Vanhauzenhuyse, ... **Publish Year:** 2018

Safety and efficacy of etomidate and propofol anesthesia ...

https://www.researchgate.net/publication/304401097_Safety_and_efficacy_of_etomidate...

Request PDF on ResearchGate | **Safety and efficacy of etomidate and propofol anesthesia** in elderly patients undergoing **gastroscopy**: A **double-blind randomized** clinical study | The aim of the present

Effect of propofol combined with opioids on cough reflex ...

<https://bmjopen.bmj.com/content/7/9/e014881> ▾

Sep 01, 2017 · We will observe the incidence and **degree of reflex coughing** and gagging under **sedation** when using **propofol combined** with one of the above **drugs** or **propofol alone**. Methods and analysis This will be a prospective, randomised, **double-blind**, controlled trial. **ASA I-II level** patients aged 18–65 years and scheduled for **gastroscopy** will be included.

Cited by: 1 **Author:** Ning Yin, Jiangyan Xia, Yi-Zhi Cao, Xinjia...

Publish Year: 2017

Sedation in gastrointestinal endoscopy: Current issues

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3558570>

Jan 28, 2013 · The 6.5-mg/kg **dose of fospropofol** provides the ideal balance of **efficacy and safety** for patients undergoing colonoscopy. In a double-blind trial evaluating 127 patients who received fospropofol (2, 5, 6.5 or 8 mg/kg) or midazolam 0.02 mg/kg following **pre-treatment** with fentanyl, fospropofol yielded a significant **dose-dependent increase** in sedation success from 24% (2 mg/kg), 35% (5 mg/kg) and ...

Cited by: 135 **Author:** John K Triantafillidis, Emmanuel Merikas,...

Publish Year: 2013

Sedation for endoscopy: The safe use of propofol by ...

https://www.researchgate.net/publication/11450428_Sedation_for_endoscopy_The_safe_use...

The recovery time, patient satisfaction, nasal intubation rate, the safety and the efficacy of PDDC

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 49721

Manuscript Type: ORIGINAL ARTICLE

Randomized Controlled Trial

Combination of propofol and dezocine to improve the safety and efficacy of anesthesia for gastroscopy and colonoscopy in adults: A randomized, double-blind, controlled trial

Li XT *et al.* Propofol and dezocine in endoscopy

Xue-Ting Li, Chao-Qun Ma, Si-Hua Qi, Li-Min Zhang

Abstract

BACKGROUND

Gastroscopy and colonoscopy are important and common endoscopic methods for

Match Overview

1	Crossref 150 words Minqiang Liu, Hongyan Wu, Danling Yang, Fengxian Li, Zhi chao Li, Song Wang, Renliang He. "Effects of small-dose r	5%
2	Crossref 133 words Ning Yin, Jiangyan Xia, Yi-Zhi Cao, Xinjian Lu, Jing Yuan, Ju e Xie. "Effect of propofol combined with opioids on cough ...	5%
3	Crossref 21 words Zhu, Han. "Analgesic, sedative effects and safety of dezocine in patients undergoing endoscopic retrograde cholangi ...	1%
4	Crossref 20 words Ayse Oner, Ali Akal, Nuri Erdogan, Hakki Dogan, Mithat One r. "Color Doppler Imaging of Ocular Hemodynamic Chan...	1%
5	Internet 17 words crawled on 25-Jan-2019 anesthesiology.pubs.asahq.org	1%
6	Internet 17 words crawled on 11-Aug-2016 www.wjgnet.com	1%



All

Images

Videos

翻译成中文

关闭取词

6,030 Results

Any time ▾

Effect of propofol combined with opioids on cough reflex ...

<https://bmjopen.bmj.com/content/7/9/e014881> ▾

Sep 01, 2017 · We will observe the incidence and **degree of reflex coughing** and gagging under **sedation** when using **propofol combined** with one of the above **drugs** or **propofol alone**. Methods and analysis This will be a prospective, randomised, **double-blind**, controlled trial. ASA I-II level patients aged 18–65 years and scheduled for **gastroscopy** will be included.

Cited by: 2

Author: Ning Yin, Jiangyan Xia, Yi-Zhi Cao, Xinjia...

Publish Year: 2017

Safety and efficacy of combined use of propofol and ...

https://www.researchgate.net/publication/333142530_Safety_and_efficacy_of_combined_use...

Download Citation on ResearchGate | **Safety and efficacy of combined use of propofol** and etomidate for sedation during **gastroscopy**: Systematic review and meta-analysis | Background: Sedation with ...

Intravenous infusion of lidocaine significantly reduces ...

[https://bjanaesthesia.org/article/S0007-0912\(18\)30530-0/fulltext](https://bjanaesthesia.org/article/S0007-0912(18)30530-0/fulltext)

Propofol use during sedation for colonoscopy can result in cardiopulmonary complications. **Intravenous lidocaine** can alleviate visceral pain and **decrease propofol requirements** during surgery. We tested the hypothesis that i.v. **lidocaine reduces propofol requirements** during colonoscopy and **improves post-colonoscopy recovery**.

Author: C. Forster, Audrey Vanhaudenhuyse, ...

Publish Year: 2018

Sedation in gastrointestinal endoscopy: Current issues

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3558570>

Jan 28, 2013 · In a randomised trial comparing the **efficacy** and recovery time of two **sedation regimens** consisting of **midazolam in combination** with either meperidine or fentanyl in patients submitted for colonoscopy, it was found that the use of fentanyl in **combination** with **low-dose midazolam** resulted in a significantly faster recovery from sedation compared with meperidine, without any apparent loss of **analgesic effect**.

Cited by: 142

Author: John K Triantafillidis, Emmanuel Merikas,...

Publish Year: 2013