

World Journal of *Meta-Analysis*

World J Meta-Anal 2019 July 31; 7(7): 339-372





EDITORIAL

- 339 Asymptomatic bacteriuria among hospitalized diabetic patients: Should they be treated?
Ramos-Ramirez MJ, Surani S

- 343 Artificial intelligence for endoscopy
Imaeda H, Tsuzuki Y, Miyaguchi K, Ashitani K, Ohgo H, Nakamoto H

OPINION REVIEW

- 346 Phantom of the inflammasome in the gut: Cytomegalovirus
Arslan F, Vahaboglu H

SYSTEMATIC REVIEWS

- 350 Pediatric recurrent *Clostridium difficile* infections in immunocompetent children: Lessons learned from case reports of the first twelve consecutive patients
Chu A, Michail S

META-ANALYSIS

- 358 Effect of dl-3-n-butylphthalide on infarction volume in animal models of ischemic stroke: A meta-analysis
Luan D, Wu ZY, Zhang YX, Yuan LL, Xu Y, Chu ZH, Ma LS, Wang YP, Zhao SC

ABOUT COVER

Editorial Board of *World Journal of Meta-Analysis*, Edmundas Kadusevicius, PhD, Professor, Department of Clinical Pharmacology, Institute of Physiology and Pharmacology, Lithuanian University of Health Sciences, Kaunas LT-50009, Lithuania

AIMS AND SCOPE

The *World Journal of Meta-Analysis* (WJMA) is an online, open-access, single-blind peer-reviewed journal that is published by Baishideng Publishing Group. Accepting both solicited and unsolicited manuscripts, the WJMA mainly publishes high quality meta-analysis and systematic review articles to lead the development of various fields of clinical medicine. The type of articles published in the WJMA include the editorial, minireview, review, meta-analysis, meta-regression, systematic review, and evidence-based practice.

The primary aim of the WJMA is to provide scholars and readers from various fields of clinical medicine with a platform to publish high quality meta-analysis and systematic review articles and communicate their research findings online.

The WJMA mainly publishes research results and findings obtained through meta-analysis and systematic review in a wide range of areas, including medicine, pharmacy, etc.

INDEXING/ABSTRACTING

The WJMA is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: Ji-Hong Liu

Proofing Production Department Director: Yun-Xiaojuan Wu

NAME OF JOURNAL

World Journal of Meta-Analysis

ISSN

ISSN 2308-3840 (online)

LAUNCH DATE

May 26, 2013

FREQUENCY

Irregular

EDITORS-IN-CHIEF

Giuseppe Biondi-Zoccai

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2308-3840/editorialboard.htm>

EDITORIAL OFFICE

Jin-Lei Wang, Director

PUBLICATION DATE

July 31, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Artificial intelligence for endoscopy

Hiroyuki Imaeda, Yoshikazu Tsuzuki, Kazuya Miyaguchi, Keigo Ashitani, Hideki Ohgo, Hidetomo Nakamoto

ORCID number: Hiroyuki Imaeda (0000-0003-0898-6540); Yoshikazu Tsuzuki (0000-0003-3029-7469); Kazuya Miyaguchi (0000-0003-0701-3608); Keigo Ashitani (0000-0002-5043-892X); Hideki Ohgo (0000-0002-3916-2918); Hidetomo Nakamoto (0000-0003-3975-5067).

Author contributions: Imaeda H made the conception, designed the study and drafted of the article; Miyaguchi K, Ashitani K, Tsuzuki Y, Nakamoto H critical revised the article for important intellectual content; Imaeda H, Tsuzuki Y, Miyaguchi K, Ashitani K, Nakamoto H made the final approval of the article.

Conflict-of-interest statement: The authors have no conflict of interest to declare.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: June 25, 2019

Peer-review started: June 25, 2019

First decision: July 20, 2019

Revised: July 27, 2019

Hiroyuki Imaeda, Yoshikazu Tsuzuki, Hideki Ohgo, Department of Gastroenterology, Saitama Medical University, Iruma-gun, Saitama 3500495, Japan

Kazuya Miyaguchi, Keigo Ashitani, Hidetomo Nakamoto, Department of General Internal Medicine, Saitama Medical University, Iruma-gun, Saitama 3500495, Japan

Corresponding author: Hiroyuki Imaeda, MD, Professor, Department of Gastroenterology, Saitama Medical University, Morohongo 38, Moroyama-machi, Iruma-gun, Saitama 3500495, Japan. imaedahi@yahoo.co.jp

Telephone: +81-49-2761667

Fax: +81-49-2761667

Abstract

In recent times, there has been progressive development in artificial intelligence (AI) following the introduction of deep learning in the medical field including gastroenterology and endoscopy. Most of the reported studies were based on retrospective data. Several prospective studies of real-time diagnosis of moving images using the AI system are expected to match the real clinical situation and to aid the endoscopists in the detection and diagnosis of neoplasms without missing any lesion. AI can read a large number of endoscopic images in a few minutes and make a diagnosis; therefore, it is expected to cover the lack of support for the screening esophagogastroduodenoscopy in the health check-up and a large number of capsule images, thereby freeing the endoscopists from this burden. AI can help make the diagnosis during the endoscopic procedure and thereby prevent an unnecessary biopsy for patients taking antithrombotic drugs. AI can also be useful for education and training in endoscopy. Trainees can learn to perform endoscopy and the detection and diagnosis of lesions by the support of AI. In the near future, real-time endoscopic diagnosis using AI is expected to lessen the burden of endoscopists, to enhance the quality level of endoscopists, to overcome the miss of lesions and to make optimal diagnosis.

Key words: Artificial Intelligence; Endoscopy; Gastric cancer; Colonic neoplasm

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Artificial intelligence (AI) has an increasing role in medical imaging in recent times. It has numerous benefits in the field of endoscopy. It aids in the accurate identification and diagnosis of lesions. AI also helps in reading and accurately interpreting large volumes of endoscopic images. It can play a role in the training of endoscopists as well.

Accepted: July 29, 2019**Article in press:** July 29, 2019**Published online:** July 31, 2019**P-Reviewer:** De Quadros LG, Jun C, Skok P, Barreto SG**S-Editor:** Dou Y**L-Editor:** A**E-Editor:** Liu JH**Citation:** Imaeda H, Tsuzuki Y, Miyaguchi K, Ashitani K, Ohgo H, Nakamoto H. Artificial intelligence for endoscopy. *World J Meta-Anal* 2019; 7(7): 343-345**URL:** <https://www.wjnet.com/2308-3840/full/v7/i7/343.htm>**DOI:** <https://dx.doi.org/10.13105/wjma.v7.i7.343>

ARTIFICIAL INTELLIGENCE FOR ENDOSCOPY

In recent times, there has been progressive development in artificial intelligence (AI) following the introduction of deep learning in various fields. As expected, AI has been introduced into the medical field, and many papers have reported on its use in different specialties in the medical field, including gastroenterology and endoscopy^[1].

In the field of endoscopy, conventional endoscopy, magnifying endoscopy and endocytoscopy using white light images and optical digital images for gastrointestinal neoplasia, gastritis related to *Helicobacter pylori* infection, and ulcerative colitis have been reported. Most of the reported studies were based on retrospective data. In addition, the results from these retrospective studies have been based on good-quality still images and not on either low-quality images or moving images^[2,3]. Real-time imaging involves moving images from different levels in the endoscopic fields and not still images. Therefore, the results from retrospective studies might not match the real images. Several prospective studies of real-time diagnosis using the AI system have also been reported^[4]. The data from these studies are expected to match the real clinical situation and to be more beneficial. For more robust clinical verification, well-designed multicenter prospective studies with adequate inclusion/ exclusion criteria that represent the target population are needed^[1]. Moreover, the efficiency and accuracy of AI increases as the amount of data increases: For example, the use of moving images. Based on the prospective studies, real-time detection and diagnosis of lesions during endoscopic procedure are expected to become feasible. At present there is high sensitivity for the detection of gastric cancers; however, the specificity is not so good^[5], because most gastric cancers involve gastric inflammation as well, which makes the gastric mucosa red or white, irregular, and granular. Gastric inflammation causes erosion, ulcer, or polyp, and is relatively similar in appearance to the mucosa in gastric cancer. On the other hand, the diagnosis of colonic neoplasms using narrow band imaging (NBI), magnifying endoscopy with NBI, or endocytoscopy shows high sensitivity as well as specificity. Recently, EndoBRAIN software (Olympus Medical Systems, Japan) which is an AI system used in endocytoscopy for colonic neoplasms has become available in Japan. This allows the differential diagnosis of a colonic lesion and the confirmation of a colonic neoplasm to be made in a very short time. The endocytoscopy involves a special scope which magnifies the target lesion 520 times and enables observation at the cell level as with a microscope, whereas, EndoBRAIN software has not detected colonic polyps automatically yet.

Until now, it usually took huge time and effort for endoscopists to learn about the many gastrointestinal diseases and train in the endoscopic detection and diagnosis of gastric cancer or colonic neoplasm. Even when the endoscopists are experts, they might sometimes miss the detection and diagnosis of the neoplasms due to similar color of the lesions to the surrounding area, small size of the lesion, difficult location such as behind the folds or on the bending site, and lesions endoscopically observed in a moment among a lot of visual images. AI is expected to aid the endoscopists in the detection and diagnosis of neoplasms without missing any lesion.

AI also helps in the quick interpretation of endoscopic images taken in screening esophagogastroduodenoscopy (EGD) without waste of time. In Japan, screening EGD is conducted as part of the health check-up, and a double check of the screening EGD images is necessary. However, the number of endoscopists is not enough, especially in the local suburbs. Extensive time is required to interpret numerous endoscopic images, and the interpretation of endoscopic images is a burden for Japanese endoscopists. AI can read a large number of endoscopic images in a few minutes and make a diagnosis; therefore, it is expected to cover the lack of support for the screening EGD in the health check-up, thereby freeing the endoscopists from this burden. Moreover, there are numerous software that can easily detect lesions in capsule endoscopy. The interpretation of a large number of capsule images by the endoscopists takes a lot of time, and AI can free the endoscopists from the burden of capsule endoscopy image interpretation.

Lately, there has been a gradual increase in the number of patients taking antithrombotic drugs, and there is a hesitancy in performing a biopsy in these

patients. Although the guidelines for patients taking antithrombotic drugs is available not only in Western countries^[6], but also in Eastern countries^[7,8], and endoscopic biopsies are allowed in these patients, bleeding following a biopsy is sometimes observed. In addition, colonic polyps were sometimes resected and discarded during colonoscopic procedure, therefore, it is important to decide the indication of resection and discard for colonic polyps. AI can help make the diagnosis in vivo during the endoscopic procedure and thereby prevent an unnecessary biopsy and resection, and careless discard.

AI can also be useful for education and training in endoscopy. Generally, instructors teach trainees on the various aspects of endoscopy. Even in the absence of an instructor in some hospitals, trainees can learn to perform endoscopy and the detection and diagnosis of lesions by the support of AI. However, if they always rely on AI for the diagnosis, the diagnostic acumen of the trainees does not improve. The trainees need to learn detection and diagnosis of lesions without the use of AI as well.

On the other hand, as deep learning algorithm is black-box, the machine-generated decision is sometimes hard to understand for endoscopists and it does not match the diagnosis by endoscopists. Therefore, the level of detection and diagnosis sometimes has to be checked periodically.

In the near future, real-time endoscopic diagnosis during endoscopic procedures with real moving images using AI is expected to become widespread in all endoscopic fields, to lessen the burden of endoscopists and to enhance the quality level of endoscopists. AI is desired to overcome the miss of lesions by endoscopy and to make endoscopic diagnosis comparable with optimal diagnostic accuracy by histopathological findings, and to reduce medical costs by avoiding pathological examination and unnecessary resection.

REFERENCES

- 1 **Yang YJ**, Bang CS. Application of artificial intelligence in gastroenterology. *World J Gastroenterol* 2019; **25**: 1666-1683 [PMID: [31011253](#) DOI: [10.3748/wjg.v25.i14.1666](#)]
- 2 **Mori Y**, Kudo SE, Mohamed HEN, Misawa M, Ogata N, Itoh H, Oda M, Mori K. Artificial intelligence and upper gastrointestinal endoscopy: Current status and future perspective. *Dig Endosc* 2019; **31**: 378-388 [PMID: [30549317](#) DOI: [10.1111/den.13317](#)]
- 3 **Kudo SE**, Mori Y, Misawa M, Takeda K, Kudo T, Itoh H, Oda M, Mori K. Artificial intelligence and colonoscopy: Current status and future perspectives. *Dig Endosc* 2019; **31**: 363-371 [PMID: [30624835](#) DOI: [10.1111/den.13340](#)]
- 4 **Mori Y**, Kudo SE, Chiu PW, Singh R, Misawa M, Wakamura K, Kudo T, Hayashi T, Katagiri A, Miyachi H, Ishida F, Maeda Y, Inoue H, Nimura Y, Oda M, Mori K. Impact of an automated system for endocytoscopic diagnosis of small colorectal lesions: an international web-based study. *Endoscopy* 2016; **48**: 1110-1118 [PMID: [27494455](#) DOI: [10.1055/s-0042-113609](#)]
- 5 **Hirasawa T**, Aoyama K, Tanimoto T, Ishihara S, Shichijo S, Ozawa T, Ohnishi T, Fujishiro M, Matsuo K, Fujisaki J, Tada T. Application of artificial intelligence using a convolutional neural network for detecting gastric cancer in endoscopic images. *Gastric Cancer* 2018; **21**: 653-660 [PMID: [29335825](#) DOI: [10.1007/s10120-018-0793-2](#)]
- 6 **ASGE Standards of Practice Committee**. Acosta RD, Abraham NS, Chandrasekhara V, Chathadi KV, Early DS, Eloubeidi MA, Evans JA, Faulx AL, Fisher DA, Fonkalsrud L, Hwang JH, Khashab MA, Lightdale JR, Muthusamy VR, Pasha SF, Saltzman JR, Shaikat A, Shergill AK, Wang A, Cash BD, DeWitt JM. The management of antithrombotic agents for patients undergoing GI endoscopy. *Gastrointest Endosc* 2016; **83**: 3-16 [PMID: [26621548](#) DOI: [10.1016/j.gie.2015.09.035](#)]
- 7 **Fujimoto K**, Fujishiro M, Kato M, Higuchi K, Iwakiri R, Sakamoto C, Uchiyama S, Kashiwagi A, Ogawa H, Murakami K, Mine T, Yoshino J, Kinoshita Y, Ichinose M, Matsui T; Japan Gastroenterological Endoscopy Society. Guidelines for gastroenterological endoscopy in patients undergoing antithrombotic treatment. *Dig Endosc* 2014; **26**: 1-14 [PMID: [24215155](#) DOI: [10.1111/den.12183](#)]
- 8 **Kato M**, Uedo N, Hokimoto S, Ieko M, Higuchi K, Murakami K, Fujimoto K. Guidelines for Gastroenterological Endoscopy in Patients Undergoing Antithrombotic Treatment: 2017 Appendix on Anticoagulants Including Direct Oral Anticoagulants. *Dig Endosc* 2018; **30**: 433-440 [PMID: [29733468](#) DOI: [10.1111/den.13184](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

