



All

Images

Videos

关闭取词

36,900 Results

Any time ▾

Preventing Suicide Among Inpatients

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4079240>

Patients with a **psychiatric** history or attempt were "known at risk for suicide" but 14% of general **hospital suicides** were "unknown at risk." 31, p 1 Previous attempted **suicide**, especially if recent, being on antidepressants, having physical health problems (such as **chronic pain**), poor health prognosis, social stressors, hopelessness, and substance abuse constituted risks. Methods were by hanging, ...

Cited by: 24

Author: Isaac Sakinofsky

Publish Year: 2014

Suicide attempts and related factors in patients admitted ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3078091>

Mar 31, 2011 · A variety of factors are **associated with an increased risk of suicide** and attempted **suicide**, including **psychiatric disorders** [9-14], feelings of **hopelessness** and impulsivity [15,16], history of previous **suicide attempts** [9,17,18], age, **sex and race** [19-24], marital status [25,26], occupation, comorbidity [28,29], **adverse childhood** experiences, family history [31,32], and ...

Cited by: 65

Author: Jesús Alberdi-Sudupe, Salvador Pita-Fern...

Publish Year: 2011

Prevalence and Determinants of Depression and Anxiety ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4852078>

Jun 27, 2015 · Mood disorders are prevalent in **hospitalized patients**. However, risk factors for early diagnosis have not been studied exclusively in surgical **patients**. Our study aimed to investigate the **prevalence** and determinants of depression and anxiety symptoms in surgical **patients**...

Cited by: 8

Author: Saeed Shoar, Mohammad Naderan, Mota...

Publish Year: 2016

Prevalence and correlates of suicide attempt among Chinese ...

<https://www.nature.com/articles/s41598-019-52440-x>

7 days ago · These studies reported a wide range of lifetime **prevalence** of attempted **suicide** in **patients** with opioid dependence (8.0–48.0%) and a variety of risk factors **associated with suicide** ...

Prevalence and factors associated with depression among ...

<https://bmcrenotes.biomedcentral.com/articles/10.1186/s13104-019-4109-3> ▾

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 52152

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Prevalence and associated factors of suicide among hospitalized schizophrenic patients

Woottiluk P *et al.* Suicide among hospitalized schizophrenic patients

Pakapan Woottiluk, Benchalak Maneeton, Natthanan Jaiyen, Wajana Khemawichanurat, Suttipong Kawilapat, Narong Maneeton

Abstract

BACKGROUND

Previous studies found several factors associated with suicide in schizophrenic

Match Overview

1	Internet 57 words crawled on 13-Mar-2016 www.dovepress.com	2%
2	Internet 42 words crawled on 06-Dec-2018 link.springer.com	1%
3	Crossref 36 words Beckwée, David, Ivan Bautmans, Thierry Scheerlinck, and Peter Vaes. "Exercise in knee osteoarthritis – preliminary fi	1%
4	Publications 12 words DEMIRPENCE, Mustafa, AKMAN, Tulay, OZTOP, Ilhan, UNE K, Ilkay T., YAVUZSEN, Tugba and YILMAZ, Ahmet U.. "Impa	<1%



ALL

IMAGES

VIDEOS

关闭取词

47,800 Results

Any time ▾

The Prevalence of Suicide Attempts and Independent ...

https://www.researchgate.net/publication/333175146_The_Prevalence_of_Suicide_Attempts...

The Prevalence of Suicide Attempts and Independent Demographic and Clinical Correlates among Chronic Schizophrenia Patients in Agricultural Areas of China Article in Psychiatric Quarterly 90(3 ...

The Prevalence of Suicide Attempts and Independent ...

<https://link.springer.com/article/10.1007/s11126-019-09644-1> ▾

May 17, 2019 · Lifetime SA is common among hospitalized schizophrenia patients living in agricultural areas of China. For suicide prevention, regular assessments, appropriate interventions and clinical management should be integrated into a community-based psychiatric service model for this population.

Author: Yi Zhong, Lei Xia, Tong Tong Zhao, Yu... Publish Year: 2019

Preventing Suicide Among Inpatients

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4079240>

Patients with a psychiatric history or attempt were "known at risk for suicide" but 14% of general hospital suicides were "unknown at risk." 31, p 1 Previous attempted suicide, especially if recent, being on antidepressants, having physical health problems (such as chronic pain), poor health prognosis, social stressors, hopelessness, and substance abuse constituted risks. Methods were by hanging, ...

Cited by: 24 Author: Isaac Sakinofsky

Publish Year: 2014

Suicide attempts and related factors in patients admitted ...

<https://bmcpsy psychiatry.biomedcentral.com/articles/10.1186/1471-244X-11-51> ▾

Mar 31, 2011 · A variety of factors are associated with an increased risk of suicide and attempted suicide, including psychiatric disorders [9–14], feelings of hopelessness and impulsivity [15, 16], history of previous suicide attempts [9, 17, 18], age, sex and race [19–24], marital status [25, 26], occupation, comorbidity [28, 29], adverse childhood experiences, family history [31, 32], and ...

Cited by: 65 Author: Jesús Alberdi-Sudupe, Salvador Pita-Fern...

Publish Year: 2011

Prevalence and factors associated with depression among ...

<https://bmccresnotes.biomedcentral.com/articles/10.1186/s13104-019-4109-3> ▾

Feb 04, 2019 · Identifying factors associated with depression will help health care providers to design



44,500 Results

Any time ▼

The prevalence and associated factors of depression among ...

<https://link.springer.com/article/10.1186/s12888-019-2419-6> ▼

Background Depression is common among people with schizophrenia and associated with severe positive and negative symptoms, higher rates of disability, treatment resistance and mortality related to... The prevalence and associated factors of depression among patients with schizophrenia in Addis Ababa, Ethiopia, cross-sectional study | SpringerLink

Preventing Suicide Among Inpatients

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4079240>

Patients with a psychiatric history or attempt were "known at risk for suicide" but 14% of general hospital suicides were "unknown at risk." 31, p 1 Previous attempted suicide, especially if recent, being on antidepressants, having physical health problems (such as chronic pain), poor health prognosis, social stressors, hopelessness, and substance abuse constituted risks. Methods were by hanging, ...

Cited by: 24

Author: Isaac Sakinofsky

Publish Year: 2014

Suicide attempts and related factors in patients admitted ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3078091>

Mar 31, 2011 · A variety of factors are associated with an increased risk of suicide and attempted suicide, including psychiatric disorders [9-14], feelings of hopelessness and impulsivity [15,16], history of previous suicide attempts [9,17,18], age, sex and race [19-24], marital status [25,26], occupation, comorbidity [28,29], adverse childhood experiences, family history [31,32], and ...

Cited by: 65

Author: Jesús Alberdi-Sudupe, Salvador Pita-Fern...

Publish Year: 2011

Does Compulsory Admission Prevent Inpatient Suicide ...

https://www.researchgate.net/publication/326872290_Does_Compulsory_Admission_Prevent...

The study results suggest that clozapine therapy is associated with a reduced risk of suicide among patients with schizophrenia and schizoaffective disorder.

Suicide among Psychiatric Hospital Inpatients

https://www.researchgate.net/.../225221280_Suicide_among_Psychiatric_Hospital_Inpatients

Empirical investigations support these data; hanging and jumping were the most frequent suicide methods among general hospital patients, patients in psychiatric wards and patients in psychiatric ...