



ALL

IMAGES

VIDEOS

关闭取词

11,300 Results

Any time ▾

### Predictive factors for liver volume and function recovery ...

<https://www.sciencedirect.com/science/article/pii/S1365182X19307336>

Nov 01, 2019 · Recently, **type IV collagen 7s** domain was recognized as a **liver** fibrogenesis marker for predicting postoperative **liver failure** and recurrence of hepatocellular carcinoma. 21, 22 **Liver** fibrosis influences **liver** regeneration; therefore, patients with high **type IV collagen 7s** domain levels in the blood may not regenerate livers as effectively as ...

**Author:** Yutaka Nakano, Osamu Itano, Masah... **Publish Year:** 2019

### Post hepatectomy liver failure – A comprehensive review of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6120608>

Aug 23, 2018 · **Post hepatectomy liver failure** (PHLF) comprises of a conundrum of symptoms and signs following major hepatic resections. The pathophysiology essentially revolves around disruption of the normal hepatocyte regeneration and disturbed **liver** homeostasis. Prompt identification of the pre-operative ...

**Cited by:** 7 **Author:** S. Ray, N.N. Mehta, A. Golhar, S. Nundy

**Publish Year:** 2018

### Post hepatectomy liver failure – A comprehensive review of ...

<https://www.sciencedirect.com/science/article/pii/S2049080118301468>

**Post hepatectomy liver failure** – A comprehensive review of current concepts and controversies ... needs to be directed at preservation of residual hepatocyte function and microvascular organisation and not immediate **recovery** of the total **liver** volume. 4. Risk factors of PHLF ... that have the ability to predict postoperative **liver** function



ALL

IMAGES

VIDEOS

开启歌词

12,000 Results

Any time ▾

## Post hepatectomy liver failure – A comprehensive review of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6120608>

Aug 23, 2018 · Post hepatectomy liver failure (PHLF) comprises of a conundrum of symptoms and signs following major hepatic resections. The pathophysiology essentially revolves around disruption of the normal hepatocyte regeneration and disturbed liver homeostasis. Prompt identification of ...

Cited by: 7

Author: S. Ray, N.N. Mehta, A. Golhar, S. Nundy

Publish Year: 2018

## Predictive factors for liver volume and function recovery ...

<https://www.sciencedirect.com/science/article/pii/S1365182X19307336>

Nov 01, 2019 · Recently, type IV collagen 7s domain was recognized as a liver fibrogenesis marker for predicting postoperative liver failure and recurrence of hepatocellular carcinoma. 21, 22 Liver fibrosis influences liver regeneration; therefore, patients with high type IV collagen 7s domain levels in the blood may not regenerate livers as effectively as ...

Author: Yutaka Nakano, Osamu Itano, Masahi... Publish Year: 2019

## Elevation of Mac-2 binding protein glycosylation isomer ...

<https://onlinelibrary.wiley.com/doi/pdf/10.1002/ags3.12271>

rologic fibrosis markers, 7S domain of type IV collagen (4COL7S) and hyaluronic acid were also measured at SRL. Post-hepatectomy liver failure (PHLF) was diagnosed on the basis of the International Study Group of Liver Surgery defini- tion.15 Briefly, elevated prothrombni tmie – niternatoinal nor -

Author: Daisuke Imai, Takashi Maeda, Huanlin... Publish Year: 2019

## Post hepatectomy liver failure – A comprehensive review of ...

<https://www.sciencedirect.com/science/article/pii/S2049080118301468>

Post hepatectomy liver failure – A comprehensive review of current concepts and controversies ... needs to be directed at preservation of residual hepatocyte function and microvascular organisation and not immediate recovery of the total liver volume. 4. Risk factors of PHLF ... that have the ability to predict postoperative liver function ...

Cited by: 7

Author: S. Ray, N.N. Mehta, A. Golhar, S. Nundy

Publish Year: 2018

## Elevation of Mac-2 binding protein glycosylation isomer ...

<https://onlinelibrary.wiley.com/doi/full/10.1002/ags3.12271>

**Name of Journal:** *World Journal of Gastroenterology*

**Manuscript NO:** 52223

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Cohort Study*

**Pre-hepatectomy type IV collagen 7S predicts post-hepatectomy liver failure and recovery**

Masatsugu Ishii, Osamu Itano, Masahiro Shinoda, Minoru Kitago, Yuta Abe, Taizo Hibi, Hiroshi Yagi, Ayano Takeuchi, Hanako Tsujikawa, Tokiya Abe, and Yuko Kitagawa

**Abstract**

#### Match Overview

1 **Crossref** 20 words  
"Posters (Abstracts 264-2239)", *Hepatology*, 2017 1%