

The diagnostic accuracy and clinical utility of non-English vers



ALL IMAGES VIDEOS

关闭取词

171,000 Results Any time

Postpartum depression in India: a systematic review and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5689195>

Oct 01, 2017 · In 19 studies, the mean age of the study mothers was ≤ 25 years. The **Edinburgh postnatal depression scale** was the most commonly used study instrument (29 studies). The median quality score for the studies was 5 (21 articles had a score of ≤ 5 and 17 had a score > 5).

Cited by: 14 **Author:** Ravi Prakash Upadhyay, Ranadip Chowd...

Publish Year: 2017

Reliability and validity of the Edinburgh Postnatal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4820998>

Apr 04, 2016 · The **Edinburgh Postnatal Depression Scale** (EPDS) is one of the most widely used screening instruments for assessing symptoms of **perinatal depression** and anxiety [3, 4]. It assesses emotional experiences over the past seven days using ten Likert-scale items (See Additional file 1).

Cited by: 35 **Author:** Sumitra Devi Shrestha, Sumitra Devi Shr...

Publish Year: 2016

WHO | Postpartum depression in India: a systematic review ...

<https://www.who.int/bulletin/volumes/95/10/17-192237/en>

Introduction Methods Results Discussion



Postpartum psychiatric disorders can be divided into three categories: postpartum blues; postpartum psychosis and postpartum depression.^{1,2} Postpartum blues, with an incidence of 300–750 per 1000 mothers globally, may resolve in a few days to a week, has few negative sequelae and usually requires only reassurance.¹ Postpartum psychosis, which has a global prevalence ranging from 0.89 to 2.6 per 1000 births, is a ...

Edinbur

The Edinburgh
was develop
of the scale co
guilt feeling, s
ideation. Over
by adding toge
indicate more
weeks postpa
during pregna



Data from: Wikip

Suggest an edit

Name of Journal: *World Journal of Psychiatry*

Manuscript NO: 53540

Manuscript Type: META-ANALYSIS

Diagnostic accuracy and clinical utility of non-English versions of Edinburgh post-natal depression scale for screening post-natal depression in India: A meta-analysis

Paul Swamidhas Sudhakar Russell, Swetha Madhuri Chikkala, Richa Earnest, Shonima Aynipully Viswanathan, Sushila Russell, Priya Mary Mammen

Abstract

Match Overview

1	Internet 38 words crawled on 07-Jan-2018 bmjopen.bmj.com	2%
2	Internet 14 words crawled on 24-Feb-2020 www.dovepress.com	1%

Diagnostic accuracy and clinical utility of non-English versions



YJ



ALL

IMAGES

VIDEOS

218,000 Results

Any time ▾

Reliability and validity of the Edinburgh Postnatal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4820998>

Apr 04, 2016 · The **Edinburgh Postnatal Depression Scale** (EPDS), originally developed in Britain, is one of the most widely used **screening** instruments for assessing symptoms of the Perinatal Common Mental Disorders (PCMDs) of **depression** and anxiety. ... There are three systematic reviews on the validity of **non-English versions** of the EPDS. ... the **accuracy** of ...

Cited by: 54

Author: Sumitra Devi Shrestha, Sumitra Devi Shr...

Publish Year: 2016

Review of the prevalence of postnatal depression across ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6141558>

Jul 20, 2018 · A total of 124 articles concerning research conducted in more than 50 countries were included in the final analysis. There were wide variations in the **screening** instruments and **diagnostic** tools used although the **Edinburgh Postnatal Depression Scale** (EPDS) was the most common instrument applied to identify PND.

Cited by: 2

Author: Siti Roshaidai Mohd Arifin, Helen Cheyne...

Publish Year: 2018

Screening and diagnosing postpartum depression: when and



Diagnostic accuracy and clinical utility of non-English ver



ALL IMAGES VIDEOS

278,000 Results Any time

Reliability and validity of the Edinburgh Postnatal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4820998>

Apr 04, 2016 · The **Edinburgh Postnatal Depression Scale** (EPDS), originally developed in Britain, is one of the most widely used **screening** instruments for assessing symptoms of the Perinatal Common Mental Disorders (PCMDs) of **depression** and anxiety. ... There are three systematic reviews on the validity of **non-English versions** of the EPDS. ... the **accuracy** of ...

Cited by: 54 **Author:** Sumitra Devi Shrestha, Sumitra Devi Shr...
Publish Year: 2016

A systematic review of studies validating the Edinburgh ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/j.1600-0447.2009.01363.x>

Edinburgh Postnatal Depression Scale cut-off points at 9/10 and 12/13 for PPD were used. As described in the Introduction, these are the widely accepted cut-off points to indicate 'possible' and 'probable' **depression** respectively. For studies in antepartum populations, a ...

Cited by: 734 **Author:** J. Gibson, K. McKenzie-McHarg, J. Shak...
Publish Year: 2009

Screening and diagnosing postpartum depression: when and ...

www.scielo.br/scielo.php?script=sci_arttext&pid=S2237-60892017000100054