

26,300 Results Any time

### Preoperative Prediction of Cervical Lymph Node Metastasis ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4670197>

Preoperative Prediction of Cervical Lymph Node Metastasis Using Primary Tumor SUVmax on 18 F-FDG PET/CT in Patients with Papillary Thyroid Carcinoma Ji-hoon Jung , Choon-Young Kim , Seung Hyun Son , Do-Hoon Kim , Shin Young Jeong , Sang-Woo Lee , Jaetae Lee , and Byeong-Cheol Ahn \*

Cited by: 7 Author: Ji-hoon Jung, Choon-Young Kim, Seung ...  
Publish Year: 2015

### Imaging of Gastric Cancer Metabolism Using 18 F-FDG PET/CT

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3996244>

Mar 31, 2014 - Until now, it has been reported that F-18 FDG PET or PET/CT has 21% to 100% of sensitivity and 78% to 100% of specificity for detecting primary tumors.3,4,5,6,7,8,9,10,11 The wide ranges of sensitivity are associated with technical and histopathological factors affecting the visibility of primary tumors on PET/CT (Fig. 1). There can be significant amounts of physiologic FDG uptake in ...

Cited by: 24 Author: Mijin Yun  
Publish Year: 2014

### CT radiomics nomogram for the preoperative prediction of ...

<https://link.springer.com/article/10.1007/s00330-019-06398-z>

Aug 29, 2019 - To investigate the role of computed tomography (CT) radiomics for the preoperative prediction of lymph node (LN) metastasis in gastric cancer. Materials and methods This retrospective study included 247 consecutive patients (training cohort, 197 patients; test cohort, 50 patients) with surgically proven gastric cancer.

Cited by: 2 Author: Yue Wang, Wei Liu, Yang Yu, Jing-juan Li...  
Publish Year: 2019

### Predictive Value of [ 18 F]FDG PET/CT for Lymph Node ...

<https://www.nature.com/articles/s41598-019-41422-8>

Mar 21, 2019 - [18F]Fluorodeoxyglucose ([18F]FDG) Positron emission tomography/computed tomography (PET/CT) is commonly used for rectal cancer staging, but improved diagnostic methods for nodal metastases are ...

Author: Sung Hoon Kim, Bong-Il Song, Beong... Publish Year: 2019  
Author: Sung Hoon Kim

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 53621

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Study*

Nomogram using F-18 FDG PET/CT for preoperative prediction of lymph node metastasis in gastric cancer

Bong-Il Song

**Abstract**

BACKGROUND

Lymph node (LN) metastasis is an important prognostic factor in patients with gastric

## Match Overview

1	<b>Internet</b> 101 words <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	3%
2	<b>Crossref</b> 79 words Sung Hoon Kim, Bong-Il Song, Beong Woo Kim, Hae Won Kim, Kyoung Sook Won, Sung Uk Bae, Woon Kyung Jeon	3%
3	<b>Crossref</b> 37 words Bong-Il Song, Hae Won Kim, Kyoung Sook Won. "Predictive Value of 18F-FDG PET/CT for Axillary Lymph Node Me ...	1%
4	<b>Internet</b> 37 words crawled on 17-Jan-2020 <a href="http://www.dovepress.com">www.dovepress.com</a>	1%
5	<b>Internet</b> 30 words crawled on 03-Dec-2018 <a href="http://link.springer.com">link.springer.com</a>	1%
6	<b>Internet</b> 17 words crawled on 04-Jul-2019 <a href="http://ccforum.biomedcentral.com">ccforum.biomedcentral.com</a>	1%



Nomogram using F-18 fluorodeoxyglucose positron emission



YJ



ALL

IMAGES

VIDEOS

22,800 Results

Any time

### Performance of pre-treatment 18F-fluorodeoxyglucose ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6189439>

Performance of pre-treatment 18 F-fluorodeoxyglucose positron emission tomography/computed tomography ... (such as mesenteric root involvement or suprarenal lymph node metastasis) with a low ... Fujibayashi Y, et al. Incremental benefits of FDG positron emission tomography over CT alone for the preoperative staging of ovarian cancer. AJR Am J ...

Cited by: 1

Author: Sangwon Han, Sungmin Woo, Chong Hy...

Publish Year: 2018

### Preoperative Prediction of Cervical Lymph Node Metastasis ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4670197>

The accuracy of (18)[F]-fluoro-2-deoxy-D-glucose-positron emission tomography/computed tomography, ultrasonography, and enhanced computed tomography alone in the preoperative diagnosis of cervical lymph node metastasis in patients with papillary thyroid carcinoma. World journal of surgery. 2010; 34 (11):2564-9.

Cited by: 7

Author: Ji-hoon Jung, Choon-Young Kim, Seung ...

Publish Year: 2015

### Predictive Value of [ 18 F]FDG PET/CT for Lymph Node



Nomogram using F-18 fluorodeoxyglucose positron emi



ALL IMAGES VIDEOS

22,700 Results Any time

### F-18 fluorodeoxyglucose positron emission tomography and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2933636>

Case presentation. We report a case of a 46-year-old woman of unknown ethnic origin diagnosed with concurrent diffuse large B-cell lymphoma of the breast and squamous cell cancer of the cervix that was detected and followed with F-18 fluorodeoxyglucose (FDG) positron emission tomography and/or computed tomography (PET/CT).

Cited by: 2 Author: Nghi C Nguyen, Christopher N Hueser, A...  
Publish Year: 2010

### Prognostic value of 18-fluorodeoxyglucose positron ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3460334>

Sep 28, 2012 · AIM: To assess the prognostic value of preoperative 18 fluorodeoxyglucose positron emission tomography (FDG-PET)/computed tomography (CT) in patients with resectable colorectal cancer. METHODS: One hundred sixty-three patients with resectable colorectal cancer who underwent FDG-PET/CT before surgery were included.

Cited by: 11 Author: Jang Eun Lee, Sang Woo Kim, Jin Su Ki...  
Publish Year: 2012

### Positron emission tomography with F-18