

Match Overview

1	Crossref 38 words Hong-Yu Chen, Xing Wang, Hao Zhang, Xu-Bao Liu, Chun-Lu Tan. " Mesenterico-portal vein invasion should be an importa	1%
2	Crossref 26 words XueCheng Xie, XiaoChun Zeng, SuJuan Cao, XiaoMao Hu, Qi aoJing Shi, Dan Li, ShiYuan Zhou, Ping Gu, ZhongShan Zhan	1%
3	Internet 19 words crawled on 14-Mar-2020 www.spandidos-publications.com	1%
4	Internet 19 words crawled on 13-Aug-2019 www.tandfonline.com	1%
5	Internet 12 words crawled on 12-Jun-2019 patents.justia.com	<1%
6	Internet 10 words crawled on 22-Jan-2020 www.jcancer.org	<1%
7	Internet 10 words crawled on 23-Jan-2020 wjso.biomedcentral.com	<1%

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 53637

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Functionality is not an independent prognostic factor for pancreatic neuroendocrine tumor

Hong-Yu Chen, Ya-Liang Zhou, Yong-Hua ¹Chen, Xing Wang, Hao Zhang, Neng-Wen Ke, Xu-Bao Liu, Chun-Lu Tan



国内版

国际版

Chat with Bing

Functionality is not an independent prognostic factor for



Sign in



ALL

IMAGES

VIDEOS

2,010 Results

Any time ▾

Non-functional neuroendocrine tumors of the pancreas ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4548112>

Aug 28, 2015 · Core tip: **Pancreatic neuroendocrine tumors (PNETs)** are a fascinating and diverse group of **neoplasms**. While the clinical features of **functional PNETs** are frequently discussed, the majority of **PNETs** are actually **non-functional**. Although typically slow growing, tumors with more aggressive biology may progress to unresectable metastatic disease.

Cited by: 31

Author: Jordan M Cloyd, George A Poultides

Publish Year: 2015

Pancreatic neuroendocrine tumors (PNETs): incidence ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2735065>

May 30, 2008 · In conclusion, this large population-based study using SEER data provides an up-to-date estimate of the **incidence and prognosis of PNETs** and suggests that the **functional status** of the **tumor** may be an **independent prognostic factor** after ...

Cited by: 641

Author: Thorvardur R. Halfdanarson, K. G. Rabe, ...

Publish Year: 2008

Pancreatic Neuroendocrine Tumor Risk Factors

American Cancer Society

Changes in the genes that cause some of these syndromes can be found by genetic testing. For more information on genetic testing, see Can Pancreatic Neuroendocrine Tumor Be Found Early? Diabetes. Pancreatic NETs are more common in people with diabetes. The reason for this is not known. Most of the risk is found in people with type 2 diabetes.



Instagram

Address: 250 Williams Street NW, Atlanta, GA 30303, United States

ContactType: customer service

Telephone: [+1-404-320-3333](tel:+14043203333)

Pancreatic Neuroendocrine Tumor Risk Factors

https://www.cancer.org/cancer/pancreatic-neuroendocrine-tumor/causes_risks_prevention/risk_factors.html

Functionality is not an independent prognostic factor for



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

23,300 Results

Any time ▾

[Liver Tumor Burden in Pancreatic Neuroendocrine Tumors: CT ...](#)

<https://www.mdpi.com> › [2072-6694](#) › [12](#) › [6](#) › [1486](#) › [htm](#)

1 day ago · **Pancreatic neuroendocrine tumors** (p-NETs) are a rare group of neoplasms that often present with liver metastases. Histological characteristics, metabolic behavior, and liver tumor burden (LTB) are important **prognostic factors**. In this study, the usefulness of texture analysis of liver metastases in evaluating the biological aggressiveness of p-NETs was assessed.

[\[PDF\] Pancreatic Neuroendocrine Tumors \(Islet Cell Tumors ...](#)

<https://sfsurgery.com> › [wp-content](#) › [uploads](#) › [2014](#) › [06](#) › ...

Surgery plays a role even in the setting of metastatic disease. The symptoms of metastatic **functional pancreatic neuroendocrine tumors** (NETs) may be ameliorated by the reduction of overall tumor burden through surgical debulking. The liver is a common site of metastasis from **pancreatic** NETs. Because of the slow growth rate of many

[Frailty Questionnaire Is Not a Strong Prognostic Factor ...](#)

<https://www.ncbi.nlm.nih.gov> › [pmc](#) › [articles](#) › [PMC6362511](#)

Feb 04, 2019 · For **functional** recovery after arthroplasty surgery, we have now shown that GFI is **not** a strong **prognostic factor**. We found that the **functional** baseline score is a strong **prognostic** score which can fairly well discriminate between good and adverse **functional** outcomes.

Author: Jennifer M. T. A. Meessen, Marta Fioc... **Publish Year:** 2019

[Neuroendocrine Differentiation - an overview ...](#)

<https://www.sciencedirect.com> › [topics](#) › [medicine-and...](#)

No **prognostic role** was found in the first study, whereas the subsequent larger two, including ours, found only a marginal **prognostic role** that was not sufficient to be useful clinically. 114-116 In the single study analyzing **neuroendocrine differentiation** in **prostate cancer** on needle biopsy, there was no relationship with **prognosis**. 117 According to the CAP Consensus Statement 1999, 118 **neuroendocrine differentiation** is still considered a category III **prognostic factor not** ...

[Frailty Questionnaire Is Not a Strong Prognostic Factor](#)