

World Journal of *Gastroenterology*

World J Gastroenterol 2020 May 14; 26(18): 2126-2275



**OPINION REVIEW**

- 2126** Annexin A2 promotion of hepatocellular carcinoma tumorigenesis *via* the immune microenvironment
Qiu LW, Liu YF, Cao XQ, Wang Y, Cui XH, Ye X, Huang SW, Xie HJ, Zhang HJ

REVIEW

- 2138** Medical management of metabolic and cardiovascular complications after liver transplantation
Becchetti C, Dirchwolf M, Banz V, Dufour JF
- 2155** Natural products that target macrophages in treating non-alcoholic steatohepatitis
Li CL, Zhou WJ, Ji G, Zhang L

MINIREVIEWS

- 2166** Hepatitis B virus recurrence after liver transplantation: An old tale or a clear and present danger?
Lenci I, Milana M, Grassi G, Manzia TM, Gazia C, Tisone G, Angelico R, Baiocchi L
- 2177** New role for ceramide in hypoxia and insulin resistance
Xia QS, Lu FE, Wu F, Huang ZY, Dong H, Xu LJ, Gong J
- 2187** Role of gut microbiota on intestinal barrier function in acute pancreatitis
Li XY, He C, Zhu Y, Lu NH

ORIGINAL ARTICLE**Basic Study**

- 2194** Genetic association analysis of *CLEC5A* and *CLEC7A* gene single-nucleotide polymorphisms and Crohn's disease
Elleisy N, Rohde S, Huth A, Gittel N, Glass A, Möller S, Lamprecht G, Schäffler H, Jaster R
- 2203** Folic acid attenuates high-fat diet-induced steatohepatitis *via* deacetylase SIRT1-dependent restoration of PPAR α
Xin FZ, Zhao ZH, Zhang RN, Pan Q, Gong ZZ, Sun C, Fan JG

Retrospective Study

- 2221** Prognostic significance of hepatic encephalopathy in patients with cirrhosis treated with current standards of care
Bohra A, Worland T, Hui S, Terbah R, Farrell A, Robertson M
- 2232** Optimal proximal resection margin distance for gastrectomy in advanced gastric cancer
Kim A, Kim BS, Yook JH, Kim BS

META-ANALYSIS

- 2247 Computed tomography *vs* liver stiffness measurement and magnetic resonance imaging in evaluating esophageal varices in cirrhotic patients: A systematic review and meta-analysis
Li Y, Li L, Weng HL, Liebe R, Ding HG

CASE REPORT

- 2268 Ectopic hepatocellular carcinoma mimicking a retroperitoneal tumor: A case report
Adachi Y, Hayashi H, Yusa T, Takematsu T, Matsumura K, Higashi T, Yamamura K, Yamao T, Imai K, Yamashita Y, Baba H

ABOUT COVER

Associate Editor of *World Journal of Gastroenterology*, Mark De Ridder, MD, PhD, Chairman, Full Professor, Department of Radiotherapy, UZ Brussel, Vrije Universiteit Brussel, Brussels 1090, Belgium

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2019 edition of Journal Citation Report® cites the 2018 impact factor for WJG as 3.411 (5-year impact factor: 3.579), ranking WJG as 35th among 84 journals in gastroenterology and hepatology (quartile in category Q2). CiteScore (2018): 3.43.

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Yan-Liang Zhang*

Proofing Production Department Director: *Yun-Xiaojuan Wu*

Responsible Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Subrata Ghosh, Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

May 14, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Hepatitis B virus recurrence after liver transplantation: An old tale or a clear and present danger?

Ilaria Lenci, Martina Milana, Giuseppe Grassi, Tommaso M Manzia, Carlo Gazia, Giuseppe Tisone, Roberta Angelico, Leonardo Baiocchi

ORCID number: Ilaria Lenci (0000-0001-5704-9890); Martina Milana (0000-0003-2027-0481); Giuseppe Grassi (0000-0001-9182-8759); Tommaso M Manzia (0000-0002-4636-3478); Carlo Gazia (0000-0002-3543-4170); Giuseppe Tisone (0000-0001-8860-5909); Roberta Angelico (0000-0002-3439-7750); Leonardo Baiocchi (0000-0003-3672-4505).

Author contributions: Lenci I made the analysis and interpretation, and drafted the manuscript; Lenci I, Milana M, Grassi G, Manzia TM, Gazia C, Tisone G, and Angelico R made the acquisition of data and critical revision; Baiocchi L made the proposal of study, study conception, correction of manuscript and critical revision.

Conflict-of-interest statement: No conflict of interest to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Ilaria Lenci, Martina Milana, Giuseppe Grassi, Leonardo Baiocchi, Hepatology and Liver Transplant Unit, University of Tor Vergata, Rome 00133, Italy

Tommaso M Manzia, Carlo Gazia, Giuseppe Tisone, Roberta Angelico, Hepato-Pancreato-Biliary and Transplant, Department of Surgery, University of Rome Tor Vergata, Rome 00133, Italy

Corresponding author: Leonardo Baiocchi, MD, PhD, Assistant Professor, Hepatology and Liver Transplant Unit, University of Tor Vergata, Viale Oxford 81, Rome 00133, Italy. baiochi@uniroma2.it

Abstract

Hepatitis B virus (HBV) recurrence after liver transplantation (LT) has been described more than 50 years ago. Similarly, to other clinical conditions, in which impairment of host immune defense favors viral replication, early reports described in details recurrence and reactivation of HBV in liver transplant recipients. The evidence of a possible, severe, clinical evolution of HBV reappearance in a significant percentage of these patients, allowed to consider, for some years, HBV positivity a contraindication for LT. Moving from the old to the new millennium this picture has changed dramatically. Several studies contributed to establish efficient prophylactic protocols for HBV recurrence and with the advent of more potent anti-viral drugs an increased control of infection was achieved in transplanted patients as well as in the general immune-competent HBV population. Success obtained in the last decade led some authors to the conclusion that HBV is now to consider just as a “mere nuisance”. However, with regard to HBV and LT, outstanding issues are still on the table: (1) A standard HBV prophylaxis protocol after transplant has not yet been clearly defined; (2) The evidence of HBV resistant strains to the most potent antiviral agents is claiming for a new generation of drugs; and (3) The possibility of prophylaxis withdrawal in some patients has been demonstrated, but reliable methods for their selection are still lacking. The evolution of LT for HBV is examined in detail in this review together with the description of the strategies adopted to prevent HBV recurrence and their pros and cons.

Key words: Liver transplant; Hepatitis B virus; Viral recurrence; Prophylaxis; Minimization; Antiviral drug

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Received: February 11, 2020
Peer-review started: February 11, 2020
First decision: March 6, 2020
Revised: March 12, 2020
Accepted: April 30, 2020
Article in press: April 30, 2020
Published online: May 14, 2020

P-Reviewer: Akamatsu N, Ho CM, Zheng H
S-Editor: Dou Y
L-Editor: A
E-Editor: Ma YJ



Core tip: Liver transplantation for hepatitis B virus (HBV) has greatly evolved in the last 50 years. Several studies contributed to establish efficient prophylactic protocols for HBV recurrence and with the advent of more potent anti-viral drugs an increased control of infection was achieved. In this review we examined in detail the results obtained in preventing HBV reappearance in liver transplanted patients and the possible future directions of research in this field.

Citation: Lenci I, Milana M, Grassi G, Manzia TM, Gazia C, Tisone G, Angelico R, Baiocchi L. Hepatitis B virus recurrence after liver transplantation: An old tale or a clear and present danger? *World J Gastroenterol* 2020; 26(18): 2166-2176

URL: <https://www.wjgnet.com/1007-9327/full/v26/i18/2166.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v26.i18.2166>

INTRODUCTION

The hepatitis B virus (HBV) is a small DNA virus belonging to the *Hepadnaviridae* family^[1]. Despite the adoption, in several countries, of an extended vaccination campaign starting in 1992, HBV infection still represents an important health problem with 350-400 million people infected in the world^[2]. Without treatment, at least one third of patients are estimated to progress to significant liver disease, including end-stage liver cirrhosis and tumors. In fact, the natural history of HBV liver disease includes a spectrum of clinical conditions ranging from a non-frequent fulminant hepatitis to HBV-related hepatocellular carcinoma and/or end-stage liver disease^[3,4]. While vaccination and the new antiviral drugs are effective, respectively, in avoiding HBV infection and preventing the most severe sequelae of HBV disease, liver transplantation (LT) remains the main therapeutic option in patients with more severe forms of HBV liver injury^[5]. However, in the early 90's the possibility to offer LT to HBV candidates was an argument of debate. In fact, it was evident that HBV disease recurrence in the graft was severe in a significant proportion of patients^[6]. Moreover, an aggressive clinical form of viral reactivation, named Fibrosing Cholestatic Hepatitis, was also described in nearly 25% of HBV transplanted patients, leading to a dramatic and rapidly progressive course^[7]. Therapeutic advancement and prophylactic strategies against HBV radically changed this picture in the last three decades, allowing the consideration of HBV recurrence after LT to no longer be of concern. In this review we will describe HBV viral features, its natural history, and current outcome of HBV after LT.

NATURAL HISTORY OF HBV

HBV, a double stranded small DNA virus, replicating by reverse transcription, is able to convert its DNA in a covalently closed circular (ccc) form when reaching the hepatocyte's nucleus. cccDNA represents a mini-chromosome containing information for antigens (HBsAg, HBeAg, and HBcAg), X protein, and polymerase production^[8]. The infection route is mainly represented by vertical transmission in endemic areas. The estimated risk of acquiring the infection from an HBeAg+ mother is around 80%^[9]. On the other hand, sexual or needle transmission are important paths in non-vaccinated adult patients of western countries^[10]. Evolution of infection is dependent on the host, the viral genetics and virus/host interaction^[3,11,12]. Vertical transmission at birth is associated (without peri-natal treatment) with a lifetime infection, usually with an immune-tolerant state^[11]. This clinical situation is characterized by HBeAg positivity, high levels of HBV-DNA and normal liver function tests. Conversely, in adult normal subjects, immuno-tolerance usually lasts for 2-4 wk, the time span corresponding to the HBV incubation phase. Activation of the immune system against HBV determines: (1) Decreased HBV-DNA levels; (2) Increased liver inflammation; and (3) Elevation of serum levels of liver function tests. These features characterize the immune-active phase. This stage may evolve into: (1) Infection resolution with production of high titers of HBsAb (this target is reached by more than 90% of healthy adult individuals within 6 mo of initial HBV contact); (2) Fulminant hepatitis (rarely, ≤ 0.5%); or (3) HBsAg persistence and evolution to chronic hepatitis^[4]. During chronic hepatitis, the seroconversion to the HBeAg negative state (with development of HBeAb titer) represents an important achievement as it corresponds to decreased

levels of HBV-DNA, liver inflammation and injury^[13-15]. Moreover, HBeAg seroconversion with the consequent drop in HBV-DNA serum levels has been related to reduced fibrosis progression, histological staging, and onset of cirrhosis and hepatocellular carcinoma^[16-18]. A subject with an acquired HBeAg negative state is usually defined as an inactive HBV carrier, referring to a remission state of the liver disease. Unfortunately, seroreversion to an HBeAg positive condition may occur over time (in approximately 20% of patients), also transiently, leading to a “*de novo*” immune-active inflammatory stage. Moreover, HBeAg loss (both spontaneous and drug induced) may determine selection in the host of pre-core mutants of HBV (not producing HBeAg). These strains are not affected, during their replicative phases, by anti-HBe antibodies, thus they determine progression of liver injury in approximately 10 to 30% of patients obtaining HBeAg loss^[3,5,19]. The main clinical and virological features, in the different phases of HBV chronic infection, are reported in [Table 1](#). From the above, it is evident that host-immune-system/virus interaction is a major determinant of the presence and severity of liver injury. This result is far more evident in subjects undergoing immune system changes related to biological or immuno-suppressive therapies, including the majority of transplanted patients. In this setting severe reactivation of HBV is an element of concern^[20].

HBV DURING IMMUNE SYSTEM SUPPRESSION OR MODIFICATION (DRUG-INDUCED IMPAIRMENT OF THE IMMUNE SYSTEM AS A RISK FACTOR FOR HBV REACTIVATION)

Reactivation of HBV is represented by sudden reappearance or increase of viral DNA in the serum of a patient with a resolved or clinically silent HBV infection^[20]. This condition, which may also occur spontaneously, has been reported more frequently in patients undergoing immunosuppressive therapy for malignant or non-malignant disease^[21]. In an early study in non-Hodgkin lymphoma patients under chemotherapy, HBV reactivation accounted for 72% of cases in HBsAg positive subjects^[22]. More importantly, viral reappearance was also observed in HBsAb/HBcAb or only HBcAb positive subjects, thus suggesting the possibility of HBV reactivation also in conditions in which the infection was considered resolved in the past. Further studies also demonstrated HBV reappearance in non-neoplastic clinical settings such as Crohn's disease^[23] or rheumatologic affections^[24]. Treatment with biological agents, such as B-cell depleting (*i.e.*, rituximab) or anti-tumor necrosis factor drugs (infliximab), carries a significant risk of HBV reactivation^[25]. However, standard steroid treatment may also be responsible for HBV reappearance^[26]. Evolution of viral reactivation is generally thought to occur in three separate phases^[20,21]. At the beginning, a rise in HBV-DNA (at least ten-fold in comparison with baseline values) is observed during immunosuppressant treatment. In the second phase, when drugs are tapered or discontinued, the inflammatory damage begins, being triggered by the host immune defense that is also, in part, restored. In the last phase, the liver damage is repaired or may progress to end-stage liver failure. The evidence of a possible dramatic evolution of HBV reactivation in liver failure prompted the adoption of strategies to counteract this preventable occurrence. First of all, an adequate screening for HBV virus, including HBcAb, is proposed in individuals undergoing chronic therapy with immunologic modifiers. Secondly, antiviral agents able to prevent or cure this clinical condition are administered according to both viral and patient's features^[27-29]. However, the clinical strategies commonly employed to prevent HBV reinfection are still lacking significant scientific evidence^[27]. Therefore, the question of the best approach in different clinical scenarios remains open.

Finally, the most important clinical setting in which HBV reappearance is a relevant issue is that of transplant. Transplanted patients usually require long-term high-dose immunosuppression to prevent rejection. In HBsAg positive patients undergoing bone marrow transplantation, HBV reactivation accounts for nearly the totality of cases^[30]. Even in HBsAb/HBcAb+ subjects, reappearance of active HBV is not rare, accounting for nearly 20% of cases^[31]. Starting from the early eighties, HBV reactivation was reported to be very frequent in the setting of kidney and heart transplantation, and it was characterized by the insurgence of HBV chronic hepatitis^[20]. Indeed, HBV reactivation or recurrence also represents an important issue in liver transplanted patients. These subjects, in fact, share the same immuno-suppressive need as other transplanted patients but, at the same time, are suffering the most important sequelae of HBV before surgery. This setting probably represents the most important clinical scenario in which dramatic HBV resurgence was observed

Table 1 Main virological and biochemical features in the different clinical phases of chronic hepatitis B in HBsAg+ patients

Phase 1: Immune tolerant	Phase 2: Immune active	Phase 3: Asymptomatic carrier	Phase 4: HBV reactivation
HBeAg+	Attempt to remove HBeAg	HBeAg-	HBeAg+ or -
HBV-DNA $\geq 10^8$ IU/mL	HBV-DNA \downarrow	HBV-DNA < 2000 IU/mL	HBV-DNA \uparrow
Normal LFTs \rightarrow no damage	Altered LFTs \rightarrow damage	Normal LFTs \rightarrow no damage	Altered LFTs \rightarrow damage

HBV: Hepatitis B virus; LFT: liver function tests

and prophylactic measures were firstly pursued. At the same time, LT was the setting in which the risk of transplant with anti-core-HBV positive liver graft was identified.

HBV RECURRENCE IN THE EARLY TIMES OF LIVER TRANSPLANTATION (THE PURSUIT OF AN EFFECTIVE PROPHYLACTIC STRATEGY)

HBV recurrence/reactivation after LT was already recognized almost 50 years ago^[32]. In the early 90's the feasibility of LT in HBV patients remained a crucial question since several reports observed viral recurrence in nearly all transplanted subjects, with an aggressive course in the larger part of them. While graft replacement was able to transiently reduce viral load, viral resurgence in the course of immunosuppression was related to significant liver damage and cirrhosis development^[6,33]. So, at that time, LT in HBsAg positive patients was considered a high risk procedure for graft and patient loss, with an unacceptable hazard in particular in HBeAg+ subjects^[34]. The disappointing results, and the need to pursue a solution for HBV patients with end-stage liver disease, stimulated the research for a possible prophylactic therapy after LT. In a pioneering study conducted at Paul Brousse Hospital (Villejuif, France) in the eighties, an extended passive immune-prophylaxis was tested in HBsAg positive patients after LT^[35]. Despite the monthly HBsAb immunoglobulin (HBIG) administration, 29% of patients experienced HBsAg and HBV-DNA reappearance in serum, however these data demonstrated the possibility to reduce HBV recurrence after LT. In a further European retrospective study on 372 HBV liver transplanted patients (between 1977 and 1990), a reduced rate of HBV reactivation was statistically associated with the absence of HBV-DNA before transplant and again to long-term passive immune prophylaxis with HBIG^[36]. The exact mechanisms of the beneficial effects of immunoglobulin in this setting are not completely clear at present. Both reduced deletion of infected hepatocytes and prevention of viral aggression of liver cells have been suggested as possible effects^[37]. Starting from the mid 90's, evidence was gathered on the role of lamivudine (Lam) treatment in repressing HBV replication^[38,39]. Since, at that time, only HBIG-based prophylaxis was available after LT, and this therapy was a life-long, suboptimal, expensive treatment, the evaluation of the Lam effect in this clinical setting began. In an English study, 17 HBsAg positive patients were enrolled to receive Lam 4 wk before liver transplant and to continue 1 year thereafter^[40]. Twelve out of seventeen patients were transplanted. In them, Lam induced a loss of HBsAg and undetectable HBV-DNA serum levels within 4 wk of treatment and after transplant. Moreover, liver histology did not show features suggesting HBV recurrence after LT, and these results were obtained without concomitant HBIG immune prophylaxis. Unfortunately, in the same study, selection of a resistant strain to Lam was observed in one patient after 20 wk of treatment. This occurrence was characterized by reactivation of HBV and evidence of chronic hepatitis on liver tissue after 1 year. Similar to that observed in HIV therapy^[41], HBV strains not-responding to Lam were characterized by mutation of polymerase at the highly conserved YMDD motif^[42-44]. With regard to liver transplanted HBV patients, extended follow up of Lam resistant patients was lately reported. Resistance to Lam began to occur, typically, six months after its introduction and was sometimes characterized by severe disease recurrence^[45,46]. A combination of Lam therapy with HBIG was then attempted in order to further reduce HBV recurrence after LT. In a study, fourteen HBsAg positive LT patients were treated with Lam plus HBIG^[47]. In a median follow-up of one year, all patients were HBV-DNA negative in serum, thus demonstrating the superiority of combination therapy in comparison with monotherapy with either Lam or HBIG. These data were also confirmed in a study with an extended (average 31 mo) follow-up^[48]. Thus, the past millennium ended with the positive perspective that prevention of HBV recurrence/reactivation in HBsAg

transplanted subjects was feasible. On the basis of these results, the possible exclusion of HBV subjects from transplant lists was largely reexamined.

HBV RECURRENCE/REACTIVATION AFTER LIVER TRANSPLANTATION IN THE THIRD MILLENNIUM (TESTING NEW THERAPEUTIC APPROACHES AND DRUGS)

The efficacy of passive immunization, in association with Lam, was again demonstrated in retrospective studies after the year 2000^[49,50]. However, since this strategy was flawed by the relevant cost of HBIG and the need of life-long administration, the possibility to induce active immunization in HBV liver transplanted patients was examined.

HBV vaccination

In a study on 17 HBsAg+, HBeAg and HBV-DNA negative liver transplanted patients (after at least 18 mo of HBIG treatment), the double dose administration of HBV vaccine at baseline, 1 and 6 mo was tested^[51]. After vaccination 84% of patients developed an HBsAb titer. During a further follow-up of 14 mo, HBsAg reappearance was not observed. These positive results were not replicated in a following study in which three reinforced and sequential cycles of HBV vaccination determined only a 17.6% HBsAb seroconversion in HBV transplanted patients^[52]. In an editorial in the same journal, the limits of this strategy in transplanted patients were discussed, underscoring the scarce vaccine efficacy during immunosuppression and the long time needed to reconstitute the immune system after its depression^[53]. In conclusion, it was confirmed that HBIG and antiviral therapy were regarded as the most appropriate measures against HBV recurrence after LT^[54].

Adefovir dipivoxil

With regard to antiviral agents, in those years, a new drug implemented the armamentarium for the therapy of HBV. Adefovir dipivoxil (ADV), a nucleotide analog inhibiting viral reverse transcriptase that was abandoned for treatment of HIV because of kidney damage when used at high dose, was licensed for HBV treatment since it was active at lower, non-toxic levels for this virus (10 mg/d). ADV treatment in the majority of immune-competent HBsAg patients (both HBeAg positive or negative) determined a clear reduction of HBV-DNA, improvement of liver histology, and normalization of liver enzymes after a 48 wk course^[55,56]. Moreover, emergence of ADV resistant mutants was not observed during these trials. Despite the fact that possible long-term viral resistance to ADV remained to be assessed, the efficacy of this new antiviral drug allowed hope for a new era in which HBV could be regarded as just a “mere nuisance”^[57]. Soon, ADV was employed for the treatment of Lam resistant HBV after transplant^[58]. Again, a significant improvement of liver function was recorded in nearly 90% of patients, and no resistant HBV strains were selected after 48 wk of therapy. However, ADV viral resistance was then observed with prolonged follow-up^[59,60]. This was characterized by a novel N236T mutation of HBV polymerase. In spite of this, the clinical evolution in patients was not worrisome since these ADV resistant strains were easily suppressed by Lam concomitant therapy. On the base of these findings, a possible Lam + ADV concomitant treatment for HBV was suggested^[61]. Data from a systematic review including 2162 HBV LT patients^[62] identified the following as possible risk factors for HBV recurrence: (1) Being HBV-DNA positive at transplant (8.5% *vs* 4%); (2) Administration of low dose HBIG in the first week after LT (6.1% *vs* 3.5%); and (3) Combination therapy with HBIG + Lam *vs* HBIG + ADV (6.1% *vs* 2%). This picture was destined to undergo further changes with the advent of new nucleos(t)ide analogues with high genetic barriers.

New high genetic barrier nucleos(t)ide analogues.

Starting from 2012, entecavir (ETV) and tenofovir dipivoxyl (TDF) were proposed by several guidelines as a first line of treatment for chronic HBV hepatitis^[5,63]. In fact, both drugs were demonstrated to be very effective in clinical studies, to have an excellent safety profile, and to be affected by a minimal or absent emergence of resistant HBV strains^[64-68]. In a systematic review^[69] on nucleos(t)ide analogues for HBV prophylaxis after LT, the comparison between Lam + HBIG *vs* the association of ETV or TDF with HBIG demonstrated the superiority of the latter treatments (HBV recurrence rate 6.1% *vs* 1%, $P < 0.001$). Moreover, in the same analysis, preliminary data evidenced slightly better results with either ETV or TDF monotherapy (after HBIG discontinuation) in comparison with the canonical Lam + HBIG prophylaxis (HBV recurrence rate 3.9% *vs* 6.1%, difference not statistically significant). These findings introduced the concept of

a possible minimization of HBV prophylaxis after LT, stimulating research with this target.

TOWARD HBV PROPHYLAXIS MINIMIZATION AFTER LT

Several strategies have been designed to minimize HBV prophylaxis after LT. The most relevant are described in the following subparagraphs with the corresponding results. Main studies on this issue are also summarized in [Table 2](#).

HBIG dose reduction

Since long-term administration of HBIG was a critical point for its high cost, that would easily reach \$100.000/pts/year^[70], several attempts were carried out to reduce HBIG administration and acceptable results obtained. In a 2004 study conducted in our Unit (Liver Transplant Center, University of Rome Tor Vergata), we evaluated the possibility to prevent HBV recurrence after LT by administering HBIG on demand (when HBsAg serum levels were ≤ 70 IU/L) instead of the standard monthly administration^[71]. Moreover, in the same study, two different HBIG doses (5000 IU or 2000 IU) were employed. In eleven HBV patients, at low risk for reactivation (HBsAg, HBV-DNA negative) and under concomitant Lam therapy, this strategy did not determine any HBV reactivation for 1 year follow up. On the other hand, the treatment based on administration of 2000 IU HBIG on demand reduced the cost of passive immune-prophylaxis by more than 50%. In 2007, the Australasian Liver Transplant Study Group assessed the association of very-low HBIG doses (400-800 IU) + Lam on HBV recurrence after LT^[72]. This strategy accounted for a modest HBV recurrence risk of 4% in 5 years, and the results were considered highly satisfactory since the majority of patients (85%) were HBV-DNA positive at transplant.

High-genetic barrier nucleos(t)ide analogues monotherapy

The advent of high-genetic barrier nucleos(t)ide analogues ETV and TDF, allowed speculation on a possible prophylaxis without HBIG. ETV monotherapy, tested on 80 patients undergoing LT for HBV, was able to suppress HBV-DNA (under the lower detection limit) in nearly 99% of cases after 24 mo^[73]. Extended follow up of this study (8 years) demonstrated a 92% loss of HBsAg, while HBV-DNA was undetectable in all^[74]. On the other hand, discontinuation of HBIG in transplanted patients treated with TDF + HBIG did not change any viral or patient profile in a 72 wk follow-up^[75]. Good results with either ETV and TDF were also replicated in other studies^[76,77]. In a 5-year follow up in patients discontinuing HBIG and commencing either ETV or TDF after LT, HBsAg+ seroconversion occurred in 8% of cases, while HBV-DNA reappearance was not observed^[78]. On the basis of these results, the most authoritative guidelines in the field now contemplate ETV or TDF monotherapy as an efficient prophylactic measure in subjects at low risk of HBV recurrence after LT^[78,79].

Complete withdrawal of HBV prophylaxis

In the past years, our group examined a more radical approach to HBV prophylaxis minimization. This was characterized by the complete withdrawal of antiviral drugs in well selected HBV transplanted patients. We started with the assumption that reappearance of HBV after transplantation was dependent on the presence of cccDNA in the graft. Contrary to a North American study, (including several HBeAg/HBV-DNA+ patients at LT) in which total HBV-DNA and cccDNA were detected in liver tissue in 83% and 18% of cases, respectively^[80], in a preliminary evaluation of HBsAg patients transplanted in our center, only 1 out of 44 was found to be positive for cccDNA^[81]. Among those that were negative for liver cccDNA, 30 were selected and underwent sequential withdrawal of HBIG and Lam. The majority of patients (83%, $n = 25$) did not experience any HBV recurrence in a median follow up longer than 2 years. Five patients came back to an HBsAg positive status. Prompt resumption of HBV prophylaxis allowed infection control, avoiding any significant clinical impairment^[82]. From this study, we concluded that complete withdrawal of HBV prophylaxis after LT was feasible in patients with negative serum HBV-DNA and tissue cccDNA at transplant. An editorial, in the same journal, wisely observed that the time had come for an individualized prophylaxis in HBV transplanted patients^[83]. In fact, recurrence of HBV was mainly reported in patients who were HBV-DNA positive at transplant (> 100.000 copies/mL) and/or HBeAg+^[36,72,84]. On the other hand, those not falling in the above category were considered at low risk for HBV recurrence. In this perspective, the target of HBV-DNA negativity was to be pursued before transplant in order to perhaps minimize prophylaxis after grafting. Conversely, for high-risk patients (HBeAg, HBV-DNA positive), more robust antiviral protocols were to be considered.

Table 2 Main studies examining prophylaxis minimization in hepatitis B virus liver transplanted patients, employing different approaches

Specific aim	Ref.	Number of patients	Method	Main results
HBIG minimization	Di Paolo <i>et al</i> ^[71]	11	HBIG administration on demand (when HBsAb < 70 IU/L) with Lam	No HBV reactivation (1 yr F/U)
	Gane <i>et al</i> ^[72]	147	Very-low HBIG dose (400-800 IU monthly) with Lam	4% of HBV recurrence (5 yr F/U)
High-genetic barrier nucleos(t)ide analogues monotherapy	Fung <i>et al</i> ^[73,74]	80	ETV monotherapy	92% HBsAg-100%HBV-DNA undetectable (8 yr F/U)
	Teperman <i>et al</i> ^[75]	40	TDF monotherapy after HBIG discontinuation	No change (72 wk F/U)
	Manini <i>et al</i> ^[77]	77	ETV or TDF monotherapy after HBIG discontinuation	100%HBV-DNA undetectable 9% HBsAg reappearance (5 yr F/U)
Complete withdrawal of HBV prophylaxis	Lenci <i>et al</i> ^[81,82]	30	Sequential discontinuation of HBIG and Lam in low risk (cccDNA negative) patients	90% successful withdrawal 60% HBsAb > 10 IU (6 yr F/U)

cccDNA: Covalently closed circular DNA; HBV: Hepatitis B virus; ETV: Entecavir; F/U: Follow-up; HBIG: HBsAb immunoglobulin; Lam: Lamivudine; TDF: Tenofovir dipivoxyl.

More recently, data on longer (6-year) follow up of this original cohort were published by our group^[85]. Only 3 patients needed prophylaxis resumption (10%). Of the whole cohort, 93% remained HBsAg negative and 100% had undetectable HBV-DNA. More interestingly, 60% of patients spontaneously developed an HBsAb titer > 10 IU/L. This was probably related to the minimization or withdrawal of anti-rejection therapy that is routinely pursued in our center in patients transplanted for several years. Comment on this study appeared in a new editorial^[86]. While these data were encouraging, it underscored that limits remained in the identification of low risk patients. cccDNA techniques, in fact, needed to be standardized to be widely and consistently applicable in clinical settings, but on the other hand, extra-hepatic HBV reservoirs were still a possible issue of concern.

CONCLUSION

Several important achievements were obtained in the last fifty years with regard to HBV liver transplanted patients. The original exclusion of these subjects from LT waiting lists, due to poor outcome, was counteracted by the adoption of effective measures to prevent HBV recurrence. At present, high genetic barrier anti-viral drugs are giving an important contribution in transplanted patients, as well as in the HBV immune-competent population. Recently, tenofovir alafenamide, a TDF analog with improved renal safety and increased ability to reduce alanine aminotransferase, was employed in LT patients with good results^[87]. However, of concern, HBV mutants with resistance to TDF (the drug with the highest genetic barrier) were recently identified, underscoring the need of a new generation of HBV agents to be employed, at least, as a rescue therapy^[88]. The future of LT for HBV is not completely predictable at this stage. It will, however, depend on the global burden of HBV and the possible discovery of HBV eradicating drugs.

REFERENCES

1. Liang TJ. Hepatitis B: the virus and disease. *Hepatology* 2009; **49**: S13-S21 [PMID: 19399811 DOI: 10.1002/hep.22881]
2. Wang FS, Fan JG, Zhang Z, Gao B, Wang HY. The global burden of liver disease: the major impact of China. *Hepatology* 2014; **60**: 2099-2108 [PMID: 25164003 DOI: 10.1002/hep.27406]
3. Lin CL, Kao JH. Natural history of acute and chronic hepatitis B: The role of HBV genotypes and mutants. *Best Pract Res Clin Gastroenterol* 2017; **31**: 249-255 [PMID: 28774406 DOI: 10.1016/j.bpg.2017.04.010]
4. Fattovich G. Natural history of hepatitis B. *J Hepatol* 2003; **39** Suppl 1: S50-S58 [PMID: 14708678 DOI: 10.1016/s0168-8278(03)00139-9]
5. Terrault NA, Bzowej NH, Chang KM, Hwang JP, Jonas MM, Murad MH; American Association for the Study of Liver Diseases. AASLD guidelines for treatment of chronic hepatitis B. *Hepatology* 2016; **63**: 261-283 [PMID: 26566064 DOI: 10.1002/hep.28156]

- 6 **Demetris AJ**, Todo S, Van Thiel DH, Fung JJ, Iwaki Y, Sysyn G, Ming W, Trager J, Starzl TE. Evolution of hepatitis B virus liver disease after hepatic replacement. Practical and theoretical considerations. *Am J Pathol* 1990; **137**: 667-676 [PMID: [2399936](#)]
- 7 **Davies SE**, Portmann BC, O'Grady JG, Aldis PM, Chaggar K, Alexander GJ, Williams R. Hepatic histological findings after transplantation for chronic hepatitis B virus infection, including a unique pattern of fibrosing cholestatic hepatitis. *Hepatology* 1991; **13**: 150-157 [PMID: [1988336](#)]
- 8 **Guo H**, Jiang D, Zhou T, Cuconati A, Block TM, Guo JT. Characterization of the intracellular deproteinized relaxed circular DNA of hepatitis B virus: an intermediate of covalently closed circular DNA formation. *J Virol* 2007; **81**: 12472-12484 [PMID: [17804499](#) DOI: [10.1128/JVI.01123-07](#)]
- 9 **Stevens CE**, Neurath RA, Beasley RP, Szmunness W. HBeAg and anti-HBe detection by radioimmunoassay: correlation with vertical transmission of hepatitis B virus in Taiwan. *J Med Virol* 1979; **3**: 237-241 [PMID: [479860](#) DOI: [10.1002/jmv.1890030310](#)]
- 10 **Spada E**, Mele A, Ciccozzi M, Tosti ME, Bianco E, Szklo A, Ragni P, Gallo G, Balocchi E, Sangalli M, Lopalco PL, Moiraghi A, Stroffolini T; SEIEVA collaborating group. Changing epidemiology of parenterally transmitted viral hepatitis: results from the hepatitis surveillance system in Italy. *Dig Liver Dis* 2001; **33**: 778-784 [PMID: [11838613](#) DOI: [10.1016/s1590-8658\(01\)80695-2](#)]
- 11 **Beasley RP**. Rocks along the road to the control of HBV and HCC. *Ann Epidemiol* 2009; **19**: 231-234 [PMID: [19344859](#) DOI: [10.1016/j.annepidem.2009.01.017](#)]
- 12 **Wu JF**, Chang MH. Natural history of chronic hepatitis B virus infection from infancy to adult life - the mechanism of inflammation triggering and long-term impacts. *J Biomed Sci* 2015; **22**: 92 [PMID: [26487087](#) DOI: [10.1186/s12929-015-0199-y](#)]
- 13 **Hsu YS**, Chien RN, Yeh CT, Sheen IS, Chiou HY, Chu CM, Liaw YF. Long-term outcome after spontaneous HBeAg seroconversion in patients with chronic hepatitis B. *Hepatology* 2002; **35**: 1522-1527 [PMID: [12029639](#) DOI: [10.1053/jhep.2002.33638](#)]
- 14 **Niederau C**, Heintges T, Lange S, Goldmann G, Niederau CM, Mohr L, Häussinger D. Long-term follow-up of HBeAg-positive patients treated with interferon alfa for chronic hepatitis B. *N Engl J Med* 1996; **334**: 1422-1427 [PMID: [8618580](#) DOI: [10.1056/NEJM199605303342202](#)]
- 15 **Liaw YF**. HBeAg seroconversion as an important end point in the treatment of chronic hepatitis B. *Hepatol Int* 2009; **3**: 425-433 [PMID: [19669245](#) DOI: [10.1007/s12072-009-9140-3](#)]
- 16 **Hui CK**, Leung N, Shek TW, Yao H, Lee WK, Lai JY, Lai ST, Wong WM, Lai LS, Poon RT, Lo CM, Fan ST, Lau GK; Hong Kong Liver Fibrosis Study Group. Sustained disease remission after spontaneous HBeAg seroconversion is associated with reduction in fibrosis progression in chronic hepatitis B Chinese patients. *Hepatology* 2007; **46**: 690-698 [PMID: [17680649](#) DOI: [10.1002/hep.21758](#)]
- 17 **Lin SM**, Yu ML, Lee CM, Chien RN, Sheen IS, Chu CM, Liaw YF. Interferon therapy in HBeAg positive chronic hepatitis reduces progression to cirrhosis and hepatocellular carcinoma. *J Hepatol* 2007; **46**: 45-52 [PMID: [17107734](#) DOI: [10.1016/j.jhep.2006.08.021](#)]
- 18 **Yang HI**, Lu SN, Liaw YF, You SL, Sun CA, Wang LY, Hsiao CK, Chen PJ, Chen DS, Chen CJ; Taiwan Community-Based Cancer Screening Project Group. Hepatitis B e antigen and the risk of hepatocellular carcinoma. *N Engl J Med* 2002; **347**: 168-174 [PMID: [12124405](#) DOI: [10.1056/NEJMoa013215](#)]
- 19 **Papathodoridis GV**, Hadziyannis SJ. Diagnosis and management of pre-core mutant chronic hepatitis B. *J Viral Hepat* 2001; **8**: 311-321 [PMID: [11555188](#) DOI: [10.1046/j.1365-2893.2001.00303.x](#)]
- 20 **Hoofnagle JH**. Reactivation of hepatitis B. *Hepatology* 2009; **49**: S156-S165 [PMID: [19399803](#) DOI: [10.1002/hep.22945](#)]
- 21 **Guo L**, Wang D, Ouyang X, Tang N, Chen X, Zhang Y, Zhu H, Li X. Recent Advances in HBV Reactivation Research. *Biomed Res Int* 2018; **2018**: 2931402 [PMID: [30687740](#) DOI: [10.1155/2018/2931402](#)]
- 22 **Lok AS**, Liang RH, Chiu EK, Wong KL, Chan TK, Todd D. Reactivation of hepatitis B virus replication in patients receiving cytotoxic therapy. Report of a prospective study. *Gastroenterology* 1991; **100**: 182-188 [PMID: [1983820](#) DOI: [10.1016/0016-5085\(91\)90599-g](#)]
- 23 **Esteve M**, Saro C, González-Huix F, Suarez F, Forné M, Viver JM. Chronic hepatitis B reactivation following infliximab therapy in Crohn's disease patients: need for primary prophylaxis. *Gut* 2004; **53**: 1363-1365 [PMID: [15306601](#) DOI: [10.1136/gut.2004.040675](#)]
- 24 **Moghhoofei M**, Mostafaei A, Ashraf-Ganjouei A, Kavosi H, Mahmoudi M. HBV reactivation in rheumatic diseases patients under therapy: A meta-analysis. *Microb Pathog* 2018; **114**: 436-443 [PMID: [29223453](#) DOI: [10.1016/j.micpath.2017.12.014](#)]
- 25 **Loomba R**, Liang TJ. Hepatitis B Reactivation Associated With Immune Suppressive and Biological Modifier Therapies: Current Concepts, Management Strategies, and Future Directions. *Gastroenterology* 2017; **152**: 1297-1309 [PMID: [28219691](#) DOI: [10.1053/j.gastro.2017.02.009](#)]
- 26 **Perrillo RP**, Gish R, Falck-Ytter YT. American Gastroenterological Association Institute technical review on prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology* 2015; **148**: 221-244.e3 [PMID: [25447852](#) DOI: [10.1053/j.gastro.2014.10.038](#)]
- 27 **Reddy KR**, Beavers KL, Hammond SP, Lim JK, Falck-Ytter YT; American Gastroenterological Association Institute. American Gastroenterological Association Institute guideline on the prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology* 2015; **148**: 215-9; quiz e16-7 [PMID: [25447850](#) DOI: [10.1053/j.gastro.2014.10.039](#)]
- 28 **Loomba R**, Rowley A, Wesley R, Liang TJ, Hoofnagle JH, Pucino F, Csako G. Systematic review: the effect of preventive lamivudine on hepatitis B reactivation during chemotherapy. *Ann Intern Med* 2008; **148**: 519-528 [PMID: [18378948](#) DOI: [10.7326/0003-4819-148-7-200804010-00008](#)]
- 29 **Pattullo V**. Hepatitis B reactivation in the setting of chemotherapy and immunosuppression - prevention is better than cure. *World J Hepatol* 2015; **7**: 954-967 [PMID: [25954478](#) DOI: [10.4254/wjh.v7.i7.954](#)]
- 30 **Lau GK**, Liang R, Chiu EK, Lee CK, Lam SK. Hepatic events after bone marrow transplantation in patients with hepatitis B infection: a case controlled study. *Bone Marrow Transplant* 1997; **19**: 795-799 [PMID: [9134171](#) DOI: [10.1038/sj.bmt.1700744](#)]
- 31 **Dhédin N**, Douvin C, Kuentz M, Saint Marc MF, Reman O, Rieux C, Bernaudin F, Norol F, Cordonnier C, Bobin D, Metreau JM, Vernant JP. Reverse seroconversion of hepatitis B after allogeneic bone marrow transplantation: a retrospective study of 37 patients with pretransplant anti-HBs and anti-HBe. *Transplantation* 1998; **66**: 616-619 [PMID: [9753342](#) DOI: [10.1097/00007890-199809150-00012](#)]
- 32 **Torisu M**, Yokoyama T, Amemiya H, Kohler PF, Schroter G, Martineau G, Penn I, Palmer W, Halgrimson CG, Putnam CW, Starzl TE. Immunosuppression, liver injury, and hepatitis in renal, hepatic, and cardiac homograft recipients: with particular reference to the Australia antigen. *Ann Surg* 1971; **174**: 620-639 [PMID: [4999690](#) DOI: [10.1097/0000658-197110000-00008](#)]

- 33 **Todo S**, Demetris AJ, Van Thiel D, Teperman L, Fung JJ, Starzl TE. Orthotopic liver transplantation for patients with hepatitis B virus-related liver disease. *Hepatology* 1991; **13**: 619-626 [PMID: [2010156](#)]
- 34 **Van Thiel DH**, Schade RR, Gavaler JS, Shaw BW, Iwatsuki S, Starzl TE. Medical aspects of liver transplantation. *Hepatology* 1984; **4**: 79S-83S [PMID: [6363264](#)]
- 35 **Samuel D**, Bismuth A, Mathieu D, Arulnaden JL, Reynes M, Benhamou JP, Brechot C, Bismuth H. Passive immunoprophylaxis after liver transplantation in HBsAg-positive patients. *Lancet* 1991; **337**: 813-815 [PMID: [1672913](#) DOI: [10.1016/0140-6736\(91\)92515-4](#)]
- 36 **Samuel D**, Muller R, Alexander G, Fassati L, Ducot B, Benhamou JP, Bismuth H. Liver transplantation in European patients with the hepatitis B surface antigen. *N Engl J Med* 1993; **329**: 1842-1847 [PMID: [8247035](#) DOI: [10.1056/NEJM199312163292503](#)]
- 37 **Chauhan R**, Lingala S, Gadiparthi C, Lahiri N, Mohanty SR, Wu J, Michalak TI, Satapathy SK. Reactivation of hepatitis B after liver transplantation: Current knowledge, molecular mechanisms and implications in management. *World J Hepatol* 2018; **10**: 352-370 [PMID: [29599899](#) DOI: [10.4254/wjh.v10.i3.352](#)]
- 38 **Benhamou Y**, Dohin E, Lunel-Fabiani F, Poynard T, Huraux JM, Katlama C, Opolon P, Gentilini M. Efficacy of lamivudine on replication of hepatitis B virus in HIV-infected patients. *Lancet* 1995; **345**: 396-397 [PMID: [7845151](#) DOI: [10.1016/s0140-6736\(95\)90388-7](#)]
- 39 **Dienstag JL**, Perrillo RP, Schiff ER, Bartholomew M, Vicary C, Rubin M. A preliminary trial of lamivudine for chronic hepatitis B infection. *N Engl J Med* 1995; **333**: 1657-1661 [PMID: [7477217](#) DOI: [10.1056/NEJM199512213332501](#)]
- 40 **Grellier L**, Mutimer D, Ahmed M, Brown D, Burroughs AK, Rolles K, McMaster P, Beranek P, Kennedy F, Kibbler H, McPhillips P, Elias E, Dusheiko G. Lamivudine prophylaxis against reinfection in liver transplantation for hepatitis B cirrhosis. *Lancet* 1996; **348**: 1212-1215 [PMID: [8898039](#) DOI: [10.1016/s0140-6736\(96\)04444-3](#)]
- 41 **Boucher CA**, Cammack N, Schipper P, Schuurman R, Rouse P, Wainberg MA, Cameron JM. High-level resistance to (-) enantiomeric 2'-deoxy-3'-thiacytidine in vitro is due to one amino acid substitution in the catalytic site of human immunodeficiency virus type 1 reverse transcriptase. *Antimicrob Agents Chemother* 1993; **37**: 2231-2234 [PMID: [7504909](#) DOI: [10.1128/aac.37.10.2231](#)]
- 42 **Ling R**, Mutimer D, Ahmed M, Boxall EH, Elias E, Dusheiko GM, Harrison TJ. Selection of mutations in the hepatitis B virus polymerase during therapy of transplant recipients with lamivudine. *Hepatology* 1996; **24**: 711-713 [PMID: [8781347](#) DOI: [10.1002/hep.510240339](#)]
- 43 **Bartholomew MM**, Jansen RW, Jeffers LJ, Reddy KR, Johnson LC, Bunzendahl H, Condreay LD, Tzakis AG, Schiff ER, Brown NA. Hepatitis-B-virus resistance to lamivudine given for recurrent infection after orthotopic liver transplantation. *Lancet* 1997; **349**: 20-22 [PMID: [8988118](#) DOI: [10.1016/S0140-6736\(96\)02266-0](#)]
- 44 **Honkoop P**, Niesters HG, de Man RA, Osterhaus AD, Schalm SW. Lamivudine resistance in immunocompetent chronic hepatitis B. Incidence and patterns. *J Hepatol* 1997; **26**: 1393-1395 [PMID: [9210628](#) DOI: [10.1016/s0168-8278\(97\)80476-x](#)]
- 45 **Mutimer D**, Pillay D, Shields P, Cane P, Ratcliffe D, Martin B, Buchan S, Boxall L, O'Donnell K, Shaw J, Hübscher S, Elias E. Outcome of lamivudine resistant hepatitis B virus infection in the liver transplant recipient. *Gut* 2000; **46**: 107-113 [PMID: [10601065](#) DOI: [10.1136/gut.46.1.107](#)]
- 46 **Perrillo R**, Rakela J, Dienstag J, Levy G, Martin P, Wright T, Caldwell S, Schiff E, Gish R, Villeneuve JP, Farr G, Anshuetz G, Crowther L, Brown N. Multicenter study of lamivudine therapy for hepatitis B after liver transplantation. Lamivudine Transplant Group. *Hepatology* 1999; **29**: 1581-1586 [PMID: [10216146](#) DOI: [10.1002/hep.510290507](#)]
- 47 **Markowitz JS**, Martin P, Conrad AJ, Markmann JF, Seu P, Yersiz H, Goss JA, Schmidt P, Pakrasi A, Artinian L, Murray NG, Imagawa DK, Holt C, Goldstein LI, Stribling R, Busuttill RW. Prophylaxis against hepatitis B recurrence following liver transplantation using combination lamivudine and hepatitis B immune globulin. *Hepatology* 1998; **28**: 585-589 [PMID: [9696028](#) DOI: [10.1002/hep.510280241](#)]
- 48 **Marzano A**, Salizzoni M, Debernardi-Venon W, Smedile A, Franchello A, Ciancio A, Gentilecore E, Piantino P, Barbui AM, David E, Negro F, Rizzetto M. Prevention of hepatitis B virus recurrence after liver transplantation in cirrhotic patients treated with lamivudine and passive immunoprophylaxis. *J Hepatol* 2001; **34**: 903-910 [PMID: [11451175](#) DOI: [10.1016/s0168-8278\(01\)00080-0](#)]
- 49 **Dumortier J**, Chevallier P, Scoazec JY, Berger F, Boillot O. Combined lamivudine and hepatitis B immunoglobulin for the prevention of hepatitis B recurrence after liver transplantation: long-term results. *Am J Transplant* 2003; **3**: 999-1002 [PMID: [12859536](#) DOI: [10.1034/j.1600-6143.2003.00191.x](#)]
- 50 **Zheng S**, Chen Y, Liang T, Lu A, Wang W, Shen Y, Zhang M. Prevention of hepatitis B recurrence after liver transplantation using lamivudine or lamivudine combined with hepatitis B immunoglobulin prophylaxis. *Liver Transpl* 2006; **12**: 253-258 [PMID: [16447195](#) DOI: [10.1002/lt.20701](#)]
- 51 **Sánchez-Fueyo A**, Rimola A, Grande L, Costa J, Mas A, Navasa M, Cirera I, Sánchez-Tapias JM, Rodés J. Hepatitis B immunoglobulin discontinuation followed by hepatitis B virus vaccination: A new strategy in the prophylaxis of hepatitis B virus recurrence after liver transplantation. *Hepatology* 2000; **31**: 496-501 [PMID: [10655276](#) DOI: [10.1002/hep.510310233](#)]
- 52 **Angelico M**, Di Paolo D, Trinito MO, Petrolati A, Araco A, Zazza S, Lionetti R, Casciani CU, Tisone G. Failure of a reinforced triple course of hepatitis B vaccination in patients transplanted for HBV-related cirrhosis. *Hepatology* 2002; **35**: 176-181 [PMID: [11786974](#) DOI: [10.1053/jhep.2002.30278](#)]
- 53 **Pruett T**. Vaccination for hepatitis B after transplantation: a realistic goal? *Hepatology* 2002; **35**: 235-237 [PMID: [11786982](#) DOI: [10.1053/jhep.2002.31024](#)]
- 54 **Lok AS**. Prevention of recurrent hepatitis B post-liver transplantation. *Liver Transpl* 2002; **8**: S67-S73 [PMID: [12362302](#) DOI: [10.1053/jlts.2002.35780](#)]
- 55 **Marcellin P**, Chang TT, Lim SG, Tong MJ, Sievert W, Shiffman ML, Jeffers L, Goodman Z, Wulfsohn MS, Xiong S, Fry J, Brosgart CL; Adefovir Dipivoxil 437 Study Group. Adefovir dipivoxil for the treatment of hepatitis B e antigen-positive chronic hepatitis B. *N Engl J Med* 2003; **348**: 808-816 [PMID: [12606735](#) DOI: [10.1056/NEJMoa020681](#)]
- 56 **Hadziyannis SJ**, Tassopoulos NC, Heathcote EJ, Chang TT, Kitis G, Rizzetto M, Marcellin P, Lim SG, Goodman Z, Wulfsohn MS, Xiong S, Fry J, Brosgart CL; Adefovir Dipivoxil 438 Study Group. Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N Engl J Med* 2003; **348**: 800-807 [PMID: [12606734](#) DOI: [10.1056/NEJMoa021812](#)]
- 57 **Mailliard ME**, Gollan JL. Suppressing hepatitis B without resistance--so far, so good. *N Engl J Med* 2003; **348**: 848-850 [PMID: [12606741](#) DOI: [10.1056/NEJMe020185](#)]
- 58 **Schiff ER**, Lai CL, Hadziyannis S, Neuhaus P, Terrault N, Colombo M, Tillmann HL, Samuel D, Zeuzem

- S, Lilly L, Rendina M, Villeneuve JP, Lama N, James C, Wulfsohn MS, Namini H, Westland C, Xiong S, Choy GS, Van Doren S, Fry J, Brosgart CL; Adefovir Dipivoxil Study 435 International Investigators Group. Adefovir dipivoxil therapy for lamivudine-resistant hepatitis B in pre- and post-liver transplantation patients. *Hepatology* 2003; **38**: 1419-1427 [PMID: [14647053](#) DOI: [10.1016/j.hep.2003.09.040](#)]
- 59 **Angus P**, Vaughan R, Xiong S, Yang H, Delaney W, Gibbs C, Brosgart C, Colledge D, Edwards R, Ayres A, Bartholomeusz A, Locarnini S. Resistance to adefovir dipivoxil therapy associated with the selection of a novel mutation in the HBV polymerase. *Gastroenterology* 2003; **125**: 292-297 [PMID: [12891527](#) DOI: [10.1016/s0016-5085\(03\)00939-9](#)]
- 60 **Villeneuve JP**, Durantel D, Durantel S, Westland C, Xiong S, Brosgart CL, Gibbs CS, Parvaz P, Werle B, Trépo C, Zoulim F. Selection of a hepatitis B virus strain resistant to adefovir in a liver transplantation patient. *J Hepatol* 2003; **39**: 1085-1089 [PMID: [14642631](#) DOI: [10.1016/j.jhep.2003.09.022](#)]
- 61 **Zoulim F**. Treatment of pre- and post-liver transplantation HBV infection: Should we aim at combination therapy? *Hepatology* 2003; **38**: 1353-1355 [PMID: [14647043](#) DOI: [10.1016/j.hep.2003.10.001](#)]
- 62 **Cholongitas E**, Goulis J, Akriviadis E, Papatheodoridis GV. Hepatitis B immunoglobulin and/or nucleos(t)ide analogues for prophylaxis against hepatitis b virus recurrence after liver transplantation: a systematic review. *Liver Transpl* 2011; **17**: 1176-1190 [PMID: [21656655](#) DOI: [10.1002/lt.22354](#)]
- 63 **European Association for the Study of the Liver**. EASL clinical practice guidelines: Management of chronic hepatitis B virus infection. *J Hepatol* 2012; **57**: 167-185 [PMID: [22436845](#) DOI: [10.1016/j.jhep.2012.02.010](#)]
- 64 **Chang TT**, Gish RG, de Man R, Gadano A, Sollano J, Chao YC, Lok AS, Han KH, Goodman Z, Zhu J, Cross A, DeHertogh D, Wilber R, Colonna R, Apelian D; BEHoLD A1463022 Study Group. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med* 2006; **354**: 1001-1010 [PMID: [16525137](#) DOI: [10.1056/NEJMoa051285](#)]
- 65 **Marcellin P**, Heathcote EJ, Buti M, Gane E, de Man RA, Krastev Z, Germanidis G, Lee SS, Flisiak R, Kaita K, Manns M, Kotzev I, Tchernev K, Buggisch P, Weilert F, Kurdas OO, Shiffman ML, Trinh H, Washington MK, Sorbel J, Anderson J, Snow-Lampart A, Mondou E, Quinn J, Rousseau F. Tenofovir disoproxil fumarate versus adefovir dipivoxil for chronic hepatitis B. *N Engl J Med* 2008; **359**: 2442-2455 [PMID: [19052126](#) DOI: [10.1056/NEJMoa0802878](#)]
- 66 **Chang TT**, Lai CL, Kew Yoon S, Lee SS, Coelho HS, Carrilho FJ, Poordad F, Halota W, Horsmans Y, Tsai N, Zhang H, Tenney DJ, Tamez R, Iloeje U. Entecavir treatment for up to 5 years in patients with hepatitis B e antigen-positive chronic hepatitis B. *Hepatology* 2010; **51**: 422-430 [PMID: [20049753](#) DOI: [10.1002/hep.23327](#)]
- 67 **Heathcote EJ**, Marcellin P, Buti M, Gane E, De Man RA, Krastev Z, Germanidis G, Lee SS, Flisiak R, Kaita K, Manns M, Kotzev I, Tchernev K, Buggisch P, Weilert F, Kurdas OO, Shiffman ML, Trinh H, Gurel S, Snow-Lampart A, Borroto-Esoda K, Mondou E, Anderson J, Sorbel J, Rousseau F. Three-year efficacy and safety of tenofovir disoproxil fumarate treatment for chronic hepatitis B. *Gastroenterology* 2011; **140**: 132-143 [PMID: [20955704](#) DOI: [10.1053/j.gastro.2010.10.011](#)]
- 68 **Locarnini S**. Primary resistance, multidrug resistance, and cross-resistance pathways in HBV as a consequence of treatment failure. *Hepatol Int* 2008; **2**: 147-151 [PMID: [19669299](#) DOI: [10.1007/s12072-008-9048-3](#)]
- 69 **Cholongitas E**, Papatheodoridis GV. High genetic barrier nucleos(t)ide analogue(s) for prophylaxis from hepatitis B virus recurrence after liver transplantation: a systematic review. *Am J Transplant* 2013; **13**: 353-362 [PMID: [23137006](#) DOI: [10.1111/j.1600-6143.2012.04315.x](#)]
- 70 **Laryea MA**, Watt KD. Immunoprophylaxis against and prevention of recurrent viral hepatitis after liver transplantation. *Liver Transpl* 2012; **18**: 514-523 [PMID: [22315212](#) DOI: [10.1002/lt.23408](#)]
- 71 **Di Paolo D**, Tisone G, Piccolo P, Lenci I, Zazza S, Angelico M. Low-dose hepatitis B immunoglobulin given "on demand" in combination with lamivudine: a highly cost-effective approach to prevent recurrent hepatitis B virus infection in the long-term follow-up after liver transplantation. *Transplantation* 2004; **77**: 1203-1208 [PMID: [15114086](#) DOI: [10.1097/01.tp.0000118904.63669.eb](#)]
- 72 **Gane EJ**, Angus PW, Strasser S, Crawford DH, Ring J, Jeffrey GP, McCaughan GW; Australasian Liver Transplant Study Group. Lamivudine plus low-dose hepatitis B immunoglobulin to prevent recurrent hepatitis B following liver transplantation. *Gastroenterology* 2007; **132**: 931-937 [PMID: [17383422](#) DOI: [10.1053/j.gastro.2007.01.005](#)]
- 73 **Fung J**, Cheung C, Chan SC, Yuen MF, Chok KS, Sharr W, Dai WC, Chan AC, Cheung TT, Tsang S, Lam B, Lai CL, Lo CM. Entecavir monotherapy is effective in suppressing hepatitis B virus after liver transplantation. *Gastroenterology* 2011; **141**: 1212-1219 [PMID: [21762659](#) DOI: [10.1053/j.gastro.2011.06.083](#)]
- 74 **Fung J**, Wong T, Chok K, Chan A, Cheung TT, Dai JW, Sin SL, Ma KW, Ng K, Ng KT, Seto WK, Lai CL, Yuen MF, Lo CM. Long-term outcomes of entecavir monotherapy for chronic hepatitis B after liver transplantation: Results up to 8 years. *Hepatology* 2017; **66**: 1036-1044 [PMID: [28370215](#) DOI: [10.1002/hep.29191](#)]
- 75 **Teperman LW**, Poordad F, Bzowej N, Martin P, Pungpapong S, Schiano T, Flaherty J, Dinh P, Rossi S, Subramanian GM, Spivey J. Randomized trial of emtricitabine/tenofovir disoproxil fumarate after hepatitis B immunoglobulin withdrawal after liver transplantation. *Liver Transpl* 2013; **19**: 594-601 [PMID: [23447407](#) DOI: [10.1002/lt.23628](#)]
- 76 **Fernández I**, Loinaz C, Hernández O, Abradelo M, Manrique A, Calvo J, Manzano M, García A, Cambra F, Castellano G, Jiménez C. Tenofovir/entecavir monotherapy after hepatitis B immunoglobulin withdrawal is safe and effective in the prevention of hepatitis B in liver transplant recipients. *Transpl Infect Dis* 2015; **17**: 695-701 [PMID: [26257166](#) DOI: [10.1111/tid.12434](#)]
- 77 **Manini MA**, Whitehouse G, Bruce M, Passerini M, Lim TY, Carey I, Considine A, Lampertico P, Suddle A, Heaton N, Heneghan M, Agarwal K. Entecavir or tenofovir monotherapy prevents HBV recurrence in liver transplant recipients: A 5-year follow-up study after hepatitis B immunoglobulin withdrawal. *Dig Liver Dis* 2018; **50**: 944-953 [PMID: [29735294](#) DOI: [10.1016/j.dld.2018.03.032](#)]
- 78 **Terrault NA**, Lok ASF, McMahon BJ, Chang KM, Hwang JP, Jonas MM, Brown RS, Bzowej NH, Wong JB. Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology* 2018; **67**: 1560-1599 [PMID: [29405329](#) DOI: [10.1002/hep.29800](#)]
- 79 **European Association for the Study of the Liver**. EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus infection. *J Hepatol* 2017; **67**: 370-398 [PMID: [28427875](#) DOI: [10.1016/j.jhep.2017.03.021](#)]
- 80 **Hussain M**, Soldevila-Pico C, Emre S, Luketic V, Lok AS; NIH HBV-OLT Study Group. Presence of

- intrahepatic (total and ccc) HBV DNA is not predictive of HBV recurrence after liver transplantation. *Liver Transpl* 2007; **13**: 1137-1144 [PMID: [17663390](#) DOI: [10.1002/lt.21179](#)]
- 81 **Lenci I**, Marcuccilli F, Tisone G, Di Paolo D, Tariciotti L, Ciotti M, Guenci T, Perno CF, Angelico M. Total and covalently closed circular DNA detection in liver tissue of long-term survivors transplanted for HBV-related cirrhosis. *Dig Liver Dis* 2010; **42**: 578-584 [PMID: [20097143](#) DOI: [10.1016/j.dld.2009.12.003](#)]
- 82 **Lenci I**, Tisone G, Di Paolo D, Marcuccilli F, Tariciotti L, Ciotti M, Svicher V, Perno CF, Angelico M. Safety of complete and sustained prophylaxis withdrawal in patients liver-transplanted for HBV-related cirrhosis at low risk of HBV recurrence. *J Hepatol* 2011; **55**: 587-593 [PMID: [21251938](#) DOI: [10.1016/j.jhep.2010.12.036](#)]
- 83 **Fox AN**, Terrault NA. Individualizing hepatitis B infection prophylaxis in liver transplant recipients. *J Hepatol* 2011; **55**: 507-509 [PMID: [21601598](#) DOI: [10.1016/j.jhep.2011.04.002](#)]
- 84 **Marzano A**, Gaia S, Ghisetti V, Carezzi S, Premoli A, Debernardi-Venon W, Alessandria C, Franchello A, Salizzoni M, Rizzetto M. Viral load at the time of liver transplantation and risk of hepatitis B virus recurrence. *Liver Transpl* 2005; **11**: 402-409 [PMID: [15776431](#) DOI: [10.1002/lt.20402](#)]
- 85 **Lenci I**, Baiocchi L, Tariciotti L, Di Paolo D, Milana M, Santopaolo F, Manzia TM, Toti L, Svicher V, Tisone G, Perno CF, Angelico M. Complete hepatitis B virus prophylaxis withdrawal in hepatitis B surface antigen-positive liver transplant recipients after longterm minimal immunosuppression. *Liver Transpl* 2016; **22**: 1205-1213 [PMID: [27272189](#) DOI: [10.1002/lt.24493](#)]
- 86 **Roche B**, Samuel D. Withdrawal of posttransplant hepatitis B virus prophylaxis: A blind test. *Liver Transpl* 2016; **22**: 1183-1185 [PMID: [27389861](#) DOI: [10.1002/lt.24511](#)]
- 87 **Sripongpun P**, Mannalithara A, Kwo PY, Kim WR. Potential Benefits of Switching Liver Transplant Recipients to Tenofovir Alafenamide Prophylaxis. *Clin Gastroenterol Hepatol* 2020; **18**: 747-749 [PMID: [31271737](#) DOI: [10.1016/j.cgh.2019.05.057](#)]
- 88 **Park ES**, Lee AR, Kim DH, Lee JH, Yoo JJ, Ahn SH, Sim H, Park S, Kang HS, Won J, Ha YN, Shin GC, Kwon SY, Park YK, Choi BS, Lee YB, Jeong N, An Y, Ju YS, Yu SJ, Chae HB, Yu KS, Kim YJ, Yoon JH, Zoulim F, Kim KH. Identification of a quadruple mutation that confers tenofovir resistance in chronic hepatitis B patients. *J Hepatol* 2019; **70**: 1093-1102 [PMID: [30794889](#) DOI: [10.1016/j.jhep.2019.02.006](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

