

# World Journal of *Stem Cells*

*World J Stem Cells* 2020 September 26; 12(9): 897-1049



## Contents

Monthly Volume 12 Number 9 September 26, 2020

### REVIEW

- 897** Dental stem cells: The role of biomaterials and scaffolds in developing novel therapeutic strategies  
*Granz CL, Gorji A*
- 922** Inflammatory niche: Mesenchymal stromal cell priming by soluble mediators  
*Jauković A, Kukulj T, Obradović H, Okić-Dorđević I, Mojsilović S, Bugarski D*
- 938** Role of the CXCR4-SDF1-HMGB1 pathway in the directional migration of cells and regeneration of affected organs  
*Haq N, Fareez IM, Fong LF, Mandal C, Abu Kasim NH, Kacharaju KR, Soesilawati P*
- 952** Mechanotransduction of stem cells for tendon repair  
*Wang HN, Huang YC, Ni GX*
- 966** Senescent mesenchymal stem/stromal cells and restoring their cellular functions  
*Meng QS, Liu J, Wei L, Fan HM, Zhou XH, Liang XT*
- 986** Mechanisms of action of neuropeptide Y on stem cells and its potential applications in orthopaedic disorders  
*Wu JQ, Jiang N, Yu B*

### MINIREVIEWS

- 1001** Stem cell treatments for oropharyngeal dysphagia: Rationale, benefits, and challenges  
*Tran EK, Juarez KO, Long JL*
- 1013** Perspectives on mesenchymal stem/progenitor cells and their derivatives as potential therapies for lung damage caused by COVID-19  
*Klimczak A*
- 1023** Mass acquisition of human periodontal ligament stem cells  
*Maeda H*

### ORIGINAL ARTICLE

#### Basic Study

- 1032** Human mesenchymal stem cells derived from umbilical cord and bone marrow exert immunomodulatory effects in different mechanisms  
*Song Y, Lim JY, Lim T, Im KI, Kim N, Nam YS, Jeon YW, Shin JC, Ko HS, Park IY, Cho SG*

**ABOUT COVER**

Editorial board member of *World Journal of Stem Cells*, Dr. Elisa Oltra is Distinguished Professor at the Catholic University of Valencia (UCV), Spain. Having received her Bachelor's and Master's degrees in Biochemistry at UCV in 1988 and her PhD degree in Cell and Molecular Biology at the University of Miami, FL, United States in 1999, she returned to UCV in 2009, where she became Director of the first official European Master's degree program in Biobanking. Her current research interests include the identification of molecular biomarkers of fibromyalgia and myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) and the study of immunomodulatory potential of mesenchymal stem cells. She has also served as Chair of the Biomarker subgroup ME/CFS Common Data Elements NINDS initiative at the National Institutes of Health (United States). (L-Editor: Filipodia)

**AIMS AND SCOPE**

The primary aim of *World Journal of Stem Cells* (*WJSC*, *World J Stem Cells*) is to provide scholars and readers from various fields of stem cells with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJSC* publishes articles reporting research results obtained in the field of stem cell biology and regenerative medicine, related to the wide range of stem cells including embryonic stem cells, germline stem cells, tissue-specific stem cells, adult stem cells, mesenchymal stromal cells, induced pluripotent stem cells, embryonal carcinoma stem cells, hemangioblasts, lymphoid progenitor cells, etc.

**INDEXING/ABSTRACTING**

The *WJSC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Biological Abstracts, BIOSIS Previews, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJSC* as 3.231; IF without journal self cites: 3.128; Ranking: 18 among 29 journals in cell and tissue engineering; Quartile category: Q3; Ranking: 113 among 195 journals in cell biology; and Quartile category: Q3.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Yan-Xia Xing; Production Department Director: Yun-Xiaojuan Wu; Editorial Office Director: Jin-Lai Wang.

**NAME OF JOURNAL**

*World Journal of Stem Cells*

**ISSN**

ISSN 1948-0210 (online)

**LAUNCH DATE**

December 31, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Carlo Ventura

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-0210/editorialboard.htm>

**PUBLICATION DATE**

September 26, 2020

**COPYRIGHT**

© 2020 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Role of the CXCR4-SDF1-HMGB1 pathway in the directional migration of cells and regeneration of affected organs

Nazmul Haque, Ismail M Fareez, Liew Fong Fong, Chanchal Mandal, Noor Hayaty Abu Kasim, Kranthi Raja Kacharaju, Pratiwi Soesilawati

**ORCID number:** Nazmul Haque 0000-0002-8191-8993; Ismail M Fareez 0000-0001-6993-5339; Liew Fong Fong 0000-0001-9165-2343; Chanchal Mandal 0000-0003-2865-2847; Noor Hayaty Abu Kasim 0000-0002-8889-842X; Kranthi Raja Kacharaju 0000-0002-2057-9473; Pratiwi Soesilawati 0000-0002-1141-5645.

**Author contributions:** Fareez IM and Fong LF contributed equally to this work; Haque N contributed to the conception and design of the study; Mandal C, Kacharaju KR and Soesilawati P drafted and wrote the article; Haque N, Fareez IM, Fong LF and Abu Kasim NH contributed to the writing of the manuscript, made critical revisions related to relevant intellectual content of the manuscript; Haque N and Abu Kasim NH approved the final version of the article.

**Supported by** MAHSA University Grant, No. RP158-05/19.

**Conflict-of-interest statement:** The authors declare no conflicts of interest.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative

**Nazmul Haque, Ismail M Fareez, Liew Fong Fong,** Department of Oral Biology and Biomedical Sciences, Faculty of Dentistry, MAHSA University, Selangor 42610, Malaysia

**Chanchal Mandal,** Biotechnology and Genetic Engineering Discipline, Life Science, Khulna University, Khulna 9208, Bangladesh

**Noor Hayaty Abu Kasim,** Faculty of Dentistry, University Kebangsaan Malaysia, Kuala Lumpur 50300, Malaysia

**Noor Hayaty Abu Kasim,** Faculty of Dental Medicine, Universitas Airlangga, Surabaya 411007, Indonesia

**Kranthi Raja Kacharaju,** Department of Conservative Dentistry, Faculty of Dentistry MAHSA University, Selangor 42610, Malaysia

**Pratiwi Soesilawati,** Department of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60115, Indonesia

**Corresponding author:** Nazmul Haque, PhD, Senior Lecturer, Department of Oral Biology and Biomedical Sciences, Faculty of Dentistry, MAHSA University, Jalan SP 2, Bandar Saujana Putra, Selangor 42610, Malaysia. [nazmul@mahsa.edu.my](mailto:nazmul@mahsa.edu.my); [tanna.bge@gmail.com](mailto:tanna.bge@gmail.com)

### Abstract

In recent years, several studies have reported positive outcomes of cell-based therapies despite insufficient engraftment of transplanted cells. These findings have created a huge interest in the regenerative potential of paracrine factors released from transplanted stem or progenitor cells. Interestingly, this notion has also led scientists to question the role of proteins in the secretome produced by cells, tissues or organisms under certain conditions or at a particular time of regenerative therapy. Further studies have revealed that the secretomes derived from different cell types contain paracrine factors that could help to prevent apoptosis and induce proliferation of cells residing within the tissues of affected organs. This could also facilitate the migration of immune, progenitor and stem cells within the body to the site of inflammation. Of these different paracrine factors present within the secretome, researchers have given proper consideration to stromal cell-derived factor-1 (SDF1) that plays a vital role in tissue-specific migration of the cells needed for regeneration. Recently researchers recognized that SDF1 could facilitate site-specific migration of cells by regulating SDF1-



Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Received:** March 3, 2020

**Peer-review started:** March 3, 2020

**First decision:** June 7, 2020

**Revised:** June 18, 2020

**Accepted:** July 18, 2020

**Article in press:** July 18, 2020

**Published online:** September 26, 2020

**P-Reviewer:** Huang YC, Politi L, Tang JM

**S-Editor:** Yan JP

**L-Editor:** Webster JR

**P-Editor:** Xing YX



CXCR4 and/or HMGB1-SDF1-CXCR4 pathways which is vital for tissue regeneration. Hence in this study, we have attempted to describe the role of different types of cells within the body in facilitating regeneration while emphasizing the HMGB1-SDF1-CXCR4 pathway that orchestrates the migration of cells to the site where regeneration is needed.

**Key Words:** C-X-C motif chemokine 12; Mesenchymal stem cells; Monocytes; Neutrophils; Peripheral blood mononuclear cells; Receptor for advanced glycation end products

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** In the last few decades, cell-based regenerative therapy has received considerable attention for the treatment of degenerative diseases or the regeneration of injured organs. However, poor cell retention is considered a major drawback associated with the short-term regenerative benefits. Furthermore, the short-term regenerative benefits are linked to paracrine factors secreted by the transplanted stem cells. To improve regenerative outcomes, researchers have identified the role of stromal cell-derived factor-1 (SDF1) as a key chemotactic factor that can facilitate site-specific migration and retention of transplanted cells, and stem or progenitor cells within the body by activating the SDF1-CXCR4 or HMGB1-SDF1-CXCR4 pathways.

**Citation:** Haque N, Fareez IM, Fong LF, Mandal C, Abu Kasim NH, Kacharaju KR, Soesilawati P. Role of the CXCR4-SDF1-HMGB1 pathway in the directional migration of cells and regeneration of affected organs. *World J Stem Cells* 2020; 12(9): 938-951

**URL:** <https://www.wjgnet.com/1948-0210/full/v12/i9/938.htm>

**DOI:** <https://dx.doi.org/10.4252/wjsc.v12.i9.938>

## INTRODUCTION

Over the past decades, non-communicable diseases, especially degenerative diseases are becoming more prevalent worldwide, which also contributes to major morbidity. During the past two decades, stem cell-based regenerative therapy has been considered hopeful in addressing the unmet needs of treating degenerative diseases<sup>[1]</sup>.

Among the different tools of regenerative medicine “stem cells” are considered the most promising due to their self-renewal capability and multi-differentiation potential. However, recent studies have shown that the positive outcomes of different types of stem cell-based regenerative therapies are not directly correlated to the engraftment of transplanted cells<sup>[2,3]</sup>. These findings have created a huge interest in the regenerative potential of paracrine factors and have led scientists to reveal the regenerative potential of proteins in the secretomes. Further studies have revealed the mitogenic, angiogenic, anti-apoptotic, anti-scarring and chemoattractant features of secretomes or cell culture supernatants that make them a potential tool for regenerative therapy<sup>[1,4]</sup>. Furthermore, the regenerative potential of the secretome from adult stem cells<sup>[5]</sup>, freshly isolated healthy peripheral blood mononuclear cells (PBMC)<sup>[6]</sup> and apoptosis-induced PBMC<sup>[7,8]</sup> has been acknowledged by several researchers. Secretomes from stem and progenitor cells have been found to be favorable for regenerating tissues or treating several disorders including neuronal disorders<sup>[9]</sup>, vascular diseases<sup>[10]</sup> and cutaneous wounds<sup>[11]</sup>. The growing evidence on the role of paracrine factors (cytokines, chemokines and growth factors) in the regeneration of affected organs has led to the introduction of cell culture supernatants or secretomes as a new therapeutic tool of regenerative medicine.

Regeneration is a complex process and several types of cells namely lymphocytes, monocytes, neutrophils, endothelial progenitor cells (EPCs), hematopoietic stem cells (HSCs), mesenchymal stem cells (MSCs), and tissue resident stem cells are involved in the process of regeneration. Recent studies reported that stromal cell-derived factor-1α (SDF1α) present in the secretome increases proliferation, viability, migration and homing of stem and progenitor cells; helps lymphoid tissue development and differentiation; and inhibits apoptosis of cells<sup>[4,12]</sup>. All these features are highly important for regeneration of damaged organs<sup>[13]</sup>. Furthermore, SDF1 and C-X-C-X-C

chemokine receptor type 4 (CXCR-4) play a vital role in the high mobility group box 1 (HMGB1)-mediated inflammatory cell recruitment to the site of damaged tissue which is also vital for tissue repair and regeneration<sup>[14]</sup>. Hence, in this review we have attempted to describe the role of different types of cells in regeneration while emphasizing activation of the HMGB1-SDF1-CXCR4 pathway which is considered a key pathway that regulates the directional migration of all cells and facilitates the process of regeneration.

## PBMC IN REGENERATION

PBMC are widely used in preclinical research and applications in vaccine trials, source of biomarkers in various infectious and chronic diseases, and are a useful tool in studying various aspects of pathology and biology *in vitro*<sup>[15]</sup>. In addition, PBMC are an easily accessible source of different types of adult stem and progenitor cells, such as HSCs, MSCs, osteoclast precursor cells, and EPCs<sup>[16]</sup>. Due to the content of different types of adult stem cells, in a favorable microenvironment the potential to differentiate into several tissue specific cells including mature blood cells, endothelial cells, hepatocytes, cardiomyocytes, smooth muscle cells, epithelial cells, neural cells, osteoblasts, osteoclasts, and myofibroblasts has been shown<sup>[16-18]</sup>. Furthermore, compared to bone marrow (BM) or other multipotent cells sources, the isolation of PBMC is less invasive. However, a series of standard procedures for PBMC collection, isolation, cryopreservation and preparation are crucial for their use in cell-based regenerative therapy<sup>[15]</sup>. PBMC contain terminally differentiated immune cells, namely monocytes and lymphocytes that also play a vital role in tissue remodeling and regeneration<sup>[19-21]</sup>.

### Monocytes

Monocytes that contribute approximately 4%-10% of leukocytes in our bloodstream are highly plastic in nature<sup>[22]</sup>. Monocytes and macrophages are the largest types of white blood cells and are involved in inflammation and elimination of harmful foreign substances<sup>[23,24]</sup>. As part of the innate immunity they are involved in tissue homeostasis and facilitate wound healing by removing apoptotic and necrotic cells<sup>[24]</sup>.

In regenerative tissues, macrophages are highly plastic and play a decisive role in tissue repair and regeneration<sup>[25]</sup>. In response to injury and subsequent healing, macrophages are capable of polarization towards a spectrum of phenotypes. Based on the environmental cues and molecular mediators, these cells will differentiate into either pro-inflammatory type I macrophage (M1) or anti-inflammatory type II macrophage (M2) phenotypes<sup>[25-27]</sup>.

Studies have reported that M1 macrophages infiltrate tissues at the earlier stages of acute injury to promote the clearance of necrotic cells or tissue debris. Moreover, following activation M1 macrophages secrete a wide range of pro-inflammatory cytokines such as interleukin (IL)-1, IL-6, IL-8, IL-12, IL-18, IL-23, tumor necrosis factor (TNF)- $\alpha$ , monocyte chemoattractant protein (MCP)-1 and macrophage inflammatory protein (MIP)-1<sup>[28]</sup>. Whereas, M2 macrophages appear within the injured tissue at later stages and release high amounts of anti-inflammatory paracrine factors such as IL-10 and transforming growth factor- $\beta$  (TGF- $\beta$ )<sup>[25,29]</sup>. They are also capable of secreting extracellular matrix (ECM) remodeling components such as fibronectin, osteopontin, fibrin cross-linker transglutaminase and promote tissue healing<sup>[30,31]</sup>. Although there are controversies regarding the sequential presence of the two different macrophages within tissues, this is because of the dynamic shift in macrophage polarization or the recruitment of new monocytes which do not invalidate the role of macrophages in tissue regeneration.

### Lymphocytes

Among the different types of lymphocytes, the regulatory T-cells (Treg) are involved in the repair and regeneration of affected tissues and organ systems. Following injury, Treg are recruited to the site to regulate inflammation and modulate the process of regeneration<sup>[32]</sup>. Following the initiation of inflammation, Treg inhibit recruitment of neutrophils by secreting IL-10 which in turn helps to minimize the secretion of inflammatory cytokines namely IL-1 $\beta$ , IL-6, interferon (IFN)- $\gamma$ , and TNF- $\alpha$ . Moreover, Treg induce apoptosis of neutrophils and clear debris by activation of M1 macrophages. In addition, they play a vital role in macrophage polarization towards the M2 phenotype by secreting anti-inflammatory cytokines such as IL-4, IL-10, and IL-13 which eventually support tissue repair and regeneration<sup>[32,33]</sup>. However, the

regenerative function of Treg follows a tissue specific manner. For instance, in skeletal muscle, Treg infiltrate the tissues in response to IL-33 following activation of the M1 population and removal of necrotic tissues by these cells. Following infiltration, Treg inhibit M1-mediated inflammation and shift the polarization of macrophages towards the M2 population<sup>[32]</sup>. Whereas, in heart tissues, recruitment of the higher number of Treg induces polarization of macrophages towards the M2 phenotype that help to inhibit inflammation, excessive matrix degradation, and adverse remodeling which eventually reduce ventricular ruptures and increases the rate of survival<sup>[34]</sup>.

## PERIPHERAL BLOOD POLYMORPHONUCLEAR CELLS IN REGENERATION

Neutrophils are the most abundantly found white blood cells in the human peripheral circulation and contribute approximately 50%-70% to all circulatory white blood cells. They are the first leukocyte population recruited to the site of injury and regulate the process of tissue regeneration positively or negatively<sup>[32]</sup>. However, the role of neutrophils in the process of regeneration is microenvironment-dependent and context-specific.

For example, in the case of skeletal muscle injury, neutrophils impair the restoration and function of muscles by releasing hypochlorous acid, nicotinamide adenine dinucleotide phosphate oxidase, and other cytokines<sup>[32,35,36]</sup>. Downregulation of lung regeneration following ischemia-reperfusion by neutrophils has also been reported<sup>[37]</sup>. It is noteworthy that neutrophils have also shown positive effects on the repair of lung epithelium and nerve cells<sup>[38,39]</sup>.

## MESENCHYMAL AND OTHER TISSUE-SPECIFIC STEM CELLS IN REGENERATION

Apart from HSCs, other adult stem cells or tissue-specific progenitor cells such as MSCs, EPCs, mammary stem cells, intestinal stem cells, and neural stem cells are found in adult tissues<sup>[40]</sup>. Tissue-specific stem cells maintain tissue homeostasis, while MSCs can differentiate into a variety of cell types. MSCs, in particular, have promising cell sources, as they can be harvested from various sources, such as BM, umbilical cord (UC), adipose tissue, and dental tissues<sup>[41-43]</sup>. Unlike embryonic stem cells (ES cells or ESCs), which are pluripotent, MSCs are multipotent cells which possess limited differentiation potential. Nevertheless, their potential to differentiate into osteoblasts and osteocytes is very well known. There is also accumulating evidence regarding their robust potential in tissue healing and regenerative medicine, in both preclinical and clinical studies<sup>[44-46]</sup>. According to a recent PubMed search conducted on November 2019, there were 110 MSC-based human clinical trials exploring the safety and efficacy of stem cells for tissue healing and the treatment of degenerative diseases. However, most of these trials were phase I and phase II, or a mixture of phase I/II studies. Whereas, phase III or phase II/III trials which investigate the long-term safety of MSC-based therapies prior to full establishment of MSCs in clinical practice are poorly documented. Thus, BM-derived MSCs (BMSCs) have been the most studied stem cells in cell therapy and tissue repair for the last 5 years, due to their multi-lineage differentiation potential<sup>[47]</sup>.

However, it is worthwhile noting that different MSC populations exhibit tissue-specific characteristics such as the expression of specific cell surface markers and transcription factors. In response to injury signals, these MSCs can potentially migrate from their niche to reach target tissues through vessel walls in the peripheral circulation<sup>[48]</sup>. Many studies have been conducted to investigate both the chemical and mechanical factors that influence the homing mechanism and engraftment of MSCs into local areas of damaged sites. The chemical factors that affect the trafficking process are the presence of a variety of chemokines, growth factors and cytokines, whereas the mechanical factors involved in the process include ECM stiffness, vascular cyclic stretching and hemodynamic forces or shear stress on the vessel walls<sup>[49]</sup>. These factors make up the vital characteristics of MSCs and result in their promising effect in tissue healing and differentiation.

The first characteristic of MSCs is their multi-lineage differentiation potential. MSCs are capable of differentiating into several mesoderm lineages, including adipogenic, osteogenic, chondrogenic and myogenic lineages, depending on the multitude of

stimuli and inhibitors present in the tissue microenvironment<sup>[50]</sup>. The microenvironment plays an important role in the activation or downregulation of transcription factors that regulate the expression of genes responsible for the induction and progression of tissue-specific differentiation<sup>[51]</sup>. MSCs can also generate neural cells in the ectodermal layer, and hepatic cells and pancreatic cells in the endodermal layer<sup>[52]</sup>. The study by Chen *et al*<sup>[53]</sup> was among the first to explore the ability of MSCs to differentiate into functional islet-like cells that might play an important role in the future treatment of diabetes. MSCs cultured in stiff scaffolds can easily differentiate into osteoblasts, and showed the potential for myogenic, adipogenic and neurogenic differentiation, respectively, but with a decrease in elasticity. Recently, Jung *et al*<sup>[54]</sup> demonstrated that ECM proteins in 3D composites were able to trigger differentiation of BMSCs into mesodermal lineages with enhanced adipogenic differentiation and IL-6 expression compared to that in 2D ECM proteins.

Secondly, MSCs are capable of dynamic interactions with their microenvironment and secrete a wide variety of paracrine factors that are required for tissue recovery or wound healing. Several studies refuted the hypothesis that direct trans-differentiation or cell fusion of MSCs was the principal mechanism underlying their therapeutic action in tissue regeneration<sup>[55]</sup>. Indeed, MSCs transplantation regulated released factors in experimental models of tissue injury, which was largely associated with suppression of immune and inflammatory reactions, inhibition of apoptosis, and enhancement of cell proliferation and angiogenesis, thereby promoting regeneration of the tissue<sup>[56]</sup>. Apart of MSCs-mediated secretion of these paracrine and autocrine factors, extracellular vesicles such as exosomes and microvesicles may also regulate these functional roles<sup>[57,58]</sup>. The MSC-mediated factors released at high levels include the following: (1) Growth factors and their receptors [*i.e.*, granulocyte-macrophage colony-stimulating factor (GM-CSF), basic fibroblast growth factor (bFGF), vascular endothelial growth factor (VEGF), insulin-like growth factor-binding proteins (IGFBP3, IGFBP4, IGFBP7) and bone morphogenetic protein 2 (BMP-2)]; (2) Extracellular matrix remodelers/mediators [*i.e.*, periostin, fibronectin, collagen, TIMP metalloproteinase inhibitor 2 (TIMP-2), metalloproteinase inhibitors, and decorin]; and (3) Immune system signaling regulatory proteins (*i.e.*, TGF- $\beta$ , MCP-1, IL-6, and IL-8)<sup>[49,59]</sup>. Various studies have demonstrated that the released pro-inflammatory cytokines up-regulate the efficacy of MSC-mediated immunomodulation and functional improvement in microvascular injury<sup>[60]</sup>, inflammatory liver disease<sup>[61]</sup>, osteoarthritis<sup>[62]</sup>, spinal cord injury<sup>[63]</sup>, brain cancer<sup>[64]</sup>, ischemic limb regeneration<sup>[65]</sup>, and asthmatic<sup>[66]</sup> models. Taken together, it is well accepted that the combination of MSCs with these trophic factors can modulate their behavior during inflammation and tissue injury. Research should now focus on the strategies to manipulate and modulate the secretion of these molecules in the infused or implanted MSCs microenvironment to enhance their functional role<sup>[1]</sup>.

Finally, MSCs exhibit immunomodulatory properties<sup>[67-69]</sup>. The immunomodulatory properties of MSCs proved effective in treating various immune disorders in both *in vivo* and human studies. MSCs modulate the functions of almost all cells of both the innate and adaptive immune systems and induce an anti-inflammatory phenotype<sup>[59]</sup>. MSCs interact with a variety of immune cells and have the capacity to inhibit the excessive response of B cells, T cells, macrophages, dendritic cells, and natural killer cells<sup>[68]</sup>. Nevertheless, the underlying molecular and cellular mechanisms behind MSC-mediated immunomodulation have not been fully elucidated. MSCs have been shown to modulate the immune response by secreting soluble factors [*e.g.*, IL-6, M-CSF, IL-10, TGF- $\beta$ , HGF, and prostaglandin E2 (PGE2)] in the presence of adhesion molecules [*i.e.*, vascular cell adhesion molecule (VCAM)-1, intercellular adhesion molecule (ICAM)-1, and lymphocyte function-associated antigen (LFA)-3]<sup>[70-72]</sup>. Through a synergy of cell contact-dependent mechanisms and these soluble factors, MSCs are able to initiate the T-cell interactions that play a prominent role in their immunomodulatory potential<sup>[71,73]</sup>. Furthermore, anti-inflammatory monocytes/macrophages and Tregs are also important in MSC-mediated immunosuppression<sup>[69,74]</sup>. Studies also linked the low immunogenic properties of MSCs to the lower level of expression of major histocompatibility complex (MHC) class I antigens, and lack of MHC class II and co-stimulatory molecules such as CD80, CD86, and CD40<sup>[75,76]</sup>. Although the immunomodulatory effect of MSCs is hypothesized to be *via* MSC-secreted cytokines in many studies, most studies documented that MSCs act differently depending on the local microenvironment and the presence of inflammatory cytokines during the pre-treatment of MSCs. An understanding of the immune suppressive role of MSCs would enhance prospective clinical applications of these cells.

Thus, the fate of MSCs is vastly influenced by their environment which includes mechanical or physical stimulation, growth factors, cell density, and cell-cell



attachment or interactions. However, this multipotency of MSCs could also be due to another reason which has been widely discussed. In fact, a debate is currently ongoing regarding the 'stem cell' status of MSCs<sup>[77]</sup>. It is postulated that MSCs are purely specific adult stem cells, which contradicts findings that MSCs are a diverse mixture of many specific lineage progenitor cells. However, these shortcomings provide a good reason for the continuous research on MSCs in stem-cell based therapy.

## CELL MIGRATION IS ESSENTIAL FOR TISSUE REGENERATION

Progenitors and MSCs migrate and initiate the homing mechanism in response to inflammatory signaling molecules and corresponding receptors around the injured tissue. MSCs are therapeutically capable of reaching and homing to sites of inflammation by various routes such as intravenous (IV), intra-arterial (IA), intraparenchymal, intracoronary (IC) local administration and into the subarachnoid and epidural spaces<sup>[48]</sup>. From the systemic circulation, MSCs migrate specifically to damaged tissue sites and exert their functional effects locally under a variety of pathologic conditions. Luger *et al.*<sup>[78]</sup> demonstrated that intravenously administered fluorescent and radiolabeled MSCs homed to regions of myocardial injury to suppress the progressive deterioration in left ventricular function and adverse remodeling in mice, and it is thought to be a feasible and effective therapeutic strategy for the treatment of patients with large infarcts and ischemic cardiomyopathy. MSCs homing involves various chemokines and their receptors (*i.e.*, SDF1, CCL5, CXCR4, CXCR5, CXCR6, CCR2, CCR3, and CCR4), matrix metalloproteinases (MMPs) [MMP-2 and membrane type 1 MMP (MT1-MMP)], receptor tyrosine kinase dependent growth factors [*e.g.*, hepatocyte growth factor-Mesenchymal Epithelial Transition Factor (c-Met) proto-oncogene/receptor tyrosine kinase (HGF/c-Met) axes, platelet-derived growth factor (PDGF) and insulin-like growth factor 1 (IGF-1)] and some other adhesion molecules (*i.e.*, integrin  $\beta$ 1, integrin  $\alpha$ 4, and VCAM)<sup>[79-82]</sup>. These homing signals are released by injured cells and/or respondent immune cells. Besides these homing signals, other molecules are implicated in different steps of the homing process such as PGE2 and hematopoietic cell E-/L-selectin ligand (HCELL) that are functionally involved in cell migration to the injured tissue<sup>[83]</sup>. These factors could be a feasible strategy to facilitate therapeutic delivery of MSCs to targeted injured tissue.

Of the different chemokines and chemokine-mediated pathways, the SDF1-CXCR4 and HMGB1-SDF1-CXCR4 axis have received considerable attention due to their potential in-site specific directional migration of stem and progenitor cells. The role of HMGB1-SDF1-CXCR4 in regeneration of injured tissues or organs is discussed further below.

## HMGB1-SDF1-CXCR4 AXIS IN FACILITATING TISSUE-SPECIFIC MIGRATION

### ***HMGB1 in orchestrating the process of migration and regeneration***

HMGB1 protein is a highly conserved non-histone nuclear protein that binds to DNA and regulates the expression of genes and the chromosomal architecture<sup>[84]</sup>. Extracellular HMGB1 is actively secreted from activated or stressed immune cells, while passively secreted from necrotic tissues<sup>[85,86]</sup>. Following secretion into the extracellular space, HMGB1 exerts chemotactic activity or acts as a damage-associated molecular pattern molecule<sup>[87]</sup>. Indeed, the overall signaling mechanism by HMGB1 interacting with target cells needs to be elucidated for future therapeutic intervention<sup>[88]</sup>.

Wound healing is a complex process that involves the ECM, cytokines, growth factors and several types of cells. The steps involved in the process of wound healing include hemostasis, inflammation, cell migration and proliferation, wound contraction, and remodeling<sup>[89,90]</sup>. During the inflammatory phase, vasodilation followed by early vasoconstriction which is mediated by histamine, leukotrienes, and prostaglandins, increases capillary permeability and cell migration into the wound site<sup>[91]</sup>. Neutrophils are the first among the infiltrated cells to the site of injury followed by monocytes and lymphocytes. Initiation of leukocyte migration is mediated by several autocrine and paracrine factors. In addition, proteases are involved in the elimination of denatured ECM components. Following infiltration into the site of injury, monocytes transform into macrophages and clear debris from the area, release

cytokines and growth factors, such as FGF, TGF- $\beta$ , PDGF, and EGF that help to initiate the formation of granulation tissue<sup>[92]</sup>. HMGB1 also acts as an important chemotactic factor that regulates the directional migration of monocytes and neutrophils<sup>[93]</sup>. Following injury or inflammation, HMGB1 is released into the extracellular space and triggers the secretion of TNF, IL-1 $\alpha$ , IL-6, and IL-8 from monocytes, macrophages and neutrophils<sup>[94-96]</sup>.

Upon interaction of HMGB1 with the advanced glycation products (RAGE), toll-like receptor (TLR) 2, TLR4, and TLR9, activate pro-inflammatory responses thereby facilitating cell migration and the release of pro-inflammatory cytokines (Figure 1)<sup>[97-99]</sup>. In 2018, Xue *et al*<sup>[100]</sup> showed that the HMGB1/RAGE axis mediated migration of neural stem cells (NSCs) by the formation of filopodia which was further linked to the activation of RAGE/Rac and CDC42 or the RAGE/MAPK signaling cascade.

### **SDF1 in upregulation of the HMGB1-SDF1-CXCR4 axis**

SDF1 $\alpha$ , known as C-X-C motif chemokine 12 (CXCL12), is a chemotactic factor encoded by the CXCL12 gene on chromosome 10<sup>[101]</sup>. Studies have reported the therapeutic potential of the SDF1-CXCR4 axis in tissue regeneration. SDF1 is capable of activation, mobilization, homing and retention of HSCs, MSCs and several progenitor cells<sup>[80,102-104]</sup>. SDF1 is able to bind to CXCR4 and CXCR7. However, the SDF1-CXCR4 axis induces the homing process by regulating the cellular secretion and cell adhesion molecules, while SDF1-CXCR7 is involved in angiogenesis and tumor development<sup>[105]</sup>.


Principally, the binding of chemokine SDF1 to the chemokine receptor CXCR4 plays an important role in homeostatic regulation of leukocyte trafficking, hematopoiesis, organogenesis, cell differentiation and tissue regeneration in response to other molecules that are involved in triggering inflammation<sup>[14,106]</sup>. The mechanism of MSCs mobilization mediated by HMGB1 is analogous to the recruitment of inflammatory cells to injured tissues for leukocyte trafficking and homing (Figure 1). As mentioned above, HMGB1 acts as a damage-associated molecular pattern which is released either from necrotic cells or by secretion from activated immune cells, hepatocytes, enterocytes, and possibly several other types of cells under distress<sup>[107]</sup>. Stress conditions that promote HMGB1 secretion include hypoxia<sup>[108]</sup>, lethal irradiation<sup>[109]</sup>, treatment with specific antitumor drugs<sup>[110]</sup> or through regulation of autophagy<sup>[111]</sup>. HMGB1-induced cell migration requires both I $\kappa$ B kinase (IKK)- $\beta$  and IKK $\alpha$ -dependent nuclear factor- $\kappa$ B (NF- $\kappa$ B) activation. IKK $\beta$ -mediated activation of NF- $\kappa$ B maintains expression of RAGE, while continuous production of SDF1 is ensured by IKK $\alpha$ -dependent NF- $\kappa$ B activation<sup>[112,113]</sup>. Moreover, HMGB1 induces both physical and functional interactions between molecules that prevent the degradation of SDF1<sup>[114]</sup>.

In 2012, Kew *et al*<sup>[115]</sup> proposed that the SDF1-CXCR4 axis works as a co-receptor signal for RAGE receptor-dependent HMGB1 migration responses. Furthermore, SDF1 binding to CXCR4 can also induce CXCR4-TCR heterodimerization, which in turn can enhance gene transcription, cytokine production, increased calcium ion concentrations, and could facilitate cell migration. However, it is possible that the SDF1-CXCR4 axis might have other indirect effects on the regulation of cell migration such as enhancing HMGB1 binding to RAGE, which require further investigation.

### **TLR and RAGE dependent or independent activation of the HMGB1-SDF1-CXCR4 axis**

There is ample evidence of the capability of stem cells to regulate numerous growth factors, cytokines and chemokines. Chemokines, specifically, regulate cell locomotion and integrin function by binding to seven transmembrane domain receptors coupled to G-protein-coupled receptors (GPCR)s, which are heterotrimeric GTP-binding proteins, that are differentially expressed in various cell types<sup>[48,107]</sup>. In addition, there is always a need to assess the consequences of the combined activity of these chemokines and other inflammatory molecules to control appropriate tissue distribution of distinct leukocyte subsets under normal and pathological conditions. One of the interesting insights in stem cell research is the effect of such paracrine factors in the HMGB1-SDF1-CXCR4 signaling pathway during tissue regeneration.

Extracellular HMGB1 can interact with different molecules to dictate their biologic effects. The role of HMGB1 as a chemokine or cytokine is determined by its oxidative state (Figure 1). The role of extracellular HMGB1 to promote cell migration was first reported in smooth muscle cells in 2010<sup>[116]</sup>. Similar involvement was also reported in different cell types in the same year by Rauvala and Rouhiainen<sup>[117]</sup>. Studies showed that HMGB1-induced cell migration requires the formation of a heterocomplex with



**Figure 1 Schematic diagram of the dynamic HMGB1–CXCL12–SDF1 axis for accelerated tissue regeneration.** Disulfide (oxidized) HMGB1 usually binds to RAGE and TLR and regulates the expression of genes with pro- or anti-inflammatory properties and partial chemotactic properties. Whereas, fully reduced HMGB1 released from necrotic or stressed cells forms a heterocomplex with SDF1 secreted from activated immune cells or from cells within the injured tissues. Later, this heterocomplex binds to the CXCR4 receptors on the cells and facilitates site-specific migration<sup>[123-125]</sup>. MSC: Mesenchymal stem cells.

SDF1 and further binding with CXCR4, and not with RAGE, TLR2, or TLR4. Furthermore, it was also reported that HMGB1 does not affect migration by other chemokines such as CXCL8, CCL2, CCL7, CCL19, and CCL21<sup>[14]</sup>.

The bonding chemistry between SDF1 and HMGB1 was analyzed by NMR chemical shift mapping and revealed that most of the amino acids present in SDF1 have the ability to bind HMGB1 or its individual HMG boxes<sup>[14]</sup>. Each HMGB1 molecule has two HMG boxes and thereby can attach two SDF1 molecules at a time. Interestingly, the first few N-terminal residues of the SDF1 molecule do not attach to the HMGB1 molecule and remain free. These free residues can access deep inside the CXCR4 transmembrane domain to initiate signaling cascades<sup>[118]</sup>. As the HMGB1-SDF1 heterocomplex can present two SDF1 ligand molecules to dimers of the CXCR4 receptor, this heterocomplex would be more efficient than SDF1 alone in inducing cellular migration<sup>[119]</sup>. Alternatively, the HMGB1-SDF1 heterocomplex may help to unlock the CXCR4 binding site to promote SDF1 binding, or help lock in SDF1 into the CXCR4 transmembrane domain by providing direct HMGB1-CXCR4 contacts.

HMGB1 induces changes in SDF1 residues that are responsible for the activation of CXCR4, the SDF1 receptor. An analysis using fluorescence resonance energy transfer (FRET) demonstrated that there are different conformational rearrangements of CXCR4 homodimers triggered by SDF1 alone or in complex with HMGB1<sup>[14]</sup>. It has also been hypothesized that the formation of a heterocomplex between HMGB1 and SDF1 acts through CXCR4 which promotes the recruitment of monocytes to the injury site<sup>[14]</sup>. The interaction of locally produced SDF1 and its receptor CXCR4 expressed on the surface of MSCs plays an important role in the homing of transplanted cells. The binding of SDF1 to both CXCR4 and CXCR7 is also responsible for the production of paracrine mediators, including VEGF, IGF-1,  $\beta$ -FGF and HGF that exert mitogenic, pro-angiogenic, anti-apoptotic, and anti-inflammatory effects<sup>[120]</sup>. Hypoxia has been shown to enhance the expression of both SDF1 receptors, CXCR4 and CXCR7, in MSCs. Liu *et al.*<sup>[121]</sup> demonstrated that SDF1 $\alpha$  is upregulated in ischemic kidneys during reduced oxygen tension. Hypoxia induces expression of CXCR4 and CXCR7 while promoting the role of both SDF1 receptors for enhanced migration, adhesion and survival of hypoxia preconditioned (HP)-MSCs and thus improves homing of systemically delivered MSCs to the ischemic kidney. In addition, in normal culture-expanded MSCs, CXCR4 expression will alleviate progressively and thus could affect its ability to migrate toward the SDF1 gradient in the ischemic tissue.

The intracellular signaling cascades have not yet been clearly demonstrated. In the case of SDF1-CXCR4 bonding, activation and coordination of focal adhesion kinase (FAK) and phosphoinositide 3-kinase (PI3K) were reported in the migration of human dental pulp stem cells<sup>[122]</sup>. Subsequently, increased  $\beta$ -catenin expression by phosphorylation of protein kinase B (Akt) at ser473 that inhibits the activation of glycogen synthase kinase 3 beta (GSK3 $\beta$ ) was also reported. All these results indicate that the SDF1-CXCR4 axis activates the FAK/PI3K/Akt and GSK3 $\beta$ / $\beta$ -catenin pathways that could facilitate the migration of human dental pulp stem cells. Whereas in the HMGB1-SDF1-CXCR4 axis, elevated extracellular signal-regulated kinase (ERK) phosphorylation and Ca<sup>2+</sup> release from stores were reported<sup>[14]</sup>. Elevated ERK phosphorylation was observed in the presence of the SDF1-HMGB1 heterocomplex but not observed in the presence of SDF1 and HMGB1 alone. In the presence of HMGB1 a suboptimal SDF1 concentration was reported with a rapid increase in intracellular Ca<sup>2+</sup>.

## CONCLUSION

Until now, the migration and retention of transplanted cells have been considered a major drawback of cell-based regenerative therapy. SDF1 and its receptor CXCR4 play an important role in maintaining homeostasis by facilitating the homing of progenitor or other adult multipotent stem cells in the BM and regulating their mobilization into peripheral tissues during injury or stress. Studies have shown the potential of the SDF1-CXCR4 axis and/or HMGB1-SDF1-CXCR4 signaling pathways in regulating the process of directional migration followed by retention which are vital for the regeneration of injured tissues or organs. In addition, these pathways could play a major role in regulating the inflammatory conditions at the site of injury. Further studies concentrating on these pathways could make cell-based regenerative therapy more efficient and fruitful.

## REFERENCES

- 1 **Haque N**, Abdullah BJJ, BJJ, Kasim NHA. Secretome: Pharmaceuticals for Cell-Free Regenerative Therapy. In: Pham PV. Stem Cell Drugs - A New Generation of Biopharmaceuticals. Cham: Springer International Publishing; 2018: 17-35 [DOI: [10.1007/978-3-319-99328-7\\_2](https://doi.org/10.1007/978-3-319-99328-7_2)]
- 2 **Crivelli B**, Chlapanidas T, Perteghella S, Lucarelli E, Pascucci L, Brini AT, Ferrero I, Marazzi M, Pessina A, Torre ML; Italian Mesenchymal Stem Cell Group (GISM). Mesenchymal stem/stromal cell extracellular vesicles: From active principle to next generation drug delivery system. *J Control Release* 2017; **262**: 104-117 [PMID: [28736264](https://pubmed.ncbi.nlm.nih.gov/28736264/) DOI: [10.1016/j.jconrel.2017.07.023](https://doi.org/10.1016/j.jconrel.2017.07.023)]
- 3 **Boido M**, Ghibaudi M, Gentile P, Favaro E, Fusaro R, Tonda-Turo C. Chitosan-based hydrogel to support the paracrine activity of mesenchymal stem cells in spinal cord injury treatment. *Sci Rep* 2019; **9**: 6402 [PMID: [31024032](https://pubmed.ncbi.nlm.nih.gov/31024032/) DOI: [10.1038/s41598-019-42848-w](https://doi.org/10.1038/s41598-019-42848-w)]
- 4 **Haque N**, Kasim NHA, Kassim NLA, Rahman MT. Autologous serum supplement favours in vitro regenerative paracrine factors synthesis. *Cell Prolif* 2017; **50**: e12354 [PMID: [28682474](https://pubmed.ncbi.nlm.nih.gov/28682474/) DOI: [10.1111/cpr.12354](https://doi.org/10.1111/cpr.12354)]
- 5 **Madrigal M**, Rao KS, Riordan NH. A review of therapeutic effects of mesenchymal stem cell secretions and induction of secretory modification by different culture methods. *J Transl Med* 2014; **12**: 260 [PMID: [25304688](https://pubmed.ncbi.nlm.nih.gov/25304688/) DOI: [10.1186/s12967-014-0260-8](https://doi.org/10.1186/s12967-014-0260-8)]
- 6 **Hoetzenecker K**, Zimmermann M, Hoetzenecker W, Schweiger T, Kollmann D, Mildner M, Hegedus B, Mitterbauer A, Hacker S, Birner P, Gabriel C, Gyöngyösi M, Blyszczuk P, Eriksson U, Ankersmit HJ. Mononuclear cell secretome protects from experimental autoimmune myocarditis. *Eur Heart J* 2015; **36**: 676-685 [PMID: [23321350](https://pubmed.ncbi.nlm.nih.gov/23321350/) DOI: [10.1093/eurheartj/ehs459](https://doi.org/10.1093/eurheartj/ehs459)]
- 7 **Hoetzenecker K**, Assinger A, Lichtenauer M, Mildner M, Schweiger T, Starlinger P, Jakab A, Berényi E, Pavo N, Zimmermann M, Gabriel C, Plass C, Gyöngyösi M, Volf I, Ankersmit HJ. Secretome of apoptotic peripheral blood cells (APOSEC) attenuates microvascular obstruction in a porcine closed chest reperfused acute myocardial infarction model: role of platelet aggregation and vasodilation. *Basic Res Cardiol* 2012; **107**: 292 [PMID: [22899170](https://pubmed.ncbi.nlm.nih.gov/22899170/) DOI: [10.1007/s00395-012-0292-2](https://doi.org/10.1007/s00395-012-0292-2)]
- 8 **Altmann P**, Mildner M, Haider T, Traxler D, Beer L, Ristl R, Golabi B, Gabriel C, Leutmezer F, Ankersmit HJ. Secretomes of apoptotic mononuclear cells ameliorate neurological damage in rats with focal ischemia. *F1000Res* 2014; **3**: 131 [PMID: [25383184](https://pubmed.ncbi.nlm.nih.gov/25383184/) DOI: [10.12688/f1000research.4219.2](https://doi.org/10.12688/f1000research.4219.2)]
- 9 **Pires AO**, Neves-Carvalho A, Sousa N, Salgado AJ. The Secretome of Bone Marrow and Wharton Jelly Derived Mesenchymal Stem Cells Induces Differentiation and Neurite Outgrowth in SH-SY5Y Cells. *Stem Cells Int* 2014; **2014**: 438352 [PMID: [25132857](https://pubmed.ncbi.nlm.nih.gov/25132857/) DOI: [10.1155/2014/438352](https://doi.org/10.1155/2014/438352)]
- 10 **Dao M**, Tate CC, McGrogan M, Case CC. Comparing the angiogenic potency of naïve marrow stromal cells and Notch-transfected marrow stromal cells. *J Transl Med* 2013; **11**: 81 [PMID: [23531336](https://pubmed.ncbi.nlm.nih.gov/23531336/) DOI: [10.1186/1479-5876-11-81](https://doi.org/10.1186/1479-5876-11-81)]
- 11 **Mildner M**, Hacker S, Haider T, Gschwandtner M, Werba G, Barresi C, Zimmermann M, Golabi B, Tschachler E, Ankersmit HJ. Secretome of peripheral blood mononuclear cells enhances wound healing.



- PLoS One* 2013; **8**: e60103 [PMID: 23533667 DOI: 10.1371/journal.pone.0060103]
- 12 **Zhang M**, Mal N, Kiedrowski M, Chacko M, Askari AT, Popovic ZB, Koc ON, Penn MS. SDF-1 expression by mesenchymal stem cells results in trophic support of cardiac myocytes after myocardial infarction. *FASEB J* 2007; **21**: 3197-3207 [PMID: 17496162 DOI: 10.1096/fj.06-6558com]
  - 13 **Abduelmula A**, Huang R, Pu Q, Tamamura H, Morosan-Puopolo G, Brand-Saberi B. SDF-1 controls the muscle and blood vessel formation of the somite. *Int J Dev Biol* 2016; **60**: 29-38 [PMID: 26934287 DOI: 10.1387/ijdb.150132rh]
  - 14 **Schiraldi M**, Raucci A, Muñoz LM, Livoti E, Celona B, Venereau E, Apuzzo T, De Marchis F, Pedotti M, Bachi A, Thelen M, Varani L, Mellado M, Proudfoot A, Bianchi ME, Uguccioni M. HMGB1 promotes recruitment of inflammatory cells to damaged tissues by forming a complex with CXCL12 and signaling via CXCR4. *J Exp Med* 2012; **209**: 551-563 [PMID: 22370717 DOI: 10.1084/jem.20111739]
  - 15 **Betsou F**, Gaignaux A, Ammerlaan W, Norris PJ, Stone M. Biospecimen Science of Blood for Peripheral Blood Mononuclear Cell (PBMC) Functional Applications. *Curr Pathobiol Rep* 2019; **7**: 17-27 [DOI: 10.1007/s40139-019-00192-8]
  - 16 **Zhang M**, Huang B. The multi-differentiation potential of peripheral blood mononuclear cells. *Stem Cell Res Ther* 2012; **3**: 48 [PMID: 23200034 DOI: 10.1186/scrt139]
  - 17 **Hopper N**, Wardale J, Brooks R, Power J, Rushton N, Henson F. Peripheral Blood Mononuclear Cells Enhance Cartilage Repair in in vivo Osteochondral Defect Model. *PLoS One* 2015; **10**: e0133937 [PMID: 26252391 DOI: 10.1371/journal.pone.0133937]
  - 18 **Xian B**, Zhang Y, Peng Y, Huang J, Li W, Wang W, Zhang M, Li K, Zhang H, Zhao M, Liu X, Huang B. Adult Human Peripheral Blood Mononuclear Cells Are Capable of Producing Neurocyte or Photoreceptor-Like Cells That Survive in Mouse Eyes After Preinduction With Neonatal Retina. *Stem Cells Transl Med* 2016; **5**: 1515-1524 [PMID: 27458266 DOI: 10.5966/scrm.2015-0395]
  - 19 **Wang X**, Balaji S, Steen EH, Li H, Rae MM, Blum AJ, Miao Q, Butte MJ, Bollyky PL, Keswani SG. T Lymphocytes Attenuate Dermal Scarring by Regulating Inflammation, Neovascularization, and Extracellular Matrix Remodeling. *Adv Wound Care (New Rochelle)* 2019; **8**: 527-537 [PMID: 31637099 DOI: 10.1089/wound.2019.0981]
  - 20 **Wynn TA**, Vannella KM. Macrophages in Tissue Repair, Regeneration, and Fibrosis. *Immunity* 2016; **44**: 450-462 [PMID: 26982353 DOI: 10.1016/j.immuni.2016.02.015]
  - 21 **Ogle ME**, Segar CE, Sridhar S, Botchwey EA. Monocytes and macrophages in tissue repair: Implications for immunoregenerative biomaterial design. *Exp Biol Med (Maywood)* 2016; **241**: 1084-1097 [PMID: 27229903 DOI: 10.1177/1535370216650293]
  - 22 **Yang J**, Zhang L, Yu C, Yang XF, Wang H. Monocyte and macrophage differentiation: circulation inflammatory monocyte as biomarker for inflammatory diseases. *Biomark Res* 2014; **2**: 1 [PMID: 24398220 DOI: 10.1186/2050-7771-2-1]
  - 23 **Chiu S**, Bharat A. Role of monocytes and macrophages in regulating immune response following lung transplantation. *Curr Opin Organ Transplant* 2016; **21**: 239-245 [PMID: 26977996 DOI: 10.1097/MOT.0000000000000313]
  - 24 **Gordon S**, Taylor PR. Monocyte and macrophage heterogeneity. *Nat Rev Immunol* 2005; **5**: 953-964 [PMID: 16322748 DOI: 10.1038/nri1733]
  - 25 **Atri C**, Guerfali FZ, Laouini D. Role of Human Macrophage Polarization in Inflammation during Infectious Diseases. *Int J Mol Sci* 2018; **19**: 1801 [PMID: 29921749 DOI: 10.3390/ijms19061801]
  - 26 **Mosser DM**, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 2008; **8**: 958-969 [PMID: 19029990 DOI: 10.1038/nri2448]
  - 27 **Gu B**, Kaneko T, Zaw SYM, Sone PP, Murano H, Sueyama Y, Zaw ZCT, Okiji T. Macrophage populations show an M1-to-M2 transition in an experimental model of coronal pulp tissue engineering with mesenchymal stem cells. *Int Endod J* 2019; **52**: 504-514 [PMID: 30387178 DOI: 10.1111/iej.13033]
  - 28 **McWhorter FY**, Davis CT, Liu WF. Physical and mechanical regulation of macrophage phenotype and function. *Cell Mol Life Sci* 2015; **72**: 1303-1316 [PMID: 25504084 DOI: 10.1007/s00018-014-1796-8]
  - 29 **Rigamonti E**, Zordan P, Sciorati C, Rovere-Querini P, Brunelli S. Macrophage plasticity in skeletal muscle repair. *Biomed Res Int* 2014; **2014**: 560629 [PMID: 24860823 DOI: 10.1155/2014/560629]
  - 30 **Gratchev A**, Kzhyshkowska J, Köthe K, Muller-Moliniet I, Kannookadan S, Utikal J, Goerdts S. Mphi1 and Mphi2 can be re-polarized by Th2 or Th1 cytokines, respectively, and respond to exogenous danger signals. *Immunobiology* 2006; **211**: 473-486 [PMID: 16920487 DOI: 10.1016/j.imbio.2006.05.017]
  - 31 **Krzyszczuk P**, Schloss R, Palmer A, Berthiaume F. The Role of Macrophages in Acute and Chronic Wound Healing and Interventions to Promote Pro-wound Healing Phenotypes. *Front Physiol* 2018; **9**: 419 [PMID: 29765329 DOI: 10.3389/fphys.2018.00419]
  - 32 **Li J**, Tan J, Martino MM, Lui KO. Regulatory T-Cells: Potential Regulator of Tissue Repair and Regeneration. *Front Immunol* 2018; **9**: 585 [PMID: 29662491 DOI: 10.3389/fimmu.2018.00585]
  - 33 **Tiemessen MM**, Jagger AL, Evans HG, van Herwijnen MJ, John S, Taams LS. CD4+CD25+Foxp3+ regulatory T cells induce alternative activation of human monocytes/macrophages. *Proc Natl Acad Sci USA* 2007; **104**: 19446-19451 [PMID: 18042719 DOI: 10.1073/pnas.0706832104]
  - 34 **Weirather J**, Hofmann UD, Beyersdorf N, Ramos GC, Vogel B, Frey A, Ertl G, Kerkau T, Frantz S. Foxp3+ CD4+ T cells improve healing after myocardial infarction by modulating monocyte/macrophage differentiation. *Circ Res* 2014; **115**: 55-67 [PMID: 24786398 DOI: 10.1161/CIRCRESAHA.115.303895]
  - 35 **Toumi H**, F'guyer S, Best TM. The role of neutrophils in injury and repair following muscle stretch. *J Anat* 2006; **208**: 459-470 [PMID: 16637872 DOI: 10.1111/j.1469-7580.2006.00543.x]
  - 36 **Pizza FX**, Peterson JM, Baas JH, Koh TJ. Neutrophils contribute to muscle injury and impair its resolution after lengthening contractions in mice. *J Physiol* 2005; **562**: 899-913 [PMID: 15550464 DOI: 10.1113/jphysiol.2004.073965]
  - 37 **Yang Z**, Sharma AK, Linden J, Kron IL, Laubach VE. CD4+ T lymphocytes mediate acute pulmonary ischemia-reperfusion injury. *J Thorac Cardiovasc Surg* 2009; **137**: 695-702; discussion 702 [PMID: 19258091 DOI: 10.1016/j.jtcvs.2008.10.044]
  - 38 **Lindborg JA**, Mack M, Zigmond RE. Neutrophils Are Critical for Myelin Removal in a Peripheral Nerve

- Injury Model of Wallerian Degeneration. *J Neurosci* 2017; **37**: 10258-10277 [PMID: [28912156](#) DOI: [10.1523/JNEUROSCI.2085-17.2017](#)]
- 39 **Zemans RL**, Briones N, Campbell M, McClendon J, Young SK, Suzuki T, Yang IV, De Langhe S, Reynolds SD, Mason RJ, Kahn M, Henson PM, Colgan SP, Downey GP. Neutrophil transmigration triggers repair of the lung epithelium via beta-catenin signaling. *Proc Natl Acad Sci USA* 2011; **108**: 15990-15995 [PMID: [21880956](#) DOI: [10.1073/pnas.1110144108](#)]
- 40 **Saez B**, Yusuf RZ, Scadden DT. Harnessing the Biology of Stem Cells' Niche. *Biology and Engineering of Stem Cell Niches*: Elsevier; 2017: 15-31 [DOI: [10.1016/B978-0-12-802734-9.00002-0](#)]
- 41 **Mushahary D**, Spittler A, Kasper C, Weber V, Charwat V. Isolation, cultivation, and characterization of human mesenchymal stem cells. *Cytometry A* 2018; **93**: 19-31 [PMID: [29072818](#) DOI: [10.1002/cyto.a.23242](#)]
- 42 **Haque N**, Abu Kasim NH. Pooled Human Serum Increases Regenerative Potential of In Vitro Expanded Stem Cells from Human Extracted Deciduous Teeth. *Adv Exp Med Biol* 2018; **1083**: 29-44 [PMID: [28730381](#) DOI: [10.1007/5584\\_2017\\_74](#)]
- 43 **Haque N**, Widera D, Abu Kasim NH. Stem Cells from Human Extracted Deciduous Teeth Expanded in Foetal Bovine and Human Sera Express Different Paracrine Factors After Exposure to Freshly Prepared Human Serum. *Adv Exp Med Biol* 2019; **1084**: 175-186 [PMID: [30771186](#) DOI: [10.1007/5584\\_2018\\_299](#)]
- 44 **Grayson WL**, Bunnell BA, Martin E, Frazier T, Hung BP, Gimble JM. Stromal cells and stem cells in clinical bone regeneration. *Nat Rev Endocrinol* 2015; **11**: 140-150 [PMID: [25560703](#) DOI: [10.1038/nrendo.2014.234](#)]
- 45 **Trounson A**, McDonald C. Stem Cell Therapies in Clinical Trials: Progress and Challenges. *Cell Stem Cell* 2015; **17**: 11-22 [PMID: [26140604](#) DOI: [10.1016/j.stem.2015.06.007](#)]
- 46 **Gu W**, Hong X, Potter C, Qu A, Xu Q. Mesenchymal stem cells and vascular regeneration. *Microcirculation* 2017; **24** [PMID: [27681821](#) DOI: [10.1111/micc.12324](#)]
- 47 **Shi Z**, Wang Q, Jiang D. Extracellular vesicles from bone marrow-derived multipotent mesenchymal stromal cells regulate inflammation and enhance tendon healing. *J Transl Med* 2019; **17**: 211 [PMID: [31238964](#) DOI: [10.1186/s12967-019-1960-x](#)]
- 48 **Nitzsche F**, Müller C, Lukomska B, Jolkonen J, Deten A, Boltze J. Concise Review: MSC Adhesion Cascade-Insights into Homing and Transendothelial Migration. *Stem Cells* 2017; **35**: 1446-1460 [PMID: [28316123](#) DOI: [10.1002/stem.2614](#)]
- 49 **Fu X**, Liu G, Halim A, Ju Y, Luo Q, Song AG. Mesenchymal Stem Cell Migration and Tissue Repair. *Cells* 2019; **8**: 784 [PMID: [31357692](#) DOI: [10.3390/cells8080784](#)]
- 50 **Visweswaran M**, Pohl S, Arfuso F, Newsholme P, Dilley R, Pervaiz S, Dharmarajan A. Multi-lineage differentiation of mesenchymal stem cells - To Wnt, or not Wnt. *Int J Biochem Cell Biol* 2015; **68**: 139-147 [PMID: [26410622](#) DOI: [10.1016/j.biocel.2015.09.008](#)]
- 51 **Almalki SG**, Llamas Valle Y, Agrawal DK. MMP-2 and MMP-14 Silencing Inhibits VEGFR2 Cleavage and Induces the Differentiation of Porcine Adipose-Derived Mesenchymal Stem Cells to Endothelial Cells. *Stem Cells Transl Med* 2017; **6**: 1385-1398 [PMID: [28213979](#) DOI: [10.1002/sctm.16-0329](#)]
- 52 **Zhang Y**, Khan D, Delling J, Tobiasch E. Mechanisms underlying the osteo- and adipo-differentiation of human mesenchymal stem cells. *ScientificWorldJournal* 2012; **2012**: 793823 [PMID: [22500143](#) DOI: [10.1100/2012/793823](#)]
- 53 **Chen LB**, Jiang XB, Yang L. Differentiation of rat marrow mesenchymal stem cells into pancreatic islet beta-cells. *World J Gastroenterol* 2004; **10**: 3016-3020 [PMID: [15378785](#) DOI: [10.3748/wjg.v10.i20.3016](#)]
- 54 **Jung JP**, Bache-Wiig MK, Provenzano PP, Ogle BM. Heterogeneous Differentiation of Human Mesenchymal Stem Cells in 3D Extracellular Matrix Composites. *Biores Open Access* 2016; **5**: 37-48 [PMID: [26862471](#) DOI: [10.1089/biores.2015.0044](#)]
- 55 **Gruh I**, Martin U. Transdifferentiation of stem cells: a critical view. *Adv Biochem Eng Biotechnol* 2009; **114**: 73-106 [PMID: [19343303](#) DOI: [10.1007/10\\_2008\\_49](#)]
- 56 **Bassi EJ**, de Almeida DC, Moraes-Vieira PM, Câmara NO. Exploring the role of soluble factors associated with immune regulatory properties of mesenchymal stem cells. *Stem Cell Rev Rep* 2012; **8**: 329-342 [PMID: [21881832](#) DOI: [10.1007/s12015-011-9311-1](#)]
- 57 **Merino-González C**, Zuñiga FA, Escudero C, Ormazabal V, Reyes C, Nova-Lamperti E, Salomón C, Aguayo C. Mesenchymal Stem Cell-Derived Extracellular Vesicles Promote Angiogenesis: Potencial Clinical Application. *Front Physiol* 2016; **7**: 24 [PMID: [26903875](#) DOI: [10.3389/fphys.2016.00024](#)]
- 58 **Lee Y**, El Andaloussi S, Wood MJ. Exosomes and microvesicles: extracellular vesicles for genetic information transfer and gene therapy. *Hum Mol Genet* 2012; **21**: R125-R134 [PMID: [22872698](#) DOI: [10.1093/hmg/dd317](#)]
- 59 **Haque N**, Khan IM, Abu Kasim NH. Survival and immunomodulation of stem cells from human extracted deciduous teeth expanded in pooled human and foetal bovine sera. *Cytokine* 2019; **120**: 144-154 [PMID: [31071675](#) DOI: [10.1016/j.cyt.2019.04.018](#)]
- 60 **Chang PY**, Zhang BY, Cui S, Qu C, Shao LH, Xu TK, Qu YQ, Dong LH, Wang J. MSC-derived cytokines repair radiation-induced intra-villi microvascular injury. *Oncotarget* 2017; **8**: 87821-87836 [PMID: [29152123](#) DOI: [10.18632/oncotarget.21236](#)]
- 61 **de Witte SFH**, Merino AM, Franquesa M, Strini T, van Zoggel JAA, Korevaar SS, Luk F, Garghesha M, O'Flynn L, Roy D, Elliman SJ, Newsome PN, Baan CC, Hoogduijn MJ. Cytokine treatment optimises the immunotherapeutic effects of umbilical cord-derived MSC for treatment of inflammatory liver disease. *Stem Cell Res Ther* 2017; **8**: 140 [PMID: [28595619](#) DOI: [10.1186/s13287-017-0590-6](#)]
- 62 **Toh WS**, Lai RC, Hui JHP, Lim SK. MSC exosome as a cell-free MSC therapy for cartilage regeneration: Implications for osteoarthritis treatment. *Semin Cell Dev Biol* 2017; **67**: 56-64 [PMID: [27871993](#) DOI: [10.1016/j.semedb.2016.11.008](#)]
- 63 **Cantinieux D**, Quertainmont R, Blacher S, Rossi L, Wanet T, Noël A, Brook G, Schoenen J, Franzen R. Conditioned medium from bone marrow-derived mesenchymal stem cells improves recovery after spinal cord injury in rats: an original strategy to avoid cell transplantation. *PLoS One* 2013; **8**: e69515 [PMID: [24013448](#) DOI: [10.1371/journal.pone.0069515](#)]
- 64 **Motain H**, Turnsek TL. Cytokines play a key role in communication between mesenchymal stem cells and

- brain cancer cells. *Protein Pept Lett* 2015; **22**: 322-331 [PMID: [25642990](#) DOI: [10.2174/0929866522666150131123808](#)]
- 65 **Bhang SH**, Cho SW, Lim JM, Kang JM, Lee TJ, Yang HS, Song YS, Park MH, Kim HS, Yoo KJ, Jang Y, Langer R, Anderson DG, Kim BS. Locally delivered growth factor enhances the angiogenic efficacy of adipose-derived stromal cells transplanted to ischemic limbs. *Stem Cells* 2009; **27**: 1976-1986 [PMID: [19544425](#) DOI: [10.1002/stem.115](#)]
- 66 **Zhang LB**, He M. Effect of mesenchymal stromal (stem) cell (MSC) transplantation in asthmatic animal models: A systematic review and meta-analysis. *Pulm Pharmacol Ther* 2019; **54**: 39-52 [PMID: [30496803](#) DOI: [10.1016/j.pupt.2018.11.007](#)]
- 67 **Yaochite JN**, de Lima KW, Caliri-Oliveira C, Palma PV, Couri CE, Simões BP, Covas DT, Voltarelli JC, Oliveira MC, Donadi EA, Malmegrim KC. Multipotent mesenchymal stromal cells from patients with newly diagnosed type 1 diabetes mellitus exhibit preserved in vitro and in vivo immunomodulatory properties. *Stem Cell Res Ther* 2016; **7**: 14 [PMID: [26781648](#) DOI: [10.1186/s13287-015-0261-4](#)]
- 68 **Palomares Cabeza V**, Hoogduijn MJ, Kraaijeveld R, Franquesa M, Witte-Bouma J, Wolvius EB, Farrell E, Brama PAJ. Pediatric Mesenchymal Stem Cells Exhibit Immunomodulatory Properties Toward Allogeneic T and B Cells Under Inflammatory Conditions. *Front Bioeng Biotechnol* 2019; **7**: 142 [PMID: [31245368](#) DOI: [10.3389/fbioe.2019.00142](#)]
- 69 **Haque N**, Ramasamy TS, Kasim NHA. Mechanisms of Mesenchymal Stem Cells for Autoimmune Disease Treatment. In: Pham PV. *Stem Cell Transplantation for Autoimmune Diseases and Inflammation*. Cham: Springer International Publishing; 2019: 27-44 [DOI: [10.1007/978-3-030-23421-8\\_2](#)]
- 70 **Chen X**, Wang S, Cao W. Mesenchymal stem cell-mediated immunomodulation in cell therapy of neurodegenerative diseases. *Cell Immunol* 2018; **326**: 8-14 [PMID: [28778534](#) DOI: [10.1016/j.cellimm.2017.06.006](#)]
- 71 **Gebler A**, Zabel O, Seliger B. The immunomodulatory capacity of mesenchymal stem cells. *Trends Mol Med* 2012; **18**: 128-134 [PMID: [22118960](#) DOI: [10.1016/j.molmed.2011.10.004](#)]
- 72 **Coulson-Thomas VJ**, Coulson-Thomas YM, Gesteira TF, Kao WW. Extrinsic and Intrinsic Mechanisms by Which Mesenchymal Stem Cells Suppress the Immune System. *Ocul Surf* 2016; **14**: 121-134 [PMID: [26804815](#) DOI: [10.1016/j.jtos.2015.11.004](#)]
- 73 **Najar M**, Raicevic G, Fayyad-Kazan H, Bron D, Tounouz M, Lagneaux L. Mesenchymal stromal cells and immunomodulation: A gathering of regulatory immune cells. *Cytotherapy* 2016; **18**: 160-171 [PMID: [26794710](#) DOI: [10.1016/j.jcyt.2015.10.011](#)]
- 74 **Dazzi F**, Lopes L, Weng L. Mesenchymal stromal cells: a key player in 'innate tolerance'? *Immunology* 2012; **137**: 206-213 [PMID: [22804624](#) DOI: [10.1111/j.1365-2567.2012.03621.x](#)]
- 75 **Krampera M**, Glennie S, Dyson J, Scott D, Laylor R, Simpson E, Dazzi F. Bone marrow mesenchymal stem cells inhibit the response of naive and memory antigen-specific T cells to their cognate peptide. *Blood* 2003; **101**: 3722-3729 [PMID: [12506037](#) DOI: [10.1182/blood-2002-07-2104](#)]
- 76 **Romieu-Mourez R**, François M, Boivin MN, Stagg J, Galipeau J. Regulation of MHC class II expression and antigen processing in murine and human mesenchymal stromal cells by IFN-gamma, TGF-beta, and cell density. *J Immunol* 2007; **179**: 1549-1558 [PMID: [17641021](#) DOI: [10.4049/jimmunol.179.3.1549](#)]
- 77 **Bhartiya D**. The need to revisit the definition of mesenchymal and adult stem cells based on their functional attributes. *Stem Cell Res Ther* 2018; **9**: 78 [PMID: [29587828](#) DOI: [10.1186/s13287-018-0833-1](#)]
- 78 **Luger D**, Lipinski MJ, Westman PC, Glover DK, Dimastromatteo J, Frias JC, Albelda MT, Sikora S, Kharazi A, Vertelov G, Waksman R, Epstein SE. Intravenously Delivered Mesenchymal Stem Cells: Systemic Anti-Inflammatory Effects Improve Left Ventricular Dysfunction in Acute Myocardial Infarction and Ischemic Cardiomyopathy. *Circ Res* 2017; **120**: 1598-1613 [PMID: [28232595](#) DOI: [10.1161/CIRCRESAHA.117.310599](#)]
- 79 **Chen W**, Zhou D, Li YR, Liu JX, Wang XM, Zhu F, Xu KL. [Effect of Lentiviral Vector-Mediated CXCR4 Gene Overexpression on Mesenchymal Stem Cell Homing Capacity]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 2018; **26**: 1543-1547 [PMID: [30295281](#) DOI: [10.7534/j.issn.1009-2137.2018.05.047](#)]
- 80 **Jin W**, Liang X, Brooks A, Futrega K, Liu X, Doran MR, Simpson MJ, Roberts MS, Wang H. Modelling of the SDF-1/CXCR4 regulated *in vivo* homing of therapeutic mesenchymal stem/stromal cells in mice. *PeerJ* 2018; **6**: e6072 [PMID: [30564525](#) DOI: [10.7717/peerj.6072](#)]
- 81 **Sordi V**. Mesenchymal stem cell homing capacity. *Transplantation* 2009; **87**: S42-S45 [PMID: [19424004](#) DOI: [10.1097/TP.0b013e3181a28533](#)]
- 82 **Hocking AM**. The Role of Chemokines in Mesenchymal Stem Cell Homing to Wounds. *Adv Wound Care (New Rochelle)* 2015; **4**: 623-630 [PMID: [26543676](#) DOI: [10.1089/wound.2014.0579](#)]
- 83 **Sackstein R**. The biology of CD44 and HCELL in hematopoiesis: the 'step 2-bypass pathway' and other emerging perspectives. *Curr Opin Hematol* 2011; **18**: 239-248 [PMID: [21546828](#) DOI: [10.1097/MOH.0b013e31823476140](#)]
- 84 **Martinotti S**, Patrone M, Ranzato E. Emerging roles for HMGB1 protein in immunity, inflammation, and cancer. *Immunotargets Ther* 2015; **4**: 101-109 [PMID: [27471716](#) DOI: [10.2147/ITT.S58064](#)]
- 85 **Yang H**, Wang H, Chavan SS, Andersson U. High Mobility Group Box Protein 1 (HMGB1): The Prototypical Endogenous Danger Molecule. *Mol Med* 2015; **21** Suppl 1: S6-S12 [PMID: [26605648](#) DOI: [10.2119/molmed.2015.00087](#)]
- 86 **Rauci A**, Di Maggio S, Scavella F, D'Ambrosio A, Bianchi ME, Capogrossi MC. The Janus face of HMGB1 in heart disease: a necessary update. *Cell Mol Life Sci* 2019; **76**: 211-229 [PMID: [30306212](#) DOI: [10.1007/s00018-018-2930-9](#)]
- 87 **Kohno T**, Anzai T, Naito K, Miyasho T, Okamoto M, Yokota H, Yamada S, Maekawa Y, Takahashi T, Yoshikawa T, Ishizaka A, Ogawa S. Role of high-mobility group box 1 protein in post-infarction healing process and left ventricular remodelling. *Cardiovasc Res* 2009; **81**: 565-573 [PMID: [18984601](#) DOI: [10.1093/cvr/cvn291](#)]
- 88 **Andersson U**, Erlandsson-Harris H, Yang H, Tracey KJ. HMGB1 as a DNA-binding cytokine. *J Leukoc Biol* 2002; **72**: 1084-1091 [PMID: [12488489](#)]
- 89 **Frangogiannis NG**. Fibroblast-Extracellular Matrix Interactions in Tissue Fibrosis. *Curr Pathobiol Rep* 2016; **4**: 11-18 [PMID: [27171595](#) DOI: [10.1007/s40139-016-0099-1](#)]

- 90 **Klingberg F**, Hinz B, White ES. The myofibroblast matrix: implications for tissue repair and fibrosis. *J Pathol* 2013; **229**: 298-309 [PMID: [22996908](#) DOI: [10.1002/path.4104](#)]
- 91 **Zaja-Milatovic S**, Richmond A. CXC chemokines and their receptors: a case for a significant biological role in cutaneous wound healing. *Histol Histopathol* 2008; **23**: 1399-1407 [PMID: [18785122](#) DOI: [10.14670/HH-23.1399](#)]
- 92 **Behm B**, Babilas P, Landthaler M, Schreml S. Cytokines, chemokines and growth factors in wound healing. *J Eur Acad Dermatol Venereol* 2012; **26**: 812-820 [PMID: [22211801](#) DOI: [10.1111/j.1468-3083.2011.04415.x](#)]
- 93 **Wang H**, Bloom O, Zhang M, Vishnubhakat JM, Ombrellino M, Che J, Frazier A, Yang H, Ivanova S, Borovikova L, Manogue KR, Faist E, Abraham E, Andersson J, Andersson U, Molina PE, Abumrad NN, Sama A, Tracey KJ. HMG-1 as a late mediator of endotoxin lethality in mice. *Science* 1999; **285**: 248-251 [PMID: [10398600](#) DOI: [10.1126/science.285.5425.248](#)]
- 94 **Andersson U**, Wang H, Palmblad K, Aveberger AC, Bloom O, Erlandsson-Harris H, Janson A, Kokkola R, Zhang M, Yang H, Tracey KJ. High mobility group 1 protein (HMG-1) stimulates proinflammatory cytokine synthesis in human monocytes. *J Exp Med* 2000; **192**: 565-570 [PMID: [10952726](#) DOI: [10.1084/jem.192.4.565](#)]
- 95 **Kokkola R**, Andersson A, Mullins G, Ostberg T, Treutiger CJ, Arnold B, Nawroth P, Andersson U, Harris RA, Harris HE. RAGE is the major receptor for the proinflammatory activity of HMGB1 in rodent macrophages. *Scand J Immunol* 2005; **61**: 1-9 [PMID: [15644117](#) DOI: [10.1111/j.0300-9475.2005.01534.x](#)]
- 96 **Park JS**, Arcaroli J, Yum HK, Yang H, Wang H, Yang KY, Choe KH, Strassheim D, Pitts TM, Tracey KJ, Abraham E. Activation of gene expression in human neutrophils by high mobility group box 1 protein. *Am J Physiol Cell Physiol* 2003; **284**: C870-C879 [PMID: [12620891](#) DOI: [10.1152/ajpcell.00322.2002](#)]
- 97 **Palumbo R**, De Marchis F, Pusterla T, Conti A, Alessio M, Bianchi ME. Src family kinases are necessary for cell migration induced by extracellular HMGB1. *J Leukoc Biol* 2009; **86**: 617-623 [PMID: [19401391](#) DOI: [10.1189/jlb.0908581](#)]
- 98 **Palumbo R**, Galvez BG, Pusterla T, De Marchis F, Cossu G, Marcu KB, Bianchi ME. Cells migrating to sites of tissue damage in response to the danger signal HMGB1 require NF-kappaB activation. *J Cell Biol* 2007; **179**: 33-40 [PMID: [17923528](#) DOI: [10.1083/jcb.200704015](#)]
- 99 **Yang J**, Chen L, Yang J, Ding J, Rong H, Dong W, Li X. High mobility group box-1 induces migration of vascular smooth muscle cells via TLR4-dependent PI3K/Akt pathway activation. *Mol Biol Rep* 2012; **39**: 3361-3367 [PMID: [21698364](#) DOI: [10.1007/s11033-011-1106-6](#)]
- 100 **Xue X**, Chen X, Fan W, Wang G, Zhang L, Chen Z, Liu P, Liu M, Zhao J. High-mobility group box 1 facilitates migration of neural stem cells via receptor for advanced glycation end products signaling pathway. *Sci Rep* 2018; **8**: 4513 [PMID: [29540727](#) DOI: [10.1038/s41598-018-22672-4](#)]
- 101 **Shirozu M**, Nakano T, Inazawa J, Tashiro K, Tada H, Shinohara T, Honjo T. Structure and chromosomal localization of the human stromal cell-derived factor 1 (SDF1) gene. *Genomics* 1995; **28**: 495-500 [PMID: [7490086](#) DOI: [10.1006/geno.1995.1180](#)]
- 102 **Grunewald M**, Avraham I, Dor Y, Bachar-Lustig E, Itin A, Jung S, Chimenti S, Landsman L, Abramovitch R, Keshet E. VEGF-induced adult neovascularization: recruitment, retention, and role of accessory cells. *Cell* 2006; **124**: 175-189 [PMID: [16413490](#) DOI: [10.1016/j.cell.2005.10.036](#)]
- 103 **Ratajczak MZ**, Kim CH, Abdel-Latif A, Schneider G, Kucia M, Morris AJ, Laughlin MJ, Ratajczak J. A novel perspective on stem cell homing and mobilization: review on bioactive lipids as potent chemoattractants and cationic peptides as underappreciated modulators of responsiveness to SDF-1 gradients. *Leukemia* 2012; **26**: 63-72 [PMID: [21886175](#) DOI: [10.1038/leu.2011.242](#)]
- 104 **Cheng M**, Qin G. Progenitor cell mobilization and recruitment: SDF-1, CXCR4,  $\alpha$ 4-integrin, and c-kit. *Prog Mol Biol Transl Sci* 2012; **111**: 243-264 [PMID: [22917234](#) DOI: [10.1016/B978-0-12-398459-3.00011-3](#)]
- 105 **Burns JM**, Summers BC, Wang Y, Melikian A, Berahovich R, Miao Z, Penfold ME, Sunshine MJ, Littman DR, Kuo CJ, Wei K, McMaster BE, Wright K, Howard MC, Schall TJ. A novel chemokine receptor for SDF-1 and I-TAC involved in cell survival, cell adhesion, and tumor development. *J Exp Med* 2006; **203**: 2201-2213 [PMID: [16940167](#) DOI: [10.1084/jem.20052144](#)]
- 106 **Ponte AL**, Marais E, Gallay N, Langonné A, Delorme B, Héralut O, Charbord P, Domenech J. The in vitro migration capacity of human bone marrow mesenchymal stem cells: comparison of chemokine and growth factor chemotactic activities. *Stem Cells* 2007; **25**: 1737-1745 [PMID: [17395768](#) DOI: [10.1634/stemcells.2007-0054](#)]
- 107 **Gauley J**, Pisetsky DS. The translocation of HMGB1 during cell activation and cell death. *Autoimmunity* 2009; **42**: 299-301 [PMID: [19811282](#) DOI: [10.1080/08916930902831522](#)]
- 108 **Tang Z**, Jiang M, Ou-Yang Z, Wu H, Dong S, Hei M. High mobility group box 1 protein (HMGB1) as biomarker in hypoxia-induced persistent pulmonary hypertension of the newborn: a clinical and *in vivo* pilot study. *Int J Med Sci* 2019; **16**: 1123-1131 [PMID: [31523175](#) DOI: [10.7150/ijms.34344](#)]
- 109 **Zhao Y**, Yi J, Tao L, Huang G, Chu X, Song H, Chen L. Wnt signaling induces radioresistance through upregulating HMGB1 in esophageal squamous cell carcinoma. *Cell Death Dis* 2018; **9**: 433 [PMID: [29567990](#) DOI: [10.1038/s41419-018-0466-4](#)]
- 110 **Yadav SS**, Kumar M, Varshney A, Yadava PK. KLF4 sensitizes the colon cancer cell HCT-15 to cisplatin by altering the expression of HMGB1 and hTERT. *Life Sci* 2019; **220**: 169-176 [PMID: [30716337](#) DOI: [10.1016/j.lfs.2019.02.005](#)]
- 111 **Foglio E**, Pellegrini L, Germani A, Russo MA, Limana F. HMGB1-mediated apoptosis and autophagy in ischemic heart diseases. *Vasc Biol* 2019; **1**: H89-H96 [DOI: [10.1530/VB-19-0013](#)]
- 112 **Penzo M**, Habel DM, Ramadass M, Kew RR, Marcu KB. Cell migration to CXCL12 requires simultaneous IKK $\alpha$  and IKK $\beta$ -dependent NF- $\kappa$ B signaling. *Biochim Biophys Acta* 2014; **1843**: 1796-1804 [PMID: [24747690](#) DOI: [10.1016/j.bbamer.2014.04.011](#)]
- 113 **Penzo M**, Molteni R, Suda T, Samaniego S, Raucci A, Habel DM, Miller F, Jiang HP, Li J, Pardi R, Palumbo R, Olivetto E, Kew RR, Bianchi ME, Marcu KB. Inhibitor of NF-kappa B kinases alpha and beta are both essential for high mobility group box 1-mediated chemotaxis [corrected]. *J Immunol* 2010; **184**: 4497-4509 [PMID: [20231695](#) DOI: [10.4049/jimmunol.0903131](#)]



- 114 **Lolmede K**, Campana L, Vezzoli M, Bosurgi L, Tonlorenzi R, Clementi E, Bianchi ME, Cossu G, Manfredi AA, Brunelli S, Rovere-Querini P. Inflammatory and alternatively activated human macrophages attract vessel-associated stem cells, relying on separate HMGB1- and MMP-9-dependent pathways. *J Leukoc Biol* 2009; **85**: 779-787 [PMID: [19197071](#) DOI: [10.1189/jlb.0908579](#)]
- 115 **Kew RR**, Penzo M, Habel DM, Marcu KB. The IKK $\alpha$ -dependent NF- $\kappa$ B p52/RelB noncanonical pathway is essential to sustain a CXCL12 autocrine loop in cells migrating in response to HMGB1. *J Immunol* 2012; **188**: 2380-2386 [PMID: [22287708](#) DOI: [10.4049/jimmunol.1102454](#)]
- 116 **Degryse B**, Bonaldi T, Scaffidi P, Müller S, Resnati M, Sanvito F, Arrighi G, Bianchi ME. The high mobility group (HMG) boxes of the nuclear protein HMGB1 induce chemotaxis and cytoskeleton reorganization in rat smooth muscle cells. *J Cell Biol* 2001; **152**: 1197-1206 [PMID: [11257120](#) DOI: [10.1083/jcb.152.6.1197](#)]
- 117 **Rauvala H**, Rouhiainen A. Physiological and pathophysiological outcomes of the interactions of HMGB1 with cell surface receptors. *Biochim Biophys Acta* 2010; **1799**: 164-170 [PMID: [19914413](#) DOI: [10.1016/j.bbagr.2009.11.012](#)]
- 118 **Kofuku Y**, Yoshiura C, Ueda T, Terasawa H, Hirai T, Tominaga S, Hirose M, Maeda Y, Takahashi H, Terashima Y, Matsushima K, Shimada I. Structural basis of the interaction between chemokine stromal cell-derived factor-1/CXCL12 and its G-protein-coupled receptor CXCR4. *J Biol Chem* 2009; **284**: 35240-35250 [PMID: [19837984](#) DOI: [10.1074/jbc.M109.024851](#)]
- 119 **Venereau E**, Schiraldi M, Uguccioni M, Bianchi ME. HMGB1 and leukocyte migration during trauma and sterile inflammation. *Mol Immunol* 2013; **55**: 76-82 [PMID: [23207101](#) DOI: [10.1016/j.molimm.2012.10.037](#)]
- 120 **Lee G**, Espirito Santo AI, Zwingenberger S, Cai L, Vogl T, Feldmann M, Horwood NJ, Chan JK, Nanchahal J. Fully reduced HMGB1 accelerates the regeneration of multiple tissues by transitioning stem cells to G<sub>Alert</sub>. *Proc Natl Acad Sci USA* 2018; **115**: E4463-E4472 [PMID: [29674451](#) DOI: [10.1073/pnas.1802893115](#)]
- 121 **Liu H**, Liu S, Li Y, Wang X, Xue W, Ge G, Luo X. The role of SDF-1-CXCR4/CXCR7 axis in the therapeutic effects of hypoxia-preconditioned mesenchymal stem cells for renal ischemia/reperfusion injury. *PLoS One* 2012; **7**: e34608 [PMID: [22511954](#) DOI: [10.1371/journal.pone.0034608](#)]
- 122 **Li M**, Sun X, Ma L, Jin L, Zhang W, Xiao M, Yu Q. SDF-1/CXCR4 axis induces human dental pulp stem cell migration through FAK/PI3K/Akt and GSK3 $\beta$ / $\beta$ -catenin pathways. *Sci Rep* 2017; **7**: 40161 [PMID: [28067275](#) DOI: [10.1038/srep40161](#)]
- 123 **Lau TT**, Wang DA. Stromal cell-derived factor-1 (SDF-1): homing factor for engineered regenerative medicine. *Expert Opin Biol Ther* 2011; **11**: 189-197 [PMID: [21219236](#) DOI: [10.1517/14712598.2011.546338](#)]
- 124 **Vénereau E**, Ceriotti C, Bianchi ME. DAMPs from Cell Death to New Life. *Front Immunol* 2015; **6**: 422 [PMID: [26347745](#) DOI: [10.3389/fimmu.2015.00422](#)]
- 125 **Zhao W**, Jin K, Li J, Qiu X, Li S. Delivery of stromal cell-derived factor 1 $\alpha$  for *in situ* tissue regeneration. *J Biol Eng* 2017; **11**: 22 [PMID: [28670340](#) DOI: [10.1186/s13036-017-0058-3](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

