

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2020 August 15; 12(8): 791-941



REVIEW

- 791 Gastrointestinal neuroendocrine tumors in 2020
Ahmed M
- 808 Early stage colon cancer: Current treatment standards, evolving paradigms, and future directions
Chakrabarti S, Peterson CY, Sriram D, Mahipal A
- 833 One size does not fit all for pancreatic cancers: A review on rare histologies and therapeutic approaches
Niger M, Prisciandaro M, Antista M, Monica MAT, Cattaneo L, Prinzi N, Manglaviti S, Nichetti F, Brambilla M, Torchio M, Corti F, Pusceddu S, Coppa J, Mazzaferro V, de Braud F, Di Bartolomeo M

MINIREVIEWS

- 850 Gastric neuroendocrine tumor: A practical literature review
Roberto GA, Rodrigues CMB, Peixoto RD, Younes RN

ORIGINAL ARTICLE**Basic Study**

- 857 Identification of an immune-related gene-based signature to predict prognosis of patients with gastric cancer
Qiu XT, Song YC, Liu J, Wang ZM, Niu X, He J
- 877 Interleukin-1 receptor antagonist enhances chemosensitivity to fluorouracil in treatment of Kras mutant colon cancer
Yan Y, Lin HW, Zhuang ZN, Li M, Guo S

Case Control Study

- 893 Clinical and pathological characteristics and prognosis of 132 cases of rectal neuroendocrine tumors
Yu YJ, Li YW, Shi Y, Zhang Z, Zheng MY, Zhang SW

Retrospective Cohort Study

- 903 Comparison of hyperthermic intraperitoneal chemotherapy regimens for treatment of peritoneal-metastasized colorectal cancer
Spiegelberg J, Neeff H, Holzner P, Runkel M, Fichtner-Feigl S, Glatz T

Retrospective Study

- 918 Endoscopic mucosal resection *vs* endoscopic submucosal dissection for superficial non-ampullary duodenal tumors
Esaki M, Haraguchi K, Akahoshi K, Tomoeda N, Aso A, Itaba S, Ogino H, Kitagawa Y, Fujii H, Nakamura K, Kubokawa M, Harada N, Minoda Y, Suzuki S, Ihara E, Ogawa Y

- 931 Accurate ultrasonography-based portal pressure assessment in patients with hepatocellular carcinoma
Zhang Y, Wang Z, Yue ZD, Zhao HW, Wang L, Fan ZH, Wu YF, He FL, Liu FQ

ABOUT COVER

Editorial board member of *World Journal of Gastrointestinal Oncology*, Dr. Cao is an Assistant Research Fellow at the Second Affiliated Hospital of Soochow University in Suzhou, China. Having received his Bachelor's degree from Soochow University in 2006, Dr. Cao undertook his postgraduate training at Soochow University, receiving his Master's degree in 2009 and his PhD in 2015. He works in the Department of Pathology, the Second Affiliated Hospital of Soochow University, Soochow University, where his ongoing research interests involve the molecular pathological mechanisms of malignant tumors, particularly in relation to oncogenesis, radiation resistance and cellular signal transduction involving tumors of the digestive system. To date, he has published 16 SCI papers as corresponding author or first author.

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Oncology* (*WJGO*, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

INDEXING/ABSTRACTING

The *WJGO* is now indexed in Science Citation Index Expanded (also known as SciSearch®), PubMed, and PubMed Central. The 2020 edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJGO* as 2.898; IF without journal self cites: 2.880; 5-year IF: 3.316; Ranking: 143 among 244 journals in oncology; Quartile category: Q3; Ranking: 55 among 88 journals in gastroenterology and hepatology; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Li-Li Wang; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Rosa M Jimenez Rodriguez, Pashtoon Kasi, Monjur Ahmed

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

PUBLICATION DATE

August 15, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Gastric neuroendocrine tumor: A practical literature review

Gabriel Antonio Roberto, Carolina Magalhães Britto Rodrigues, Renata D'Alpino Peixoto, Riad Naim Younes

ORCID number: Gabriel Antonio Roberto 0000-0001-9814-9038; Carolina Magalhães Britto Rodrigues 0000-0003-4666-7065; Renata D'Alpino Peixoto 0000-0003-0053-7951; Riad Naim Younes 0000-0001-9844-6390.

Conflict-of-interest statement: Authors declare no conflict of interests for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: March 20, 2020

Peer-review started: March 20, 2020

First decision: April 26, 2020

Revised: May 5, 2020

Accepted: July 19, 2020

Article in press: July 19, 2020

Published online: August 15, 2020

P-Reviewer: Fernandes J

Gabriel Antonio Roberto, Carolina Magalhães Britto Rodrigues, Bachelor Medicine, União das Faculdades dos Grandes Lagos, São José do Rio Preto 15030070, Brazil

Renata D'Alpino Peixoto, Department of Oncology, Hospital Alemão Oswaldo Cruz, São Paulo 01327001, Brazil

Renata D'Alpino Peixoto, Universidade Nove de Julho, São Paulo 01327001, Brazil

Riad Naim Younes, Department of Thoracic Surgery, Hospital Alemão Oswaldo Cruz, São Paulo 01327001, Brazil

Corresponding author: Gabriel Antonio Roberto, BM BCh, Bmed, Academic Research, Bachelor Medicine, União das Faculdades dos Grandes Lagos, Rua Dr. Eduardo Nielsem, 960 Jd. N. Aeroporto, São Jose do Rio Preto 15030070, Brazil. contatogabrielroberto@gmail.com

Abstract

Gastric neuroendocrine tumors are gastric neoplasms originating from enterochromaffin type cells and are inserted in a larger group, named gastroenteropancreatic neuroendocrine tumors. They are considered rare and variable in terms of their clinical, morphological and functional characteristics and may be indolent or aggressive. They are classified into types I, II and III, according to their pathophysiology, behavior and treatment. Their diagnosis occurs, in most cases, incidentally during upper digestive endoscopies, presenting as simple gastric polyps. Most cases (type I and type II) are related to hypergastrinemia, can be multiple and are treated by endoscopic resection, whenever possible. The use of somatostatin analogs for tumor control may be one of the options for therapy, in addition to total or subtotal gastrectomy for selected cases. Adjuvant chemotherapy is only reserved for poorly differentiated neuroendocrine carcinomas. Although rare, gastric neuroendocrine tumors have an increasing incidence over the years, therefore deserving more comprehensive studies on its adequate treatment. The present study reviews and updates management recommendations for gastric neuroendocrine tumors.

Key words: Gastric neuroendocrine tumor; Gastroenteropancreatic tumor; Hypergastrinemia; Gastric carcinoid; Endoscopic resection

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Gastric neuroendocrine tumors are rare lesions that are part of the

S-Editor: Zhang L
L-Editor: Filipodia
P-Editor: Wang LL



gastroenteropancreatic neuroendocrine tumors group. They are classified into types I, II and III according to their clinical and pathophysiological characteristics. Their diagnosis is usually made incidentally by upper gastrointestinal endoscopy, and most cases are treated by endoscopic resection. Surgical resections, as well as somatostatin analog treatments, are reserved for selected cases. Although rare, gastric neuroendocrine tumors need further research as their incidence has increased over the years.

Citation: Roberto GA, Rodrigues CMB, Peixoto RD, Younes RN. Gastric neuroendocrine tumor: A practical literature review. *World J Gastrointest Oncol* 2020; 12(8): 850-856

URL: <https://www.wjgnet.com/1948-5204/full/v12/i8/850.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v12.i8.850>

INTRODUCTION

Gastric neuroendocrine tumors (G-NETs), once called gastric carcinoids, are neoplasms derived from enterochromaffin-like cells (ECF) of the stomach mucosa and correspond to less than 2% of all gastric neoplasms^[1]. They are part of a larger group called gastroenteropancreatic NETs (GEP-NET), which comprise well-differentiated NETs from the gastrointestinal tract. Well-differentiated NETs, together with poorly differentiated neuroendocrine carcinomas (NECs) form the neuroendocrine neoplasms. In immunohistochemistry, like other GEP-NET, G-NETs usually express neuroendocrine markers, such as chromogranin and synaptophysin. They are considered rare and of heterogeneous spectrum with a wide variety of morphological, functional and clinical characteristics^[2-4]. Their behavior is generally indolent, although may be highly aggressive^[5].

The real prevalence of NETs is unknown due to a worldwide difficulty in standardizing and categorizing the data. Nonetheless, increasing incidence over time is certainly related to a greater access to endoscopic and imaging methods, favoring its diagnosis^[1,6-9]. A 2015 multicenter study involving national registries from several countries estimated that the prevalence of G-NET in Europe is 0.32 per 10000 inhabitants, while in the United States it is 0.17 and 0.05 in Japan^[10]. Most G-NETs are incidentally diagnosed as simple gastric polyps during endoscopies of the upper gastrointestinal tract, corresponding to 0.6% to 2% of gastric polyp cases^[6,9,11-16].

The present review of the English literature presents updated definitions as well as epidemiology, diagnosis and management recommendations for G-NET.

DISCUSSION

In order to standardize the classification of GEP-NETs and facilitate their understanding, the World Health Organization in 2010 divided GEP-NETs (including G-NETs) into three histological grades (G1, G2 and G3) based on the mitotic index (number of mitoses per ten high magnification fields) and/or on the Ki-67 index (mitotic and cellular proliferative index) (Table 1). This division was important due to the clinical and prognostic variability between G1, G2 and G3 groups. G1 and G2 GEP-NETs were considered well differentiated while high-grade NECs (G3) were considered poorly differentiated with significantly more aggressive behavior. In 2019, World Health Organization revised the classification and recognized a new category of high-grade but still well-differentiated GEP-NET (G3 NET-Neuroendocrine Tumors) (Table 2). Unlike G3 NECs, G3 NETs usually have a Ki-67 index below 55% and a prognosis not as poor as G3 NECs^[17]. In addition to the grade classification established by the World Health Organization, which is fundamental for all GEP-NETs, well-differentiated G-NETs are also clinically divided into three types according to their pathophysiology and behavior, which influences treatment recommendations (Table 3).

Below we will describe the three types of G-NETs with their clinical characteristics and approach to localized disease.

Type I

Type I tumors correspond to the majority of G-NETs. They constitute about 70%-80%

Table 1 Classification of gastroenteropancreatic neuroendocrine tumors according to the World Health Organization 2010

	Grade I	Grade II	Grade III
Tumor size in cm	≤ 2	> 2	Any
Mitoses/10 HPF	< 2	2–20	> 20
Ki 67 index, %	< 3	3–20	> 20
Differentiation	Well differentiated	Well differentiated	Poorly differentiated

Adapted from^[18]. HPF: High-power fields.

Table 2 Classification of gastroenteropancreatic neuroendocrine tumors: Neuroendocrine neoplasms according to the World Health Organization 2019

Terminology	Differentiation	Grade	Mitotic rate	Ki 67 index, %
NET, G1	Well differentiated	Low	< 2	< 3
NET, G2	Well differentiated	Intermediate	2-20	3-20
NET, G3	Well differentiated	High	> 20	> 20
NEC, SCNEC	Poorly differentiated	High	> 20	> 20
NEC, LCNEC	Poorly differentiated	High	> 20	> 20
MiNEN	Well or poorly differentiated	Variable	Variable	Variable

Adapted from^[17]. NET: Neuroendocrine tumor; NEC: Neuroendocrine carcinoma; SCNEC: Small cell neuroendocrine carcinoma; LCNEC: Large cell neuroendocrine carcinoma; MiNEN: Mixed neuroendocrine non-neuroendocrine neoplasm.

of the lesions and are associated with chronic autoimmune atrophic gastritis^[18-23].

The destruction of parietal cells leads to achlorhydria, which stimulates the production of gastrin. This results in hypergastrinemia as a physiological response to the demand generated by the shortage of HCl. The excess of gastrin generates hypertrophy and hyperplasia of the ECFs, favoring the appearance of innumerable small lesions, which are usually not very aggressive^[18,20,22,23]. Serum gastrin is always elevated in type I G-NETs. Vitamin B12 deficiency with or without macrocytic anemia (pernicious or megaloblastic) may be present due to the reduction of the intrinsic factor, with a consequent reduction in the absorption of vitamin B12^[18,20,22-24]. Parallel to this, serum antiparietal cell antibodies are positive in 80% of cases^[20,24-26].

The diagnosis is made by upper digestive endoscopy with biopsy. There are pale, yellowish and transparent blood vessels that contrast with the smooth and red mucosa of areas not affected by the tumor, presenting as red, small and numerous polyps^[11,19,20,22,24-26]. Histological analysis of the gastric mucosa shows atrophy of mucosal cells, hyperplasia of neuroendocrine cells and absence of parietal cells.

For type I G-NETs, treatment is generally more conservative to avoid gastrectomy because they are smaller and more defined lesions. The prognosis is good. The treatment of choice is endoscopic resection for lesions ≥ 0.5 cm and endoscopic observation in smaller ones. In lesions smaller than 2 cm, the risk of metastasis is less than 10%^[27]. In general, for lesions smaller than 1 cm, no other complementary imaging exam is necessary. However, for lesions ≥ 1 cm, echo-endoscopy is recommended to identify the depth of tumor invasion in the gastric wall and the possible involvement of regional lymph nodes. Gastrectomy is reserved for submucosa tumors and/or lymph node involvement and/or positive margin in the polypectomy sample^[19,22,28]. Patients with small type I G-NETs are managed by regular endoscopic follow-up.

When the lesions are multiple and impossible to resect endoscopically or when there are repeated recurrences after endoscopic treatment, both gastrectomy and prescription of somatostatin analogs can be used to reduce serum gastrin and tumor control^[29,30]. Reports of the use of somatostatin analogues in small groups of patients showed that the interruption after 12 mo caused the serum gastrin to rise again without the reappearance of new lesions^[11,31,32]. However, data are still insufficient to show the long-term efficacy of pharmacological treatment of localized type I G-NETs^[21,22]. More rarely, antrectomy may be indicated in an attempt to reduce

Table 3 Types of gastric neuroendocrine tumors

	Type I	Type II	Type III
Prevalence, %	70-80	5-10	10-20
Background	Chronic atrophic gastritis	Gastrinomas (Zollinger-Ellison syndrome)	Normal mucosa
Other syndromes	Autoimmune polyglandular syndrome	MEN-1 syndrome	
Number of lesions	Multiple	Multiple	Single
Site of tumor	Fundus/body	Fundus/body	Fundus/body
Cell of origin	ECL	ECL	ECL, EC or X cell
Serum gastrin levels	Elevated	Elevated	Normal
Gastric PH	High	Low	Normal
Underlying mucosa	Atrophic	Hypertrophic	Normal
Size of tumors, usual	1-2 cm	1 cm	> 2 cm
Invasion	Rare	More common	Common
Metastases			
Lymph nodes	5%-10%	10%-20% (duodenal tumors)	50%-100%
Liver	2%-5%	10%	22%-75%
Prognosis	Excellent	Very good	Similar to gastric adenocarcinoma

Adapted from^[18]. ECL: Enterochromaffin-like; EC: Endocrine.

hypergastrinemia.

Type II

They correspond to 5%-10% of G-NETs. In type II, hypergastrinemia also occurs, but it is due to the presence of Zollinger-Ellison syndrome mostly in the context of MEN-1 syndrome. Therefore, in the suspicion of a type II G-NET, it is recommended to determine the serum concentration of both pituitary and parathyroid hormones as well as serum calcium and gastrin levels to assess the possibility of MEN-1 syndrome. The patient may experience abdominal pain and diarrhea in addition to peptic ulcers. Similar to type I G-NETs, excess gastrin causes hypertrophy and hyperplasia of the ECFs. In these cases, it is also common for lesions to be small and multiple^[2,18,33-35].

Upon diagnosis, upper endoscopy reveals normal or hypertrophic gastric mucosa in addition to hypergastrinemia and low pH due to hyperchlorhydria. Unlike type I, type II G-NETs tend to be slightly larger, affect younger patients and have a slightly worse prognosis with the risk of lymph node metastases reaching 30%^[27].

In general, the management of type II G-NETs is similar to type I, except for the need to also resect the gastrinoma. Most cases are treated endoscopically with resections. Surgery is rarely necessary. The use of somatostatin analogues is still debated as well as in type I G-NETs^[20,22].

When confirming the diagnosis, the primary gastrinoma should be located and resected, although it is not always possible to locate it and multiple lesions may exist. For that, we include computed tomography, magnetic resonance imaging, endoscopic ultrasound, scintigraphy with octreotide, selective angiography, positron emission tomography and/or intraoperative ultrasound in the workup. It is also possible to use an anatomical reference known as the gastrinoma triangle composed of the junction of the cystic duct with the common liver, the transition from the second to the third duodenal portion and the pancreatic neck^[11,20,35].

Type III

G-NETs of this type are sporadic and not associated with any known clinical condition. They correspond to 10%-15% of all G-NETs. The production of gastrin and HCl is within normal values, except in rare cases where the tumor itself can produce gastrin^[36]. They are generally characterized by being single lesions, larger than 1 cm in size and with greater likelihood of evolving to regional and systemic metastases^[2,20,33,34]. More than half of patients with type III G-NET are metastatic at

diagnosis, mainly to the liver. In these cases, carcinoid syndrome may be present, which is a paraneoplastic syndrome caused by endogenous secretion of serotonin and kallikrein secondary to carcinoid tumors. It becomes manifest when those vasoactive substances from the tumors enter the systemic circulation escaping hepatic degradation. Clinical components of the typical carcinoid syndrome are flushing, diarrhea and abdominal pain. It occurs more frequently in the context of high-volume hepatic metastases and primary tumors located in the small bowel, although it may happen with G-NETs, when atypical symptoms, such as bronchoconstriction, may be present due to the release of histamine.

Recently, some groups have suggested the existence of a type IV G-NET, which consists of the same characteristics described above for type III but being neuroendocrine carcinomas or mixed neuroendocrine non-neuroendocrine neoplasm. Therefore, they have a more aggressive behavior and even worse prognosis^[37]. However, the subclassification of type IV is still not well established.

Type III lesions are also investigated by upper endoscopy with biopsy, which shows a single lesion with normal mucosa. The pH is < 4, which is normal for the gastric pattern^[2,18,20,33,34]. In addition to the neoplastic lesion, the adjacent normal mucosa should also be biopsied in order to assess whether there is atrophic gastritis, intestinal metaplasia and ECF hyperplasia, which are not usually present^[2,18,19,33,34].

Total or subtotal gastrectomy is performed together with lymphadenectomy, as recommended in gastric adenocarcinomas^[22,38]. For patients with any surgical contraindication, endoscopic resection may be an alternative, but the risk of regional lymph node spread is high. When the anatomopathological part of the resection specimen shows a slightly differentiated NEC, adjuvant chemotherapy based on platinum, such as cisplatin and etoposide, is used (similar to small-cell lung carcinomas).

Treatment of metastatic disease

The goal of metastatic G-NET therapy is to control symptoms by reducing circulating hormones (when present) and tumor growth in order to increase quality of life and ensure greater survival^[39]. In general, the treatment of well-differentiated metastatic disease (G1, G2 or G3 NET) is usually similar to other NETs, taking into account the patient's performance, available drugs, toxicity profile, the volume and extent of the metastatic disease, the expression of somatostatin receptors in functional images (Octreoscan or ⁶⁸Ga-Dotatate) and the presence/lack of a functioning syndrome. Surgical resection of metastases, local-regional therapies such as embolization or ablation when there is exclusive liver involvement, somatostatin analogs, target-molecular drugs (everolimus), ¹⁷⁷Lu-Octreotate or even chemotherapy regimens when G3 should be considered when possible^[40]. Despite the low response rates, the somatostatin analogue (Octreotide or Lanreotide) is usually the initial treatment of choice because it is well tolerated^[41,42]. In the presence of carcinoid syndrome (8% to 35% of G-NETs), the use of the somatostatin analog is mandatory to reduce symptoms and decrease the long-term risks of an uncontrolled carcinoid syndrome. The ideal sequencing for patients with G-NETs, as in other NETs, remains unknown.

In the case of metastatic NEC, the treatment usually follows the protocols of small-cell lung carcinomas, in which the most commonly administered regimen is the combination of cisplatin and etoposide^[43]. In these cases, despite good initial response rates, the prognosis is often poor.

CONCLUSION

Although relatively rare, the incidence of G-NETs has increased over time. They comprise a diverse entity of three subtypes with different pathophysiology, prognosis and management. Further studies are needed for further advances in the treatment of G-NETs.

REFERENCES

- 1 **Strosberg JR**, Benson AB, Huynh L, Duh MS, Goldman J, Sahai V, Rademaker AW, Kulke MH. Clinical benefits of above-standard dose of octreotide LAR in patients with neuroendocrine tumors for control of carcinoid syndrome symptoms: a multicenter retrospective chart review study. *Oncologist* 2014; **19**: 930-936 [PMID: 25096997 DOI: 10.1634/theoncologist.2014-0120]
- 2 **Berruti A**, Fazio N, Ferrero A, Brizzi MP, Volante M, Nobili E, Tozzi L, Bodei L, Torta M, D'Avolio A,

- Priola AM, Birocco N, Amoroso V, Biasco G, Papotti M, Dogliotti L. Bevacizumab plus octreotide and metronomic capecitabine in patients with metastatic well-to-moderately differentiated neuroendocrine tumors: the XELBEVOCT study. *BMC Cancer* 2014; **14**: 184 [PMID: [24628963](#) DOI: [10.1186/1471-2407-14-184](#)]
- 3 **Wolin EM**, Jarzab B, Eriksson B, Walter T, Toumpanakis C, Morse MA, Tomassetti P, Weber MM, Fogelman DR, Ramage J, Poon D, Gadbow B, Li J, Pasiaka JL, Mahamat A, Swahn F, Newell-Price J, Mansoor W, Öberg K. Phase III study of pasireotide long-acting release in patients with metastatic neuroendocrine tumors and carcinoid symptoms refractory to available somatostatin analogues. *Drug Des Devel Ther* 2015; **9**: 5075-5086 [PMID: [26366058](#) DOI: [10.2147/DDDT.S84177](#)]
 - 4 **Yamaguchi T**, Machida N, Morizane C, Kasuga A, Takahashi H, Sudo K, Nishina T, Tobimatsu K, Ishido K, Furuse J, Boku N, Okusaka T. Multicenter retrospective analysis of systemic chemotherapy for advanced neuroendocrine carcinoma of the digestive system. *Cancer Sci* 2014; **105**: 1176-1181 [PMID: [24975505](#) DOI: [10.1111/cas.12473](#)]
 - 5 **Yao JC**, Guthrie KA, Moran C, Strosberg JR, Kulke MH, Chan JA, LoConte N, McWilliams RR, Wolin EM, Mattar B, McDonough S, Chen H, Blanke CD, Hochster HS. Phase III Prospective Randomized Comparison Trial of Depot Octreotide Plus Interferon Alfa-2b Versus Depot Octreotide Plus Bevacizumab in Patients With Advanced Carcinoid Tumors: SWOG S0518. *J Clin Oncol* 2017; **35**: 1695-1703 [PMID: [28384065](#) DOI: [10.1200/JCO.2016.70.4072](#)]
 - 6 **Al-Efraij K**, Aljama MA, Kennecke HF. Association of dose escalation of octreotide long-acting release on clinical symptoms and tumor markers and response among patients with neuroendocrine tumors. *Cancer Med* 2015; **4**: 864-870 [PMID: [25727756](#) DOI: [10.1002/cam4.435](#)]
 - 7 **Bajetta E**, Catena L, Fazio N, Puseddu S, Biondani P, Blanco G, Ricci S, Aieta M, Pucci F, Valente M, Bianco N, Mauri CM, Spada F. Everolimus in combination with octreotide long-acting repeatable in a first-line setting for patients with neuroendocrine tumors: an ITMO group study. *Cancer* 2014; **120**: 2457-2463 [PMID: [24752410](#) DOI: [10.1002/cncr.28726](#)]
 - 8 **Yao JC**, Phan A, Hoff PM, Chen HX, Charnsangavej C, Yeung SC, Hess K, Ng C, Abbruzzese JL, Ajani JA. Targeting vascular endothelial growth factor in advanced carcinoid tumor: a random assignment phase II study of depot octreotide with bevacizumab and pegylated interferon alpha-2b. *J Clin Oncol* 2008; **26**: 1316-1323 [PMID: [18323556](#) DOI: [10.1200/JCO.2007.13.6374](#)]
 - 9 **McMullen T**, Al-Jahdali A, de Gara C, Ghosh S, McEwan A, Schiller D. A population-based study of outcomes in patients with gastrointestinal neuroendocrine tumours. *Can J Surg* 2017; **60**: 192-197 [PMID: [28327275](#) DOI: [10.1503/cjs.007616](#)]
 - 10 **Boyce M**, Thomsen L. Gastric neuroendocrine tumors: prevalence in Europe, USA, and Japan, and rationale for treatment with a gastrin/CCK2 receptor antagonist. *Scand J Gastroenterol* 2015; **50**: 550-559 [PMID: [25665655](#) DOI: [10.3109/00365521.2015.1009941](#)]
 - 11 **Massironi S**, Zilli A, Fanetti I, Ciafardini C, Conte D, Peracchi M. Intermittent treatment of recurrent type-1 gastric carcinoids with somatostatin analogues in patients with chronic autoimmune atrophic gastritis. *Dig Liver Dis* 2015; **47**: 978-983 [PMID: [26321479](#) DOI: [10.1016/j.dld.2015.07.155](#)]
 - 12 **Pavel ME**, Hainsworth JD, Baudin E, Peeters M, Hörsch D, Winkler RE, Klimovsky J, Leblwohl D, Jehl V, Wolin EM, Öberg K, Van Cutsem E, Yao JC; RADIANT-2 Study Group. Everolimus plus octreotide long-acting repeatable for the treatment of advanced neuroendocrine tumours associated with carcinoid syndrome (RADIANT-2): a randomised, placebo-controlled, phase 3 study. *Lancet* 2011; **378**: 2005-2012 [PMID: [22119496](#) DOI: [10.1016/S0140-6736\(11\)61742-X](#)]
 - 13 **Strosberg J**, El-Haddad G, Wolin E, Hendifar A, Yao J, Chasen B, Mittra E, Kunz PL, Kulke MH, Jacene H, Bushnell D, O'Dorisio TM, Baum RP, Kulkarni HR, Caplin M, Lebtahi R, Hobday T, Delpassand E, Van Cutsem E, Benson A, Srirajakanthan R, Pavel M, Mora J, Berlin J, Grande E, Reed N, Seregni E, Öberg K, Lopera Sierra M, Santoro P, Thevenet T, Erion JL, Ruszniewski P, Kwkkeboom D, Krenning E; NETTER-1 Trial Investigators. Phase 3 Trial of ¹⁷⁷Lu-Dotatate for Midgut Neuroendocrine Tumors. *N Engl J Med* 2017; **376**: 125-135 [PMID: [28076709](#) DOI: [10.1056/NEJMoa1607427](#)]
 - 14 **Yao JC**, Fazio N, Singh S, Buzzoni R, Carnaghi C, Wolin E, Tomasek J, Raderer M, Lahner H, Voi M, Pacaud LB, Rouyre N, Sachs C, Valle JW, Fave GD, Van Cutsem E, Tesselaar M, Shimada Y, Oh DY, Strosberg J, Kulke MH, Pavel ME; RAD001 in Advanced Neuroendocrine Tumours, Fourth Trial (RADIANT-4) Study Group. Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study. *Lancet* 2016; **387**: 968-977 [PMID: [26703889](#) DOI: [10.1016/S0140-6736\(15\)00817-X](#)]
 - 15 **Kvols LK**, Oberg KE, O'Dorisio TM, Mohideen P, de Herder WW, Arnold R, Hu K, Zhang Y, Hughes G, Anthony L, Wiedenmann B. Pasireotide (SOM230) shows efficacy and tolerability in the treatment of patients with advanced neuroendocrine tumors refractory or resistant to octreotide LAR: results from a phase II study. *Endocr Relat Cancer* 2012; **19**: 657-666 [PMID: [22807497](#) DOI: [10.1530/ERC-11-0367](#)]
 - 16 **Yao JC**, Phan AT, Chang DZ, Wolff RA, Hess K, Gupta S, Jacobs C, Mares JE, Landgraf AN, Rashid A, Meric-Bernstam F. Efficacy of RAD001 (everolimus) and octreotide LAR in advanced low- to intermediate-grade neuroendocrine tumors: results of a phase II study. *J Clin Oncol* 2008; **26**: 4311-4318 [PMID: [18779618](#) DOI: [10.1200/JCO.2008.16.7858](#)]
 - 17 **Klimstra DS**, Kloppell G, La Rosa S, Rindi G. Classification of neuroendocrine neoplasms of the digestive system. In: WHO Classification of Tumours: Digestive System Tumours, 5th ed, WHO Classification of Tumours Editorial Board (Ed), International Agency for Research on Cancer. Lyon: WHO, 2019: 16
 - 18 **Gluckman CR**, Metz DC. Gastric Neuroendocrine Tumors (Carcinoids). *Curr Gastroenterol Rep* 2019; **21**: 13 [PMID: [30868284](#) DOI: [10.1007/s11894-019-0684-7](#)]
 - 19 **Murugesan SV**, Steele IA, Dimaline R, Poston GJ, Shrotri M, Campbell F, Varro A, Pritchard DM. Correlation between a short-term intravenous octreotide suppression test and response to antrectomy in patients with type-1 gastric neuroendocrine tumours. *Eur J Gastroenterol Hepatol* 2013; **25**: 474-481 [PMID: [23249603](#) DOI: [10.1097/MEG.0b013e32835ccc52](#)]
 - 20 **Gladdy RA**, Strong VE, Coit D, Allen PJ, Gerdes H, Shia J, Klimstra DS, Brennan MF, Tang LH. Defining surgical indications for type I gastric carcinoid tumor. *Ann Surg Oncol* 2009; **16**: 3154-3160 [PMID: [19311111](#) DOI: [10.1245/2009.01.100](#)]

- 19727959 DOI: [10.1245/s10434-009-0687-y](https://doi.org/10.1245/s10434-009-0687-y)]
- 21 **Uygun A**, Kadayifci A, Polat Z, Yilmaz K, Gunal A, Demir H, Bagci S. Long-term results of endoscopic resection for type I gastric neuroendocrine tumors. *J Surg Oncol* 2014; **109**: 71-74 [PMID: [24165913](https://pubmed.ncbi.nlm.nih.gov/24165913/) DOI: [10.1002/jso.23477](https://doi.org/10.1002/jso.23477)]
 - 22 **Manfredi S**, Walter T, Baudin E, Coriat R, Ruzniewski P, Lecomte T, Laurenty AP, Goichot B, Rohmer V, Roquin G, Cojocararu OZ, Lombard-Bohas C, Lepage C, Morcet J, Cadiot G. Management of gastric neuroendocrine tumours in a large French national cohort (GTE). *Endocrine* 2017; **57**: 504-511 [PMID: [28664309](https://pubmed.ncbi.nlm.nih.gov/28664309/) DOI: [10.1007/s12020-017-1355-9](https://doi.org/10.1007/s12020-017-1355-9)]
 - 23 **Pellat A**, Dreyer C, Couffignal C, Walter T, Lombard-Bohas C, Niccoli P, Seitz JF, Hentic O, André T, Coriat R, Faivre S, Zappa M, Ruzniewski P, Pote N, Couvelard A, Raymond E. Clinical and Biomarker Evaluations of Sunitinib in Patients with Grade 3 Digestive Neuroendocrine Neoplasms. *Neuroendocrinology* 2018; **107**: 24-31 [PMID: [29518779](https://pubmed.ncbi.nlm.nih.gov/29518779/) DOI: [10.1159/000487237](https://doi.org/10.1159/000487237)]
 - 24 **Hung OY**, Maithel SK, Willingham FF, Farris AB 3rd, Kauh JS. Hypergastrinemia, type 1 gastric carcinoid tumors: diagnosis and management. *J Clin Oncol* 2011; **29**: e713-e715 [PMID: [21747088](https://pubmed.ncbi.nlm.nih.gov/21747088/) DOI: [10.1200/JCO.2011.35.3235](https://doi.org/10.1200/JCO.2011.35.3235)]
 - 25 **Liu Y**, Uemura N, Xiao SD, Tytgat GN, Kate FJ. Agreement between endoscopic and histological gastric atrophy scores. *J Gastroenterol* 2005; **40**: 123-127 [PMID: [15770394](https://pubmed.ncbi.nlm.nih.gov/15770394/) DOI: [10.1007/s00535-004-1511-x](https://doi.org/10.1007/s00535-004-1511-x)]
 - 26 **Soga J**. Early-stage carcinoids of the gastrointestinal tract: an analysis of 1914 reported cases. *Cancer* 2005; **103**: 1587-1595 [PMID: [15742328](https://pubmed.ncbi.nlm.nih.gov/15742328/) DOI: [10.1002/ncr.20939](https://doi.org/10.1002/ncr.20939)]
 - 27 **Rindi G**, Bordi C, Rappel S, La Rosa S, Stolte M, Solcia E. Gastric carcinoids and neuroendocrine carcinomas: pathogenesis, pathology, and behavior. *World J Surg* 1996; **20**: 168-172 [PMID: [8661813](https://pubmed.ncbi.nlm.nih.gov/8661813/) DOI: [10.1007/s002689900026](https://doi.org/10.1007/s002689900026)]
 - 28 **Delle Fave G**, O'Toole D, Sundin A, Taal B, Ferolla P, Ramage JK, Ferone D, Ito T, Weber W, Zheng-Pei Z, De Herder WW, Pascher A, Ruzniewski P; Vienna Consensus Conference participants. ENETS Consensus Guidelines Update for Gastroduodenal Neuroendocrine Neoplasms. *Neuroendocrinology* 2016; **103**: 119-124 [PMID: [26784901](https://pubmed.ncbi.nlm.nih.gov/26784901/) DOI: [10.1159/000443168](https://doi.org/10.1159/000443168)]
 - 29 **Campana D**, Nori F, Pezzilli R, Piscitelli L, Santini D, Brocchi E, Corinaldesi R, Tomassetti P. Gastric endocrine tumors type I: treatment with long-acting somatostatin analogs. *Endocr Relat Cancer* 2008; **15**: 337-342 [PMID: [18310299](https://pubmed.ncbi.nlm.nih.gov/18310299/) DOI: [10.1677/ERC-07-0251](https://doi.org/10.1677/ERC-07-0251)]
 - 30 **Jianu CS**, Fossmark R, Syversen U, Hauso Ø, Fykse V, Waldum HL. Five-year follow-up of patients treated for 1 year with octreotide long-acting release for enterochromaffin-like cell carcinoids. *Scand J Gastroenterol* 2011; **46**: 456-463 [PMID: [21133821](https://pubmed.ncbi.nlm.nih.gov/21133821/) DOI: [10.3109/00365521.2010.539255](https://doi.org/10.3109/00365521.2010.539255)]
 - 31 **Grozinsky-Glasberg S**, Kaltsas G, Gur C, Gal E, Thomas D, Fichman S, Alexandraki K, Barak D, Glaser B, Shimon I, Gross DJ. Long-acting somatostatin analogues are an effective treatment for type I gastric carcinoid tumours. *Eur J Endocrinol* 2008; **159**: 475-482 [PMID: [18662970](https://pubmed.ncbi.nlm.nih.gov/18662970/) DOI: [10.1530/EJE-08-0420](https://doi.org/10.1530/EJE-08-0420)]
 - 32 **Khuroo MS**, Khuroo MS, Khuroo NS. Treatment of type I gastric neuroendocrine tumors with somatostatin analogs. *J Gastroenterol Hepatol* 2010; **25**: 548-554 [PMID: [20074162](https://pubmed.ncbi.nlm.nih.gov/20074162/) DOI: [10.1111/j.1440-1746.2009.06131.x](https://doi.org/10.1111/j.1440-1746.2009.06131.x)]
 - 33 **Min BH**, Hong M, Lee JH, Rhee PL, Sohn TS, Kim S, Kim KM, Kim JJ. Clinicopathological features and outcome of type 3 gastric neuroendocrine tumours. *Br J Surg* 2018; **105**: 1480-1486 [PMID: [29893418](https://pubmed.ncbi.nlm.nih.gov/29893418/) DOI: [10.1002/bjs.10901](https://doi.org/10.1002/bjs.10901)]
 - 34 **Fykse V**, Sandvik AK, Waldum HL. One-year follow-up study of patients with enterochromaffin-like cell carcinoids after treatment with octreotide long-acting release. *Scand J Gastroenterol* 2005; **40**: 1269-1274 [PMID: [16334435](https://pubmed.ncbi.nlm.nih.gov/16334435/) DOI: [10.1080/00365520510023684](https://doi.org/10.1080/00365520510023684)]
 - 35 **Pritchard DM**. Zollinger-Ellison syndrome: still a diagnostic challenge in the 21st century? *Gastroenterology* 2011; **140**: 1380-1383 [PMID: [21443889](https://pubmed.ncbi.nlm.nih.gov/21443889/) DOI: [10.1053/j.gastro.2011.03.026](https://doi.org/10.1053/j.gastro.2011.03.026)]
 - 36 **Wardlaw R**, Smith JW. Gastric carcinoid tumors. *Ochsner J* 2008; **8**: 191-196 [PMID: [21603501](https://pubmed.ncbi.nlm.nih.gov/21603501/)]
 - 37 **Tan HY**, Lou YN, Luo J, Liu JX, Jia LQ. The typing and treatment of gastric neuroendocrine tumors. *Zhongguo Yixue Qianyan Zazhi* 2014; **6**: 4-8
 - 38 **Shah MH**, Goldner WS, Halfdanarson TR, Bergsland E, Berlin JD, Halperin D, Chan J, Kulke MH, Benson AB, Blaszkowsky LS, Eads J, Engstrom PF, Fanta P, Giordano T, He J, Heslin MJ, Kalemkerian GP, Kandeel F, Khan SA, Kidwai WZ, Kunz PL, Kuvshinov BW, Lieu C, Pillarisetty VG, Saltz L, Sosa JA, Strosberg JR, Sussman CA, Trikalinos NA, Uboha NA, Whisenant J, Wong T, Yao JC, Burns JL, Ogba N, Zuccarino-Catania G. NCCN Guidelines Insights: Neuroendocrine and Adrenal Tumors, Version 2.2018. *J Natl Compr Canc Netw* 2018; **16**: 693-702 [PMID: [29891520](https://pubmed.ncbi.nlm.nih.gov/29891520/) DOI: [10.6004/jnccn.2018.0056](https://doi.org/10.6004/jnccn.2018.0056)]
 - 39 **Bajetta E**, Procopio G, Catena L, Martinetti A, De Dosso S, Ricci S, Lecchi AS, Boscani PF, Iacobelli S, Carteni G, De Braud F, Loli P, Tartaglia A, Bajetta R, Ferrari L. Lanreotide autogel every 6 weeks compared with Lanreotide microparticles every 3 weeks in patients with well differentiated neuroendocrine tumors: a Phase III Study. *Cancer* 2006; **107**: 2474-2481 [PMID: [17054107](https://pubmed.ncbi.nlm.nih.gov/17054107/) DOI: [10.1002/ncr.22272](https://doi.org/10.1002/ncr.22272)]
 - 40 **Kunz PL**. Carcinoid and neuroendocrine tumors: building on success. *J Clin Oncol* 2015; **33**: 1855-1863 [PMID: [25918282](https://pubmed.ncbi.nlm.nih.gov/25918282/) DOI: [10.1200/JCO.2014.60.2532](https://doi.org/10.1200/JCO.2014.60.2532)]
 - 41 **Baldelli R**, Barnabei A, Rizza L, Isidori AM, Rota F, Di Giacinto P, Paoloni A, Torino F, Corsello SM, Lenzi A, Appetecchia M. Somatostatin analogs therapy in gastroenteropancreatic neuroendocrine tumors: current aspects and new perspectives. *Front Endocrinol (Lausanne)* 2014; **5**: 7 [PMID: [24570674](https://pubmed.ncbi.nlm.nih.gov/24570674/) DOI: [10.3389/fendo.2014.00007](https://doi.org/10.3389/fendo.2014.00007)]
 - 42 **Caplin ME**, Pavel M, Ćwikla JB, Phan AT, Raderer M, Sedláčková E, Cadiot G, Wolin EM, Capdevila J, Wall L, Rindi G, Langley A, Martinez S, Blumberg J, Ruzniewski P; CLARINET Investigators. Lanreotide in metastatic enteropancreatic neuroendocrine tumors. *N Engl J Med* 2014; **371**: 224-233 [PMID: [25014687](https://pubmed.ncbi.nlm.nih.gov/25014687/) DOI: [10.1056/NEJMoa1316158](https://doi.org/10.1056/NEJMoa1316158)]
 - 43 **Strosberg JR**, Coppola D, Klimstra DS, Phan AT, Kulke MH, Wiseman GA, Kvols LK; North American Neuroendocrine Tumor Society (NANETS). The NANETS consensus guidelines for the diagnosis and management of poorly differentiated (high-grade) extrapulmonary neuroendocrine carcinomas. *Pancreas* 2010; **39**: 799-800 [PMID: [20664477](https://pubmed.ncbi.nlm.nih.gov/20664477/) DOI: [10.1097/MPA.0b013e3181ebb56f](https://doi.org/10.1097/MPA.0b013e3181ebb56f)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

