

World Journal of *Gastroenterology*

World J Gastroenterol 2020 September 7; 26(33): 4889-5059



FRONTIER

- 4889 Treatment repurposing for inflammatory bowel disease using literature-related discovery and innovation
Kostoff RN, Briggs MB, Shores DR

REVIEW

- 4900 Tumor microenvironment in primary liver tumors: A challenging role of natural killer cells
Polidoro MA, Mikulak J, Cazzetta V, Lleo A, Mavilio D, Torzilli G, Donadon M

MINIREVIEWS

- 4919 Exploring the food-gut axis in immunotherapy response of cancer patients
Russo E, Nannini G, Dinu M, Pagliai G, Sofi F, Amedei A

ORIGINAL ARTICLE**Basic Study**

- 4933 Tumor necrosis factor alpha receptor 1 deficiency in hepatocytes does not protect from non-alcoholic steatohepatitis, but attenuates insulin resistance in mice
Bluemel S, Wang Y, Lee S, Schnabl B
- 4945 Resveratrol alleviates intestinal mucosal barrier dysfunction in dextran sulfate sodium-induced colitis mice by enhancing autophagy
Pan HH, Zhou XX, Ma YY, Pan WS, Zhao F, Yu MS, Liu JQ

Retrospective Study

- 4960 Effects of denosumab treatment in chronic liver disease patients with osteoporosis
Saeki C, Saito M, Oikawa T, Nakano M, Torisu Y, Saruta M, Tsubota A
- 4972 Bowel function and quality of life after minimally invasive colectomy with D3 lymphadenectomy for right-sided colon adenocarcinoma
Lee KM, Baek SJ, Kwak JM, Kim J, Kim SH
- 4983 Acute liver failure and death predictors in patients with dengue-induced severe hepatitis
Teerasartipan T, Chaiteerakij R, Komolmit P, Tangkijvanich P, Treeprasertsuk S
- 4996 Liver fat accumulation measured by high-speed T2-corrected multi-echo magnetic resonance spectroscopy can predict risk of cholelithiasis
Chen H, Zeng WK, Shi GZ, Gao M, Wang MZ, Shen J
- 5008 Radiomics of rectal cancer for predicting distant metastasis and overall survival
Li M, Zhu YZ, Zhang YC, Yue YF, Yu HP, Song B

SYSTEMATIC REVIEWS

- 5022** Neutrophil to lymphocyte ratio and albumin bilirubin grade in hepatocellular carcinoma: A systematic review

Bannaga A, Arasaradnam RP

CASE REPORT

- 5050** Surveilling Russell body *Helicobacter pylori*-negative gastritis: A case report and review of literature

Peruhova M, Peshevska-Sekulovska M, Georgieva V, Panayotova G, Dikov D

ABOUT COVER

Editorial Board of *World Journal of Gastroenterology*, Dr. Dario Sorrentino is a Professor of Medicine at Virginia Tech – Carilion School of Medicine and Research Institute (since 2013). His career research experience has ranged from the bench to the bedside focusing on IBDs, and carried out on three different continents. Fifteen years ago, he and his professional colleagues proposed a groundbreaking strategy to prevent post-surgical recurrence of Crohn's disease that has evolved into today's standard-of-care. More recently, he and his team developed a novel approach for diagnosing and treating pre-clinical Crohn's disease, representing a revolutionary approach to IBD management and research. Dr. Sorrentino has published > 150 high-quality publications and delivered speeches on his own research worldwide. His recent work in the United States has garnered awards of research funds exceeding 2 million dollars from major foundations and private sources. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology (WJG, World J Gastroenterol)* is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJG* mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2020 edition of Journal Citation Report® cites the 2019 impact factor (IF) for *WJG* as 3.665; IF without journal self cites: 3.534; 5-year IF: 4.048; Ranking: 35 among 88 journals in gastroenterology and hepatology; and Quartile category: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yu-Jie Ma*; Production Department Director: *Xiang Li*; Editorial Office Director: *Ze-Mao Gong*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, Subrata Ghosh

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

September 7, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Basic Study

Resveratrol alleviates intestinal mucosal barrier dysfunction in dextran sulfate sodium-induced colitis mice by enhancing autophagy

Hang-Hai Pan, Xin-Xin Zhou, Ying-Yu Ma, Wen-Sheng Pan, Fei Zhao, Mo-Sang Yu, Jing-Quan Liu

ORCID number: Hang-Hai Pan 0000-0002-7366-8269; Xin-Xin Zhou 0000-0003-0183-6400; Ying-Yu Ma 0000-0003-2364-7846; Wen-Sheng Pan 0000-0002-2347-1695; Fei Zhao 0000-0002-5217-4163; Mo-Sang Yu 0000-0002-5614-0227; Jing-Quan Liu 0000-0003-3588-6457.

Author contributions: Pan HH designed the experiments; Ma YY, Yu MS and Zhao F performed the experiments and analyzed the data; Pan HH and Zhou XX drafted the manuscript; Ma YY critically revised the manuscript; Liu JQ and Pan WS offered help during the experiments; all authors have read and approved the final manuscript.

Supported by the National Natural Science Foundation of China, No. 81600414; Medical Health Science and Technology Project of Zhejiang Provincial Health Commission, No. 2018255969; Zhejiang TCM Science and Technology Project, No. 2016ZA123 and No. 2018ZA013.

Institutional review board

statement: This study was reviewed and approved by the Institutional Review Board of Zhejiang Chinese Medical University.

Institutional animal care and use

Hang-Hai Pan, Wen-Sheng Pan, Fei Zhao, Department of Gastroenterology, Zhejiang Provincial People's Hospital, People's Hospital of Hangzhou Medical College, Hangzhou 310014, Zhejiang Province, China

Xin-Xin Zhou, Mo-Sang Yu, Department of Gastroenterology, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, Zhejiang Province, China

Ying-Yu Ma, Key Laboratory of Gastroenterology of Zhejiang Province, Zhejiang Provincial People's Hospital, People's Hospital of Hangzhou Medical College, Hangzhou 310014, Zhejiang Province, China

Jing-Quan Liu, Critical Care Unit, Zhejiang Provincial People's Hospital, People's Hospital of Hangzhou Medical College, Hangzhou 310014, Zhejiang Province, China

Corresponding author: Ying-Yu Ma, MD, Research Assistant, Key Laboratory of Gastroenterology of Zhejiang Province, Zhejiang Provincial People's Hospital, People's Hospital of Hangzhou Medical College, No. 158 Shangtang Road, Hangzhou 310014, Zhejiang Province, China. myy011525@163.com

Abstract**BACKGROUND**

Intestinal mucosal barrier dysfunction plays an important role in the pathogenesis of ulcerative colitis (UC). Recent studies have revealed that impaired autophagy is associated with intestinal mucosal dysfunction in the mucosa of colitis mice. Resveratrol exerts anti-inflammatory functions by regulating autophagy.

AIM

To investigate the effect and mechanism of resveratrol on protecting the integrity of the intestinal mucosal barrier and anti-inflammation in dextran sulfate sodium (DSS)-induced ulcerative colitis mice.

METHODS

Male C57BL/6 mice were divided into four groups: negative control group, DSS model group, DSS + resveratrol group, and DSS + 5-aminosalicylic acid group. The severity of colitis was assessed by the disease activity index, serum inflammatory cytokines were detected by enzyme-linked immunosorbent assay.

committee statement: All procedures involving animals were reviewed and approved by the Ethics Committee of Zhejiang Chinese Medical University.

Conflict-of-interest statement: No potential conflicts of interest exist.

Data sharing statement: No additional data are available.

ARRIVE guidelines statement: The authors have read the ARRIVE guidelines, and the manuscript was prepared and revised according to the ARRIVE guidelines.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: May 17, 2020

Peer-review started: May 17, 2020

First decision: June 18, 2020

Revised: June 27, 2020

Accepted: August 12, 2020

Article in press: August 12, 2020

Published online: September 7, 2020

P-Reviewer: Gregorio BM, Hu J, Katada, K, Li CY

S-Editor: Zhang L

L-Editor: MedE-Ma JY

P-Editor: Ma YJ



Colon tissues were stained with haematoxylin and eosin, and mucosal damage was evaluated by mean histological score. The expression of occludin and ZO-1 in colon tissue was evaluated using immunohistochemical analysis. In addition, the expression of autophagy-related genes was determined using reverse transcription-polymerase chain reaction and Western-blot, and morphology of autophagy was observed by transmission electron microscopy.

RESULTS

The resveratrol treatment group showed a 1.72-fold decrease in disease activity index scores and 1.42, 3.81, and 1.65-fold decrease in the production of the inflammatory cytokine tumor necrosis factor- α , interleukin-6 and interleukin-1 β , respectively, in DSS-induced colitis mice compared with DSS group ($P < 0.05$). The expressions of the tight junction proteins occludin and ZO-1 in DSS model group were decreased, and were increased in resveratrol-treated colitis group. Resveratrol also increased the levels of LC3B (by 1.39-fold compared with DSS group) and Beclin-1 (by 1.49-fold compared with DSS group) ($P < 0.05$), as well as the number of autophagosomes, which implies that the resveratrol may alleviate intestinal mucosal barrier dysfunction in DSS-induced UC mice by enhancing autophagy.

CONCLUSION

Resveratrol treatment decreased the expression of inflammatory factors, increased the expression of tight junction proteins and alleviated UC intestinal mucosal barrier dysfunction; this effect may be achieved by enhancing autophagy in intestinal epithelial cells.

Key words: Resveratrol; Ulcerative colitis; Autophagy; Intestinal mucosal barrier; Dextran sulfate sodium-induced colitis; Intestinal inflammation

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: We established a chronic colitis model successfully *via* administration of dextran sulfate sodium (DSS), and we found that resveratrol ameliorates the production of the inflammatory cytokines in DSS-induced colitis mice. Meanwhile, resveratrol treatment alleviated intestinal mucosal barrier dysfunction in DSS-induced colitis and increased the expression of the tight junction proteins occludin and ZO-1. Further studies showed that resveratrol treatment increased the levels of LC3B and Beclin-1 in the colons of colitis mice, as well as the number of autophagosomes, which may *via* enhancing autophagy.

Citation: Pan HH, Zhou XX, Ma YY, Pan WS, Zhao F, Yu MS, Liu JQ. Resveratrol alleviates intestinal mucosal barrier dysfunction in dextran sulfate sodium-induced colitis mice by enhancing autophagy. *World J Gastroenterol* 2020; 26(33): 4945-4959

URL: <https://www.wjgnet.com/1007-9327/full/v26/i33/4945.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v26.i33.4945>

INTRODUCTION

Inflammatory bowel disease (IBD) is a chronic intestinal inflammatory condition involving the gastrointestinal tract, comprising ulcerative colitis (UC) and Crohn's disease. UC is characterized by abdominal pain, diarrhoea, and haematochezia. The incidence of UC is rising with changes in dietary habits and the rhythm of life^[1,2]. UC is often persistent or recurrent in patients and increases the risk of colorectal cancer^[3]. Currently, drugs such as 5-aminosalicylic acid (5-ASA), corticosteroids and immunosuppressants are used, but the effect of these drugs is poor, and the long-term use of these drugs may cause severe adverse reactions, resulting in great psychological and economic burdens on the patients^[4,5]. Therefore, effective and safe drugs against UC are urgently needed.

The initiating event and pathogenesis of UC currently remain elusive. Recent work has highlighted the importance of intestinal mucosal barrier dysfunction in disease pathophysiology^[6]. Many studies have shown that the destruction of the intestinal

mucosal barrier plays an important role in the deterioration of IBD, especially in UC patients^[7-9]. Protecting the integrity of the intestinal mucosal barrier is thought to be a potential clinical treatment approach for UC.

Resveratrol (3,4',5-Trihydroxy-trans-stilbene), a natural plant polyphenol found in grapes, is the principal biologically active component in red wine. Resveratrol has anti-inflammatory and immunoregulatory functions, with the advantages of low price and few side effects^[10-12]. So far, very few clinical studies have indicated that resveratrol can alleviate clinical colitis activity and improve quality of life and in patients with active UC^[13,14]. Although these limited studies show that resveratrol may ameliorate inflammation in UC, the mechanism remains unclear. Intestinal epithelial cells (IECs) are reported to act as the first line of defence in the intestinal mucosal barrier. The IECs tight junctions may play an important role in intestinal mucosal barrier. Occludin and ZO-1 are important tight junction proteins that play a significant role in maintaining the integrity of the intestinal mucosal barrier. Thus, we focused on whether resveratrol can alleviate intestinal mucosal barrier injury and inflammation. In addition, the mechanism underlying the anti-inflammatory effect of resveratrol in colitis was explored.

Autophagy is a cellular recycling process involving self-degradation and the reconstruction of damaged organelles and proteins^[15]. Recent studies have shown that impaired autophagy is related to intestinal mucosal damage in mice with colitis, and stimulation of autophagy can prevent intestinal mucosal inflammation and ameliorates murine colitis in mice^[16]. Furthermore, the regulatory effect of resveratrol on autophagy in inflammatory diseases has gradually attracted attention^[17,18]. In the present study, we evaluated the effect of resveratrol on dextran sulfate sodium (DSS)-induced colitis in mice and explore the mechanism of resveratrol on protecting the integrity of the intestinal mucosal barrier and anti-inflammation.

MATERIALS AND METHODS

Chemicals and reagents

Resveratrol (3,4',5-Trihydroxy-trans-stilbene; C₁₄H₁₂O₃; Molecular Weight: 228.24 g/mol; **Figure 1A**) and 5-Aminosalicylic acid (5-ASA; C₇H₇NO₃; Molecular Weight: 153.14 g/mol; **Figure 1B**) were purchased from Sigma Chemical (United States). The purity was checked *via* HPLC analysis and exceeded 99%. Dextran Sulfate Sodium (DSS, Molecular Weight: 36000-50000) was purchased from MP Biomedicals (United States).

Animals

Male C57BL/6 mice aged 5 wk and weighing 17-19 g were purchased from Shanghai SLAC Laboratory Animal Co., Ltd. Animals were housed in mouse cages at a controlled temperature (21 ± 2 °C) with a pre-set 12-h light/dark cycle. The mice were fed a standard laboratory diet and normal drinking water. The normal diet included crude proteins ≥ 20%, crude fat ≥ 6%, crude fibre ≤ 5%, crude ash ≤ 8%, lysine 1.32%, calcium 1%-1.8%, and phosphorus 0.6%-1.2%, respectively. The animals were acclimated to the experimental conditions for one week before the experiment. According to the guidelines for the care and use of laboratory animals, all the procedures for the treatment and execution of mice were approved by the Animal Care and Use Committee of Zhejiang Chinese Medical University.

Experimental design

DSS-induced colitis is a widely used model that morphologically and symptomatically resembles UC in humans. In this study, we established DSS-induced chronic colitis model to study UC. Mice were randomly divided into four experimental groups (*n* = 12 mice/group): Negative control group, DSS model group, resveratrol-treated group and 5-ASA positive-control group. Mice in the negative control group received normal drinking water for 28 d. In the DSS model group, chronic colitis was induced by two cycles of giving drinking water containing 3% DSS for 7 d and normal drinking water for 7 d. Resveratrol treatment group (DSS + RES) mice received two cycles of DSS by drinking water, and treatment of resveratrol 100 mg/kg per day by gavage for two cycles followed by DSS at the same time (**Figure 1C**). 5-ASA, the first-line therapy for UC, was used as a positive control to evaluate the efficacy of resveratrol. Mice in the positive-control group (DSS + 5-ASA) were treated with two cycles of DSS, and 5-ASA 200 mg/kg per day for 28 d at the same time. Body mass was detected every 5 d, and body mass loss was calculated as (detected body mass-initial mass) / initial mass. All




Figure 1 Chemical structures of resveratrol and 5-aminosalicylic acid, and timeline of *in vivo* experiment. A: The chemical structures of resveratrol; B: The chemical structures of 5-aminosalicylic acid; and C: Timeline of *in vivo* experiment. Dextran sulfate sodium (DSS) group was induced by two cycles of intake of drinking water containing 3% DSS for 7 d and normal drinking water for 7 d. Other groups were also treated by two cycles, first received DSS by drinking water for 7 d, followed by treatment of 5-aminosalicylic acid or resveratrol by gavage for 7 d. DSS: Dextran sulfate sodium; 5-ASA: 5-aminosalicylic acid; RES: Resveratrol.

animals were euthanized with carbon dioxide and sacrificed at humane end points.

Clinical evaluation of colitis

To assess the severity of colitis, the disease activity index (DAI) was calculated daily based on weight loss, stool consistency and rectal bleeding (Table 1)^[19]. Stool consistency (0, normal; 2, loose stool; 4, watery diarrhea); bloody stools (0, normal; 2, slight bleeding; 4, gross bleeding); and body weight loss (0, < 2%; 1, decreased ≥ 2 - < 5%; 2, decreased ≥ 5 - < 10%; 3, decreased ≥ 10 - < 15%; 4, decreased ≥ 15). These values were assessed for each animal, and the sum of the 3 values constituted the DAI. For each parameter, the scores ranged from 0 to 4, resulting in a total DAI score ranging from 0 (unaffected) to 12 (severe colitis).

Inflammatory cytokine assay

Mice were sacrificed, and blood samples (1.0 mL) were collected from the heart. The expression levels of tumour necrosis factor- α (TNF- α , MTA00B, R&D Systems Europe Ltd., Abingdon, England), interleukin (IL)-6 (M6000B, R&D Systems Europe Ltd., Abingdon, England), and IL-1 β (MLB00C, RD Systems Europe Ltd., Abingdon, England) in the plasma were detected using the enzyme-linked immunosorbent assay kits according to the manufacturer's instructions. The concentration of the cytokines was expressed as pg/mL per mg protein.

Histopathological analysis

Colon length can be used as an indirect marker of inflammation. Thus, when mice were sacrificed, the length of colons from the caecum to the anus was measured. Then, the colon was fixed in 10% formalin, dehydrated at gradient concentrations of ethanol, and embedded in paraffin. Tissue sections (8 μ m) were prepared and stained with haematoxylin and eosin (HE), and mucosal damage was evaluated. To evaluate the colonic mucosal damage severity, 9 fields were randomly selected and observed under an inverted fluorescence microscope by two pathologists who were blinded to the experimental groups. Colonic mucosal damage was confirmed by the mean histological score (Table 2)^[20]. Each histological score, such as inflammation (0-3), extent (0-3), regeneration (0-4) and crypt damage (0-4), was multiplied by the percentage of compromised tissue (1 point for 25%, 2 points for 26%-50%, 3 points for 51%-75%, and 4 points for 76%-100%). Therefore, the scores of inflammation and extent range from 0 to 12, and the scores of regeneration and crypt damage range from 0 to 16, which reached a maximum score of 56.

Table 1 Assessment of the disease activity index

Body mass loss (%)	Stool consistency	Bleeding	Score
< 2%	Normal	No bleeding	0
≥ 2 - < 5%	-	-	1
≥ 5 - < 10%	Loose stools	Slight bleeding	2
≥ 10 - < 15%	-	-	3
≥ 15	Watery diarrhea	Gross bleeding	4

Table 2 Histological scoring of colitis

Features	Grade	Description
Inflammation	0	None
	1	Slight
	2	Moderate
	3	Severe
Extent	0	None
	1	Mucosa
	2	Mucosa and submucosa
	3	Transmural
Regeneration	4	No tissue repair
	3	Surface epithelium not intact
	2	Regeneration with crypt depletion
	1	Almost complete regeneration
	0	Complete regeneration or normal tissue
Crypt damage	0	None
	1	Basal 1/3 damaged
	2	Basal 2/3 damaged
	3	Only surface epithelium intact
	4	Entire crypt and epithelium lost
Percent involvement	1	1%-25%
	2	26%-50%
	3	51%-75%
	4	76%-100%

Immunohistochemical analysis

The expression of occludin and ZO-1 was detected by immunohistochemical analysis. Briefly, all blocks of colonic tissue were sectioned (3-5 μ m). Then, sections were deparaffinized, rehydrated at graded ethanol concentrations and incubated with fresh 3% hydrogen peroxide for 10 min. After being rinsed with phosphate buffer saline (PBS), each tissue section was subjected to antigen retrieval by a suitable antigen retrieval method (boiling in 0.01 M citrate buffer). After endogenous peroxidases were blocked, sections were incubated with anti-occludin antibody (ab222691, 1:200 dilution in PBS, Abcam, United States) and a monoclonal ZO-1 antibody (ab221546, 1:300, Abcam, United States) overnight at 4°C. Subsequently, the sections were incubated for 20 min at room temperature with biotin-labelled secondary antibodies, stained with 3,3'-diaminobenzidine, counterstained with haematoxylin, dehydrated and mounted. Primary antibody was replaced by PBS as the negative control. The following criterion was used for semi-quantitative staining: 0 (no staining), 1 (weak staining, light yellow),

2 (moderate staining, yellow brown), and 3 (strong staining, brown). The proportion of stained tumor cells was scored according to the proportion of positively stained tumor cells as follows: 0 (< 5%); 1 (6-25%); 2 (26-50%); and 3 (> 51%). A composite score was obtained by the staining intensity score multiplied the proportion scores. The total score ≤ 4 was defined as low expression, and > 5 was regarded as high expression.

Western blotting analysis

Briefly, proteins were extracted using radioimmunoprecipitation assay buffer, separated by sodium dodecyl sulphate-polyacrylamide gel (5%-10%) electrophoresis, and then transferred to polyvinylidene difluoride membranes (Millipore, Billerica, MA, United States). These membranes were washed with PBS and blocked with Tris-buffered saline-Tween with 5% skim milk (232100, BD, Bioscience, United States). Primary β -actin (3700S, 1:1000, Cell Signal Technology, United States), LC3B (3868T, 1:1000, Cell Signal Technology, United States) and Beclin-1 (3495T, 1:1000, Cell Signal Technology, United States) antibodies were applied, followed by horseradish peroxidase-conjugated secondary antibodies. Then, an enhanced chemiluminescence detection system (Thermo Fisher Scientific, United States) was used to detect antibody binding. Meanwhile, β -actin was used as a negative control.

Reverse transcription-polymerase chain reaction analysis

Total ribonucleic acid was extracted from tissue specimens using TRIzol reagent, and messenger ribonucleic acid (mRNA) was reverse transcribed to complementary deoxyribonucleic acid with a QuantiTect reverse transcription kit (205311, Qiagen, German) according to the manufacturer's instructions. Quantitative polymerase chain reaction (qPCR) was performed using a SuperScript III Platinum SYBR green one-step qPCR kit (11732088, Thermo Fisher). The qPCR primer sequences were as follows: actin: 5'-CAT CCG TAA AGA CCT CTA TGC CAA C-3' (forward) and 5'-ATG GAG CCA CCG ATC CAC A-3' (reverse); Beclin: 5'-ATG GAG GGG TCT AAG GCG TC-3' (forward) and 5'-TGG GCT GTG GTA AGT AAT GGA-3' (reverse); LC3B: 5'-TCA AGT CCA ACT ACC GAG TCC-3' (forward) and 5'-TCA GAG GTT TCC CAT CCA AG-3' (reverse). Relative messenger ribonucleic acid expression levels were calculated based on the computerized tomography values and normalized using actin expression. All experiments were performed in triplicate on a Roche LightCycler 480 platform.

Transmission electron microscopy

First, colon tissues were fixed with glutaraldehyde and osmium tetroxide, then dehydrated in ethanol, passed through propylene oxide, and embedded in Spurr resin. Fifty nanometres thick sections were cut by an ultramicrotome, and the sections were subsequently post-stained with 4% uranyl acetate for 10 min and Reynold's lead citrate for 1.5 min. The structure of the IECs and autolysosomes were observed by transmission electron microscopy (HITACHI H-7650).

Statistical analysis

All statistical analyses were performed using Statistical Package for the Social Sciences software (version 13.0; Statistic Package for Social Science Inc., Chicago, IL, United States). Continuous variables were presented as the mean and standard deviation and analysed using one-way Anova or Kruskal-Wallis test. All values were expressed as the mean \pm standard deviation. All statistical analyses were conducted with $\alpha = 0.05$, and $P < 0.05$ was considered significant.

RESULTS

Clinical activity of DSS-induced colitis mice

DSS-induced colitis was evaluated, and the clinical course was monitored and scored for the presence of bloody stool, watery diarrhoea and weight loss. During this experiment, three mice in the DSS group died, and one of the mice died on the 13th day, and the other two died on the 14th day; one mouse in the DSS + 5-ASA group died on the 13th day, and no mice in the DSS + RES group died. In comparison with those in the negative control group, mice in the DSS group showed progressively increased DAI score. However, the DSS + RES group showed 1.72-fold decrease in DAI score compared with the DSS group ($P < 0.05$, Figure 2A), which indicated that resveratrol may have a favourable effect on colitis.




Figure 2 Clinical evaluation of dextran sulfate sodium-induced chronic colitis by disease activity index, body mass loss and colon length ($n = 12$ for the control group, $n = 9$ for disease activity index group, $n = 11$ for dextran sulfate sodium + 5-aminosalicylic acid group, $n = 12$ for dextran sulfate sodium + resveratrol treatment group). A: The disease activity index score was reduced in the resveratrol treated group compared with the dextran sulfate sodium group ($^{\#}P < 0.05$); B and C: Resveratrol treatment increased the body mass and colon length of dextran sulfate sodium-induced colitis mice. Body mass loss was calculated as (detected body mass - initial mass)/initial mass. DSS: Dextran sulfate sodium; 5-ASA: 5-aminosalicylic acid; RES: Resveratrol.

Effects of resveratrol on body mass and colon length in DSS-induced colitis mice

Body mass could be decreased by DSS-induced inflammation; therefore, we studied the effect of resveratrol on body mass in DSS-induced colitis mice. The body mass was measured every 5 d, and the results showed that DSS-treated mice exhibited a statistically significant decrease. Resveratrol and 5-ASA treatment increased the body mass of DSS-induced colitis mice, but the difference was not significant (Figure 2B). We also measured the colon length of DSS-induced colitis mice and found that the colon length of the resveratrol-treated group was longer than that of the DSS group, but the difference was also not significant (Figure 2C).

Resveratrol ameliorates the production of inflammatory cytokines in DSS-induced colitis mice

The enzyme-linked immunosorbent assay results demonstrated that the protein expression levels of the inflammatory cytokines TNF- α , IL-6, and IL-1 β were higher in the DSS-induced colitis group than in the control group ($P < 0.05$, Figure 3). Furthermore, the levels of TNF- α and IL-6 in the 5-ASA-treatment group were lower than those in the DSS-induced colitis group ($P < 0.05$). However, the levels of TNF- α , IL-6 and IL-1 β showed 1.42, 3.81, and 1.65-fold decrease in the resveratrol group compared with the DSS group ($P < 0.05$), and level of IL-1 β also showed a 1.57-fold decrease in the resveratrol group compared with that in the DSS + 5-ASA group ($P < 0.05$).




Figure 3 Inflammatory cytokine expression in the dextran sulfate sodium-induced chronic colitis determined by enzyme-linked immunosorbent assay. The levels of interleukin-1β, interleukin-6 and tumor necrosis factor-α were decreased in the resveratrol treatment group compared with the dextran sulfate sodium group. ^a*P* < 0.05. TNF: Tumor necrosis factor; DSS: Dextran sulfate sodium; 5-ASA: 5-aminosalicylic acid; RES: Resveratrol.

Resveratrol alleviates intestinal mucosal barrier injury in DSS-induced colitis mice as determined by histopathology

We also assessed colon damage by histologic examination. DSS group showed significant colonic mucosal damages, crypt depletion, infiltration of inflammatory cells into the mucosa and submucosa, loss of epithelial barrier. The colon damage and intestinal inflammation in mice treated with resveratrol and 5-ASA were less severe than that in DSS-induced colitis mice (Figure 4A). Consistent with the histologic damage, the histological scores of colitis mice treated with resveratrol were significantly lower than those of DSS-induced colitis mice (*P* < 0.05, Figure 4B), while the histological scores of 5-ASA-treated colitis mice were not significantly different from those of mice in the DSS group. These results indicate that resveratrol may alleviate intestinal mucosal barrier dysfunction in DSS-induced colitis mice more effectively than 5-ASA.

Resveratrol increases the production of ZO-1 and occludin in the colons of DSS-treated mice

The tight junction proteins occludin and ZO-1 have been reported to play an important role in maintaining the integrity of the intestinal mucosal epithelial barrier. In the present study, we found that the expression levels of occludin and ZO-1 were higher in the resveratrol-treated DSS group than in the 5-ASA and DSS groups (Figure 5 and Table 3).

Table 3 ZO-1 and occludin expressions in colitis

Group	ZO-1 expression		χ^2/P value	Occludin expression		χ^2/P value
	Low (%)	High (%)		Low (%)	High (%)	
Control	6 (50.0%)	6 (50.0%)	8.812/0.029	7 (58.3%)	5 (41.7%)	10.987/0.010
DSS	7 (77.8%)	2 (22.2%)		8 (88.9%)	1 (11.1%)	
DSS + 5-ASA	7 (63.6%)	4 (36.4%)		9 (81.8%)	2 (18.2%)	
DSS + RES	2 (16.7%)	10 (83.3%)		3 (25.0%)	9 (75.0%)	

DSS: Dextran sulfate sodium; 5-ASA: 5-aminosalicylic acid; RES: Resveratrol.

Resveratrol may reduce inflammation by the enhancement of autophagy in DSS-induced colitis

To explore whether the inhibition of colitis by resveratrol treatment is mediated by autophagy activation, we examined the mRNA and protein levels of LC3B and Beclin-1. As determined by real-time PCR, a substantial increase in the mRNA expression level of LC3B and Beclin-1 was observed in the DSS + RES group compared with the DSS group, suggesting that resveratrol increases the mRNA levels of LC3B and Beclin-1 in the colons of DSS-treated mice ($P < 0.05$, Figure 6A). Consistent with the mRNA level analysis, the western blot results also showed that resveratrol treatment induced significant increases in the LC3-II/I ratio and Beclin-1 level in DSS-induced colitis mice ($P < 0.05$, Figure 6B and C). However, the difference in the mRNA levels of LC3B and Beclin-1 was not significant between the 5-ASA-treated colitis group and the DSS group ($P > 0.05$).

The transmission electron microscopy results showed that IECs in the control group had normal organelle structure with a normal endoplasmic reticulum and mitochondria (Figure 7A). However, in the DSS-induced colitis group, the organelle structure was markedly damaged, with swollen mitochondria, disrupted mitochondrial inner ridges and dilated rough endoplasmic reticulum (Figure 7B). Inflammation induced by the 5-ASA treatment and autophagosomes were rare (Figure 7C). Additionally, resveratrol administration increased the number of autophagosomes and improved the condition of the endoplasmic reticulum and mitochondria, which indicated that resveratrol promoted the progression of autophagy (Figure 7D).

DISCUSSION

UC is a recurrent chronic inflammation of the colon. DSS-induced colitis models, including acute and chronic colitis models, are most commonly used in UC. Acute colitis model is usually induced by administering DSS for 7 d. In this study, a chronic colitis model was successfully established by the administration of two cycles of DSS treatment. Each cycle involved the administration of 3% DSS for 7 d followed by normal drinking water for 7 d^[21]. Mice in the chronic colitis group developed clinical symptoms, such as bloody stool and weight loss; the mice also showed pathological features similar to UC patients, such as shortened colon length, intestinal mucosa congestion, oedema and erosion, ulceration, crypt damage, and the infiltration of inflammatory cells into the mucosa and submucosa; the histological scores of colitis and serum inflammation indices (TNF- α , IL-6, IL-1 β) in the DSS group increased. The chronic colitis model induced by administering repeated cycles of DSS more closely resembled human UC than the acute colitis model did.

In recent years, the role of intestinal mucosal barrier dysfunction in the pathogenesis of UC has attracted wide attention. IECs are the first-line defence in the intestinal mucosal barrier^[22]. The IECs tight junctions are key epithelial intracellular junctions^[23,24]. Occludin and ZO-1 are important tight junction proteins that play a significant role in maintaining the integrity of the intestinal mucosal barrier. The integrity of the intestinal mucosal barrier prevents and defends against the invasion of inflammatory factors and bacteria and further mitigates intestinal inflammation. Resveratrol is a well-known phytochemical with pleiotropic properties and has anti-oxidative and anti-inflammatory activity. Recently, resveratrol has also been shown to




Figure 4 Histological staining showed colitis induced dysfunction. A: Dextran sulfate sodium group showed significant colonic mucosal damages, crypt depletion, infiltration of inflammatory cells into the mucosa and submucosa, loss of epithelial barrier. Resveratrol treatment group and 5-aminosalicylic acid could alleviate colitis-induced intestinal mucosal barrier dysfunction; and B: The histological score of resveratrol treatment group was lower than the dextran sulfate sodium group. ^a*P* < 0.05. DSS: Dextran sulfate sodium; 5-ASA: 5-aminosalicylic acid; RES: Resveratrol.

alleviate intestinal inflammation^[25-27]. Consistent with these studies, our results also showed that resveratrol can reduce intestinal inflammation and alleviate intestinal mucosal barrier dysfunction in DSS-induced colitis mice. Resveratrol significantly increased levels of tight junction proteins (occludin and ZO-1) and reduced histological colonic mucosa injury scores, even than 5-ASA did. Resveratrol also decreased serum inflammation indices (TNF- α , IL-6, and IL-1 β), although there was no significant difference between resveratrol and 5-ASA treatment. These results showed that both resveratrol and 5-ASA alleviated intestinal inflammation, and resveratrol performed better at maintaining the integrity of the intestinal mucosal epithelial barrier than 5-ASA.

Autophagy is a cellular self-protective mechanism stimulated by both internal and external adverse environmental factors. In this process, degenerated proteins, aged and damaged organelles, even pathogenic microorganisms, are recycled *via* autophagic degradation to maintain cellular metabolic balance and homeostasis^[15,28,29]. Autophagy can be activated when an organ is stimulated by adverse factors. Saito M




Figure 5 Immunohistochemical staining of ZO-1 and occludin. Expressions of ZO-1 and occludin were higher in resveratrol treatment than in the dextran sulfate sodium group and 5-aminosalicylic acid treated group. Negative control: antibody was replaced by phosphate buffer saline. DSS: Dextran sulfate sodium; 5-ASA: 5-aminosalicylic acid; RES: Resveratrol.

and others found a reduced LC3-II/I ratio, expression of adhesion proteins and IECs adhesion, and a damaged mucosal epithelial barrier when IECs were exposed to TNF- α and treated with autophagy inhibitors^[30]. In this study, we found that resveratrol alleviated the intestinal mucosal barrier dysfunction and reduced the intestinal inflammatory response by enhancing autophagy in experimental chronic colitis (two cycles of DSS treatment for 28 d), as the expression of LC3B and Beclin-1, and the number of autophagosomes in resveratrol-treated group were increased when compared to that in the chronic colitis model group and the negative control group.

A previous study conducted by Zhang *et al.*^[31] reported that DSS-inducing increased the autophagy in acute colitis models (DSS treatment for 7 d), and curcumin and resveratrol could protect against colitis by reducing autophagy rather than enhancing autophagy^[31]. We speculated that this inconsistency may be caused by the different colitis models used in the two studies. The acute colitis model resembles the early stage of human UC, while the chronic colitis model resembles the advanced stage of UC. Our previous study showed that autophagy was dynamic in the progression of colitis. It was activated in IECs by inflammatory factors in the early stage of a rat model of sepsis-induced acute colitis, and decreased as the disease progressed, thereby reducing the removal of oxidative stress products and the damaged organelles and other substances in cells, which lead to the destruction of the intestinal mucosal barrier and the aggravation of intestinal inflammation^[32]. We believe that the chronic colitis mouse model is a better tool to explore the pathogenesis and drug effects of ulcerative colitis.

However, there are still many limitations in the present study. The detailed autophagy involved in resveratrol-induced protection of intestinal mucosal barrier was unclear as we did not study the effect of autophagy inhibitors on colitis treated by resveratrol. On the other hand, we only investigated the effect of resveratrol on animals, therefore, further research is still needed for clinical application.




Figure 6 Expression levels of Beclin-1 and LC3B in dextran sulfate sodium induced chronic colitis. A: A substantial increase in the messenger ribonucleic acid expression level of LC3B and Beclin-1 was observed in the dextran sulfate sodium (DSS) + resveratrol treatment group compared with the DSS group ($P < 0.05$); B and C: The Western blotting showed that resveratrol treatment induced significant increases in the LC3-II/I ratio and Beclin-1 level in DSS-induced colitis mice; Mean grey level of Beclin-1 and LC3B protein level was increased in DSS + resveratrol treatment group. ^a $P < 0.05$. DSS: Dextran sulfate sodium; 5-ASA: 5-aminosalicylic acid; mRNA: Messenger ribonucleic acid; RES: Resveratrol.




Figure 7 Structures of the intestinal epithelial cells and autolysosomes observed by transmission electron microscopy. A: Control group showed normal organelle structure; B: Dextran sulfate sodium induced colitis group showed mitochondrial swelling and ruptures of internal cristae, along with distention of the rough endoplasmic reticulum; C: 5-aminosalicylic acid group revealed that mitochondrial swelling was not obvious, and mitochondrial cristae showed blurred appearance; and D: Resveratrol treatment group showed that organelle structure was basically normal, and autophagosome could be observed. ER: Endoplasmic reticulum, M: Mitochondria.

ARTICLE HIGHLIGHTS

Research background

Intestinal mucosal barrier disorder plays a very important role in the pathogenesis of ulcerative colitis (UC). Recent studies have revealed that impaired autophagy is associated with intestinal mucosal dysfunction in the mucosa of colitis mice.

Research motivation

Resveratrol can regulate autophagy in the treatment of a few inflammatory diseases. Recently, few studies have indicated that resveratrol can alleviate clinical colitis activity in patients with active UC, while the mechanism for its anti-inflammatory effect remains elusive.

Research objectives

The aim of the study was to explore the effect and mechanism of resveratrol on protecting the integrity of the intestinal mucosal barrier and anti-inflammation in dextran sulfate sodium (DSS)-induced ulcerative colitis.

Research methods

DSS-induced ulcerative colitis was induced by DSS, then the disease activity index was used to assess the severity of colitis. Inflammatory cytokines were detected by enzyme-linked immunosorbent assay. Tissue sections were stained with haematoxylin and eosin, and mucosal damage was evaluated by mean histological score. The expression of occludin and ZO-1 was detected by immunohistochemical analysis. Reverse transcription-polymerase chain reaction and Western-blot were used to analyze autophagy-related gene expression, and morphology of autophagy was observed by transmission electron microscopy.

Research results

The resveratrol treatment group showed a 1.72-fold decrease in disease activity index scores and 1.42-, 3.81-, and 1.65-fold decrease in the production of the inflammatory cytokines tumor necrosis factor- α , interleukin-6 and interleukin-1 β , respectively, in DSS-induced colitis mice compared with DSS group ($P < 0.05$). The expression of the tight junction proteins occludin and ZO-1 in DSS model group was reduced, while in resveratrol-treated colitis group was increased. Resveratrol also increased the levels of

LC3B (1.39-fold compared with DSS group) and Beclin-1 (1.49-fold compared with DSS group) ($P < 0.05$), as well as the number of autophagosomes, which implies that the resveratrol may alleviate intestinal mucosal barrier dysfunction in DSS-induced UC mice by enhancing autophagy.

Research conclusions

Resveratrol treatment reduces the expression of inflammatory factors, increases the expression of tight junction proteins and alleviates UC intestinal mucosal barrier dysfunction; this effect is achieved *via* the regulation of autophagy in intestinal epithelial cells.

Research perspectives

This work suggests that resveratrol may be useful as a new approach to treat UC by enhancing autophagy. Further study of the functionary mechanism will help us to understand the role of resveratrol in colitis and provide a theoretical basis for future clinical application.

REFERENCES

- 1 Sairenji T, Collins KL, Evans DV. An Update on Inflammatory Bowel Disease. *Prim Care* 2017; **44**: 673-692 [PMID: 29132528 DOI: 10.1016/j.pop.2017.07.010]
- 2 Yu YR, Rodriguez JR. Clinical presentation of Crohn's, ulcerative colitis, and indeterminate colitis: Symptoms, extraintestinal manifestations, and disease phenotypes. *Semin Pediatr Surg* 2017; **26**: 349-355 [PMID: 29126502 DOI: 10.1053/j.sempedsurg.2017.10.003]
- 3 Ananthkrishnan AN. Epidemiology and risk factors for IBD. *Nat Rev Gastroenterol Hepatol* 2015; **12**: 205-217 [PMID: 25732745 DOI: 10.1038/nrgastro.2015.34]
- 4 Ungaro R, Mehandru S, Allen PB, Peyrin-Biroulet L, Colombel JF. Ulcerative colitis. *Lancet* 2017; **389**: 1756-1770 [PMID: 27914657 DOI: 10.1016/S0140-6736(16)32126-2]
- 5 Garcia-Planella E, Mañosa M, Van Domselaar M, Gordillo J, Zabana Y, Cabré E, López San Román A, Domènech E. Long-term outcome of ulcerative colitis in patients who achieve clinical remission with a first course of corticosteroids. *Dig Liver Dis* 2012; **44**: 206-210 [PMID: 22079262 DOI: 10.1016/j.dld.2011.10.004]
- 6 Vindigni SM, Zisman TL, Suskind DL, Damman CJ. The intestinal microbiome, barrier function, and immune system in inflammatory bowel disease: a tripartite pathophysiological circuit with implications for new therapeutic directions. *Therap Adv Gastroenterol* 2016; **9**: 606-625 [PMID: 27366227 DOI: 10.1177/1756283X16644242]
- 7 Merga Y, Campbell BJ, Rhodes JM. Mucosal barrier, bacteria and inflammatory bowel disease: possibilities for therapy. *Dig Dis* 2014; **32**: 475-483 [PMID: 24969297 DOI: 10.1159/000358156]
- 8 Keane TJ, Dziki J, Sobieski E, Smoulder A, Castleton A, Turner N, White LJ, Badylak SF. Restoring Mucosal Barrier Function and Modifying Macrophage Phenotype with an Extracellular Matrix Hydrogel: Potential Therapy for Ulcerative Colitis. *J Crohns Colitis* 2017; **11**: 360-368 [PMID: 27543807 DOI: 10.1093/ecco-jcc/jjw149]
- 9 Sina C, Kemper C, Derer S. The intestinal complement system in inflammatory bowel disease: Shaping intestinal barrier function. *Semin Immunol* 2018; **37**: 66-73 [PMID: 29486961 DOI: 10.1016/j.smim.2018.02.008]
- 10 Javid AZ, Hormoznejad R, Yousefimanesh HA, Haghighi-Zadeh MH, Zakerkish M. Impact of resveratrol supplementation on inflammatory, antioxidant, and periodontal markers in type 2 diabetic patients with chronic periodontitis. *Diabetes Metab Syndr* 2019; **13**: 2769-2774 [PMID: 31405706 DOI: 10.1016/j.dsx.2019.07.042]
- 11 De Oliveira MTP, de Sá Coutinho D, Tenório de Souza É, Stanisquaski Guterres S, Pohlmann AR, Silva PMR, Martins MA, Bernardi A. Orally delivered resveratrol-loaded lipid-core nanocapsules ameliorate LPS-induced acute lung injury *via* the ERK and PI3K/Akt pathways. *Int J Nanomedicine* 2019; **14**: 5215-5228 [PMID: 31371957 DOI: 10.2147/IJN.S200666]
- 12 Zimmermann-Franco DC, Esteves B, Lacerda LM, Souza IO, Santos JAD, Pinto NCC, Scio E, da Silva AD, Macedo GC. In vitro and in vivo anti-inflammatory properties of imine resveratrol analogues. *Bioorg Med Chem* 2018; **26**: 4898-4906 [PMID: 30193941 DOI: 10.1016/j.bmc.2018.08.029]
- 13 Samsami-Kor M, Daryani NE, Asl PR, Hekmatdoost A. Anti-Inflammatory Effects of Resveratrol in Patients with Ulcerative Colitis: A Randomized, Double-Blind, Placebo-controlled Pilot Study. *Arch Med Res* 2015; **46**: 280-285 [PMID: 26002728 DOI: 10.1016/j.arcmed.2015.05.005]
- 14 Samsamikor M, Daryani NE, Asl PR, Hekmatdoost A. Resveratrol Supplementation and Oxidative/Anti-Oxidative Status in Patients with Ulcerative Colitis: A Randomized, Double-Blind, Placebo-controlled Pilot Study. *Arch Med Res* 2016; **47**: 304-309 [PMID: 27664491 DOI: 10.1016/j.arcmed.2016.07.003]
- 15 Levine B, Mizushima N, Virgin HW. Autophagy in immunity and inflammation. *Nature* 2011; **469**: 323-335 [PMID: 21248839 DOI: 10.1038/nature09782]
- 16 Macias-Ceja DC, Cosin-Roger J, Ortiz-Masiá D, Salvador P, Hernández C, Esplugues JV, Calatayud S, Barrachina MD. Stimulation of autophagy prevents intestinal mucosal inflammation and ameliorates murine colitis. *Br J Pharmacol* 2017; **174**: 2501-2511 [PMID: 28500644 DOI: 10.1111/bph.13860]
- 17 Ji G, Wang Y, Deng Y, Li X, Jiang Z. Resveratrol ameliorates hepatic steatosis and inflammation in methionine/choline-deficient diet-induced steatohepatitis through regulating autophagy. *Lipids Health Dis* 2015; **14**: 134 [PMID: 26498332 DOI: 10.1186/s12944-015-0139-6]

- 18 **Chen ML**, Yi L, Jin X, Liang XY, Zhou Y, Zhang T, Xie Q, Zhou X, Chang H, Fu YJ, Zhu JD, Zhang QY, Mi MT. Resveratrol attenuates vascular endothelial inflammation by inducing autophagy through the cAMP signaling pathway. *Autophagy* 2013; **9**: 2033-2045 [PMID: 24145604 DOI: 10.4161/autophagy.26336]
- 19 **Chen L**, Zhou Z, Yang Y, Chen N, Xiang H. Therapeutic effect of imiquimod on dextran sulfate sodium-induced ulcerative colitis in mice. *PLoS One* 2017; **12**: e0186138 [PMID: 29049372 DOI: 10.1371/journal.pone.0186138]
- 20 **Nunes NS**, Kim S, Sundby M, Chandran P, Burks SR, Paz AH, Frank JA. Temporal clinical, proteomic, histological and cellular immune responses of dextran sulfate sodium-induced acute colitis. *World J Gastroenterol* 2018; **24**: 4341-4355 [PMID: 30344419 DOI: 10.3748/wjg.v24.i38.4341]
- 21 **Eichele DD**, Kharbanda KK. Dextran sodium sulfate colitis murine model: An indispensable tool for advancing our understanding of inflammatory bowel diseases pathogenesis. *World J Gastroenterol* 2017; **23**: 6016-6029 [PMID: 28970718 DOI: 10.3748/wjg.v23.i33.6016]
- 22 **Peterson LW**, Artis D. Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat Rev Immunol* 2014; **14**: 141-153 [PMID: 24566914 DOI: 10.1038/nri3608]
- 23 **Dokladny K**, Zuhl MN, Moseley PL. Intestinal epithelial barrier function and tight junction proteins with heat and exercise. *J Appl Physiol (1985)* 2016; **120**: 692-701 [PMID: 26359485 DOI: 10.1152/jappphysiol.00536.2015]
- 24 **Capaldo CT**, Powell DN, Kalman D. Layered defense: how mucus and tight junctions seal the intestinal barrier. *J Mol Med (Berl)* 2017; **95**: 927-934 [PMID: 28707083 DOI: 10.1007/s00109-017-1557-x]
- 25 **Cui X**, Jin Y, Hofseth AB, Pena E, Habiger J, Chumanevich A, Poudyal D, Nagarkatti M, Nagarkatti PS, Singh UP, Hofseth LJ. Resveratrol suppresses colitis and colon cancer associated with colitis. *Cancer Prev Res (Phila)* 2010; **3**: 549-559 [PMID: 20332304 DOI: 10.1158/1940-6207.CAPR-09-0117]
- 26 **Yao J**, Wei C, Wang JY, Zhang R, Li YX, Wang LS. Effect of resveratrol on Treg/Th17 signaling and ulcerative colitis treatment in mice. *World J Gastroenterol* 2015; **21**: 6572-6581 [PMID: 26074695 DOI: 10.3748/wjg.v21.i21.6572]
- 27 **Abdin AA**. Targeting sphingosine kinase 1 (SphK1) and apoptosis by colon-specific delivery formula of resveratrol in treatment of experimental ulcerative colitis in rats. *Eur J Pharmacol* 2013; **718**: 145-153 [PMID: 24055189 DOI: 10.1016/j.ejphar.2013.08.040]
- 28 **Deretic V**, Saitoh T, Akira S. Autophagy in infection, inflammation and immunity. *Nat Rev Immunol* 2013; **13**: 722-737 [PMID: 24064518 DOI: 10.1038/nri3532]
- 29 **Lassen KG**, Xavier RJ. Mechanisms and function of autophagy in intestinal disease. *Autophagy* 2018; **14**: 216-220 [PMID: 29130415 DOI: 10.1080/15548627.2017.1389358]
- 30 **Saito M**, Katsuno T, Nakagawa T, Sato T, Noguchi Y, Sazuka S, Saito K, Arai M, Yokote K, Yokosuka O. Intestinal epithelial cells with impaired autophagy lose their adhesive capacity in the presence of TNF- α . *Dig Dis Sci* 2012; **57**: 2022-2030 [PMID: 22466076 DOI: 10.1007/s10620-012-2133-4]
- 31 **Zhang L**, Xue H, Zhao G, Qiao C, Sun X, Pang C, Zhang D. Curcumin and resveratrol suppress dextran sulfate sodium-induced colitis in mice. *Mol Med Rep* 2019; **19**: 3053-3060 [PMID: 30816479 DOI: 10.3892/mmr.2019.9974]
- 32 **Wan SX**, Shi B, Lou XL, Liu JQ, Ma GG, Liang DY, Ma S. Ghrelin protects small intestinal epithelium against sepsis-induced injury by enhancing the autophagy of intestinal epithelial cells. *Biomed Pharmacother* 2016; **83**: 1315-1320 [PMID: 27571874 DOI: 10.1016/j.biopha.2016.08.048]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

