

World Journal of *Hepatology*

World J Hepatol 2020 November 27; 12(11): 883-1135



REVIEW

- 883 Malnutrition in cirrhosis: More food for thought
Chapman B, Sinclair M, Gow PJ, Testro AG

MINIREVIEWS

- 897 Paraneoplastic syndromes in cholangiocarcinoma
Rahman SU, Sana MK, Tahir Z, Ali A, Shah PA
- 908 Noninvasive scores for the prediction of esophageal varices and risk stratification in patients with cirrhosis
Bangaru S, Benhammou JN, Tabibian JH
- 919 Natrema and liver transplantation: The right amount of salt for a good recipe
Lenci I, Milana M, Grassi G, Signorello A, Aglitti A, Baiocchi L

ORIGINAL ARTICLE**Basic Study**

- 931 Inhibition of vascular adhesion protein-1 modifies hepatic steatosis *in vitro* and *in vivo*
Shepherd EL, Karim S, Newsome PN, Lalor PF
- 949 Aceclofenac-induced hepatotoxicity: An ameliorative effect of *Terminalia bellirica* fruit and ellagic acid
Gupta A, Pandey A
- 965 Obeticholic acid attenuates human immunodeficiency virus/alcohol metabolism-induced pro-fibrotic activation in liver cells
New-Aaron M, Ganesan M, Dagur RS, Kharbanda KK, Poluektova LY, Osna NA
- 976 Screening and identification of bioactive compounds from citrus against non-structural protein 3 protease of hepatitis C virus genotype 3a by fluorescence resonance energy transfer assay and mass spectrometry
Khan M, Rauf W, Habib F, Rahman M, Iqbal M

Retrospective Cohort Study

- 993 Cannabis use history is associated with increased prevalence of ascites among patients with nonalcoholic fatty liver disease: A nationwide analysis
Choi CJ, Weiss SH, Nasir UM, Prysopoulos NT
- 1004 Phase angle and non-alcoholic fatty liver disease before and after bariatric surgery
Teixeira J, Marroni CA, Zubiaurre PR, Henz A, Faina L, Pinheiro LK, Mottin CC, Fernandes SA

Retrospective Study

- 1020** Factors associated with 5-year survival of combined hepatocellular and cholangiocarcinoma
Sempokuya T, Wien EA, Pattison RJ, Ma J, Wong LL
- 1031** Circulating miR-21-5p level has limited prognostic value in patients with hepatocellular carcinoma and is influenced by renal function
Franck M, Thon C, Schütte K, Malfertheiner P, Link A
- 1046** Real impact of tumor marker AFP and PIVKA-II in detecting very small hepatocellular carcinoma (≤ 2 cm, Barcelona stage 0) - assessment with large number of cases
Tarao K, Nozaki A, Komatsu H, Komatsu T, Taguri M, Tanaka K, Chuma M, Numata K, Maeda S
- 1055** Non-invasive splenic parameters of portal hypertension: Assessment and utility
Ahmad AK, Atzori S, Maurice J, Taylor-Robinson SD, Lim AKP
- 1067** Outcome of gastric antral vascular ectasia and related anemia after orthotopic liver transplantation
Emhmed Ali S, Benrajab KM, Dela Cruz AC

Clinical Trials Study

- 1076** Hepatitis B surface antigen and hepatitis B core-related antigen kinetics after adding pegylated-interferon to nucleos(t)ids analogues in hepatitis B e antigen-negative patients
Broquetas T, Garcia-Retortillo M, Micó M, Canillas L, Puigvehí M, Cañete N, Coll S, Viu A, Hernandez JJ, Bessa X, Carrión JA

Observational Study

- 1089** Occurrence of seeding metastases in resectable perihilar cholangiocarcinoma and the role of low-dose radiotherapy to prevent this
Franken LC, Roos E, Saris J, van Hooft JE, van Delden OM, Verheij J, Erdmann JI, Besselink MG, Busch OR, van Tienhoven G, van Gulik TM

Randomized Controlled Trial

- 1098** Metalloproteinase expression after desflurane preconditioning in hepatectomies: A randomized clinical trial
Koraki E, Mantzoros I, Chatzakis C, Gkiouliava A, Cheva A, Lavrentieva A, Sifaki F, Argiriadou H, Kesisoglou I, Galanos-Demiris K, Bitsianis S, Tsalis K

SYSTEMATIC REVIEWS

- 1115** Clinical utility of viscoelastic testing in chronic liver disease: A systematic review
Wei H, Child LJ

CASE REPORT

- 1128** Hepatocellular carcinoma with tumor thrombus extends to the right atrium and portal vein: A case report
Gomez-Puerto D, Mirallas O, Vidal-González J, Vargas V

ABOUT COVER

Associate editor of *World Journal of Hepatology*, Dr. Yong-Ping Yang is a Distinguished Professor at Peking University Health Science Center in Beijing, China. Having received his Bachelor's degree from Yanbian University in 1985, Dr. Yang undertook his postgraduate training at PLA Medical College, receiving his Master's degree in 1992. He rose to Chief Physician in the Hepatology Division of the Fifth Medical Center of the Chinese PLA General Hospital in 2003 and has held the position since. His ongoing research interests involve liver fibrosis, cirrhosis and hepatocellular carcinoma, with a particular focus on cryoablation and cryo-immunotherapy for hepatocellular carcinoma. Currently, he serves as Chairman of the Department of Liver Disease of the Chinese PLA General Hospital and as President of the Chinese Research Hospital Association for the Study of the Liver Disease. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Hepatology* (*WJH*, *World J Hepatol*) is to provide scholars and readers from various fields of hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJH mainly publishes articles reporting research results and findings obtained in the field of hepatology and covering a wide range of topics including chronic cholestatic liver diseases, cirrhosis and its complications, clinical alcoholic liver disease, drug induced liver disease autoimmune, fatty liver disease, genetic and pediatric liver diseases, hepatocellular carcinoma, hepatic stellate cells and fibrosis, liver immunology, liver regeneration, hepatic surgery, liver transplantation, biliary tract pathophysiology, non-invasive markers of liver fibrosis, viral hepatitis.

INDEXING/ABSTRACTING

The *WJH* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Li-Li Wang*; Production Department Director: *Yun-Xiaojuan Wu*; Editorial Office Director: *Jia-Ping Yan*.

NAME OF JOURNAL

World Journal of Hepatology

ISSN

ISSN 1948-5182 (online)

LAUNCH DATE

October 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Nikolaos Pylsopoulos, Ke-Qin Hu, Koo Jeong Kang

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5182/editorialboard.htm>

PUBLICATION DATE

November 27, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Paraneoplastic syndromes in cholangiocarcinoma

Saad Ur Rahman, Muhammad Khawar Sana, Zabreen Tahir, Aamir Ali, Pir Ahmad Shah

ORCID number: Saad Ur Rahman 0000-0002-4838-0802; Muhammad Khawar Sana 0000-0003-1952-8203; Zabreen Tahir 0000-0001-5342-6847; Aamir Ali 0000-0001-8024-5865; Pir Ahmad Shah 0000-0002-2224-2401.

Author contributions: Shah PA and Tahir Z designed the study. Rahman SU and Sana MK did database screening and data extraction. Tahir Z, Shah PA, Rahman SU, Ali A, and Sana MK did the literature review, analyzed the data, and wrote the manuscript. Rahman SU, Sana MK, Ali A, Tahir Z, and Shah PA reviewed the paper for final submission.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Saad Ur Rahman, Muhammad Khawar Sana, Department of Medicine, King Edward Medical University, Lahore 54000, Punjab, Pakistan

Zabreen Tahir, Department of Neurology, Brigham and Women's Hospital, Boston, MA 02115, United States

Aamir Ali, Department of Radiology, Beth Israel Deaconess Medical Center, Boston, MA 02215, United States

Pir Ahmad Shah, Department of Medicine, University of Texas Health Science Center, San Antonio, TX 78229, United States

Corresponding author: Pir Ahmad Shah, MD, Internal Medicine Resident, Department of Medicine, University of Texas Health Science Center, 7703 Floyd Curl Dr, San Antonio, TX 78229, United States. pirashah2020@gmail.com

Abstract

Paraneoplastic syndromes are the symptoms or signs which result from damage to tissues that are distant from the site of malignancy, due to complex interactions between the body's immune system and malignant neoplasm. Cholangiocarcinoma (CCA) is an aggressive epithelial malignancy of hepatobiliary tree and it is found to be associated with various paraneoplastic syndromes. These syndromes can present as dermatological, neurological, renal, hematological, or multi-systemic manifestations. Clinical suspicion and timely recognition of these syndromes can lead to early diagnosis of covert malignancies like CCA. The management plan remains the removal of the underlying cause which in this case is CCA.

Key Words: Cholangiocarcinoma; Paraneoplastic syndrome; Malignancy; Immune system; Biliary tree; Multi-organ

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Various case reports have been published focusing on single paraneoplastic syndromes associated with cholangiocarcinoma (CCA) but none of the studies has reported these manifestations collectively. This review summarizes different paraneoplastic syndromes which are associated with CCA to give a better idea about how it affects various organ systems.

Manuscript source: Invited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: United States

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

Received: June 16, 2020

Peer-review started: June 16, 2020

First decision: July 30, 2020

Revised: August 13, 2020

Accepted: September 8, 2020

Article in press: September 8, 2020

Published online: November 27, 2020

P-Reviewer: Coulouarn C, Dourakis SP

S-Editor: Zhang H

L-Editor: A

P-Editor: Wang LL



Citation: Rahman SU, Sana MK, Tahir Z, Ali A, Shah PA. Paraneoplastic syndromes in cholangiocarcinoma. *World J Hepatol* 2020; 12(11): 897-907

URL: <https://www.wjgnet.com/1948-5182/full/v12/i11/897.htm>

DOI: <https://dx.doi.org/10.4254/wjh.v12.i11.897>

INTRODUCTION

Cholangiocarcinoma (CCA) is an aggressive epithelial malignancy of hepatobiliary tree, accounting for 10%-20% of primary liver cancers^[1,2]. It is strongly linked to chronic liver disease and is classified according to an anatomical location in the biliary tree as intrahepatic (iCCA), perihilar (pCCA) and distal (dCCA) subtypes^[3]. The incidence of CCA, particularly iCCA has increased worldwide between 1993 and 2002. Highest rates were found among Asian countries with South Korea on the top having an age-standardized incidence rate (ASR) of 2.80, followed by Thailand (2.19) and Japan (0.95)^[4]. In the USA, the incidence of iCCA has increased in the last 40 years between 1973 and 2012 from 0.44 to 1.18 cases per 100000 person-years^[5]. iCCA accounts for about 20% of the deaths from hepatobiliary cancers, which cause 13% of the total cancer mortality worldwide^[6]. Recent diagnostic techniques and early management have led to an improvement in 1-year mortality over time but the 5-year survival is still as low as 10% due to the appearance of clinical symptoms in the later course of the disease^[7].

Paraneoplastic syndromes are the symptoms or signs which result from damage to tissues that are remote from the site of malignancy, due to complex interactions between the body's immune system and malignant neoplasm^[8]. CCA has been reportedly found to be the source of various paraneoplastic manifestations including alopecia^[9], sensory neuropathies^[10], hypercalcemia, polycythemia, leukocytosis^[11] and increased parathyroid hormone-related protein (PTHrP)^[12-14].

Several individual cases of paraneoplastic syndrome with CCA have been reported in the literature. This mini-review summarizes all these cases with their clinical presentation and pathophysiology associated with CCA.

PARANEOPLASTIC SYNDROMES

Dermatological manifestations

Acanthosis nigricans: Acanthosis nigricans (AN) presents as a brown to black hyperpigmented velvety patch found on neck, groin, and axilla^[15]. However, AN associated with internal organ malignancy mostly appears as a diffuse patch over palms and soles, and can also involve the oral cavity and/or esophagus^[16]. Of all the reported cases, internal organ malignancies are found to be associated with AN of palms in 90% of the cases^[17]. Paraneoplastic syndrome that occurs as a result of CCA is associated with the production of biologically active particles in the malignant tissue. These include growth factors like epidermal growth factor (EGF) or alpha-transforming growth factor (αTGF) which are associated with the malignant proliferation of skin resulting in AN^[18]. AN resolves after effective treatment of the underlying malignancy and might progress if there is any recurrence or metastasis of the tumor as stated in the study of Ravnborg.

Alopecia: Alopecia is the most common hair disease reported in oncology patients^[19]. It usually presents as a round patch of hair loss on the top of the skull with well-demarcated edges. The type of alopecia associated with cholangiocarcinoma is usually alopecia areata^[20]. Alopecia in CCA is triggered by several neurological, hormonal, and emotional factors but in most cases, the actual cause remains undetermined. Alopecia caused by CCA has an increasing trend of incidence in the people of the United States as compared to those in Europe hinting towards the role of environmental factors as well^[21]. Although it can occur as an independent condition, but remission of cancer resulting in the resolution of alopecia, and relapse leading to recurrence favors the paraneoplastic nature as mentioned in the study of Antoniou^[20].

Dermatomyositis: It is relatively a rare paraneoplastic manifestation of CCA^[22]. It presents as weakness of the proximal group of muscles with a bluish-purple skin lesion on upper eyelids (heliotrope rash) and erythematous papules on knuckles are noticed (Gottron papules) as evident from the study of Suh^[23]. Muscle weakness in

dermatomyositis caused by CCA is attributed to increasing energy consumption of carcinoma cells than normal cells, resulting in excessive glucose breakdown^[24]. Dermatomyositis also results from an autoimmune response against highly active cancer cells that cross-react and attack the affected muscles^[25]. Myositis specific auto-antigens are also expressed in tumor cells that cross-react and cause this disease. Successful treatment of CCA results in the improvement of dermatomyositis^[26].

Disseminated superficial porokeratosis: Disseminated superficial porokeratosis (DSP) is a benign proliferation of keratinocytes resulting in a hyperkeratotic skin tumor^[27]. DSP is linked to internal organ malignancies such as hepatocellular carcinoma and CCA suggesting a paraneoplastic nature^[28]. DSP presents as well-demarcated reddish-brown papules, ranging from 0.5 cm to 1.5 cm, and become noticeable a few months before the onset of cancer-related symptoms. These are visible on the extensor surface of all limbs and trunk, consisting of multiple annular and itchy lesions with elevated borders sparing mucosa, palms, and soles as explained by Cannavo in his study^[29]. Overexpression of the *p53* gene product is associated with the appearance of widespread DSP as a manifestation of internal organ malignancy like CCA^[30]. All of these abnormalities in the *p53* pathway result in a lack of keratinocyte differentiation and dysregulation of loricrin expression. Loricrin is a precursor protein formed in the last stage of keratinization and its dysregulation results in the formation of a cornified envelope^[31].

Necrolytic Migratory Erythema: Necrolytic migratory erythema (NME) is an erythematous erosive patch with an advancing scaly border^[32]. NME in the setting of CCA, presents as annular erythematous lesions of almost 1-2 cm in size with a central glassy surface, surrounded by scaling that occurs on the face, trunk, and extremities as mentioned by Chiyomaru in his case report. NME mostly presents before CCA is clinically diagnosed^[33]. Several conditions like primary sclerosing cholangitis, liver fluke infection, and biliary malformation are associated with increased risk of CCA which results in paraneoplastic NME^[34]. Moreover, low nutritious conditions are also aiding the development of NME. Thus patients with NME, without a known cause, should always be checked and screened for malignancies periodically^[33].

Persistent erythema multiforme: Erythema multiforme (EM) is one of the cutaneous disorders that occur in the setting of a pre-existing internal organ malignancy and are collectively known as paraneoplastic dermatoses. EM is typically self-limited and benign. It usually occurs early in the disease course of CCA. Persistent erythema multiforme (PEM) is a rare form of erythema multiforme that comprises of both typical and atypical cutaneous/mucosal lesions and doesn't resolve on its own^[35]. On examination, EM appears as a painful erythematous rash with scaling that occurs in patches over the chest, upper back, and both thighs. After a few days, it changes to hemorrhagic bullae with violaceous edges. Vital signs are normal and there is no mucosal involvement as shown by the study of Tzovaras^[36]. EM is sudden in onset and usually resolves in 1- 6 d. Skin paraneoplastic syndromes are commonly associated with internal organ malignancies but EM is a rare skin manifestation in these cases. It is believed that PEM occurs in the setting of CCA as a result of continuous stimulation by antigenic tumor material^[37]. There is no direct involvement of tumor cells and no other acquired factors are present marking REM as a paraneoplastic manifestation of CCA. Moreover, the treatment of CCA results in regression of PEM, and relapse of the tumor results in its reappearance^[36].

Sweet syndrome: Sweet syndrome (SS) is an acute febrile neutrophilic dermatosis. Malignancy associated Sweet syndrome occurs in approximately 15% of the population suffering from solid tumors^[38]. It presents as rapidly growing painful erythematous plaques over the face, neck, and legs and is associated with fever, generalized malaise, cough, and arthralgia according to the case report presented by Shinjima^[39]. According to recent studies, it is observed that granulocyte-colony stimulating factor (G-CSF) has a major role in the pathogenesis of SS^[40]. Increased G-CSF results in production, activation, and chemotaxis of the neutrophils^[41]. Malignant tumors like CCA result in excessive production of G-CSF that further stimulates neutrophils resulting in paraneoplastic SS^[42].

Bazex syndrome: Bazex syndrome is characterized by the appearance of hyperkeratotic lesions on various parts of the body in association with an underlying malignancy. It clinically presents as pruritic scaly dusky red eruptions covered with adherent scales on face, ear, buttocks, palms, and soles. They are commonly associated with fatigue, recurrent abdominal pain, nausea, vomiting, constipation, weight loss,

and liver enlargement as mentioned by Karabulut in his study^[43]. Pathophysiology of bazex syndromes is cross-reaction of antigens from tumor to skin. This cellular immune system alteration results in increase release of various growth factors like epidermal growth factor- α ^[44]. Treatment of bazex syndrome shows how strongly this manifestation is paraneoplastic. Regular dermatological treatment doesn't show any response but the removal of the underlying tumor results in complete resolution of the disease^[45].

Erythema gyratum: Erythema gyratum is rapidly moving erythema and is a marker of underlying malignancy. It is most commonly associated with lung, esophageal, and breast carcinoma^[46]. Clinically it presents as a 3-wk history of rash on the lower limbs. The eruption of the rash began as small erythematous macules which gradually enlarge resulting in concentric raised scales on the lateral aspect of the right thigh as Liau mentioned in his study^[47].

Pityriasis rubra pilaris: Pityriasis rubra pylaris (PRP) is a papulosquamous dermatosis of skin. It clinically presents as a 10-d history of widespread pruritic rash that begins appearing on thighs and progresses gradually over shins, lower back, trunk, face, and forearms as mentioned in the case report presented by Bar-Ilan. It is associated with mild fever as well^[48]. Pathophysiology involved in the appearance of PRP is increased secretion of peptides and hormones from the tumor due to cross-reactivity of the antigens^[49].

Sign of Leser-trelat: Leser-trelat sign is the appearance of multiple pigmented seborrheic keratosis mostly associated with underlying malignancy. It is mostly associated with GI adenocarcinoma and rarely with CCA^[50]. It clinically presents as a 1-week history of worsening jaundice, pale stools, and recent onset abdominal pain as evident by the study of Morgenthau^[51]. The pathophysiology behind this sign is a sudden increase in cytokines and various growth factors like epidermal growth factor- α resulting in hyperpigmentation of the skin^[52].

Subacute cutaneous lupus erythematosus: Subacute cutaneous lupus erythematosus (SCLE) is an inflammatory skin disorder mimicking skin manifestations of systemic lupus erythematosus. It is one of the rare paraneoplastic manifestations linked to CCA. It clinically presents as explosive onset of new pruritic rash along with arthralgia and lower extremity edema in a patient with previous history of CCA as explained by Opneja in his study^[53]. The pathophysiology behind SCLE lies in self-activation of the body's immune system resulting in photosensitive rash as it is in the usual SLE^[54].

A summary of all the dermatological paraneoplastic syndromes is explained in [Table 1](#) at the bottom of the review

Neurological manifestations

Limbic encephalopathy: Limbic encephalopathy (LE) is the sub-acute onset of memory impairment and confusion^[55]. LE is reported in CCA but is a rare finding. In the starting phase, it presents as a polyneuropathy quite similar to diabetic polyneuropathy, with gradual progression to focal seizures. As the disease progresses, temporal lobe association is noted resulting in pilomotor erection (autonomic seizures) and eventually symptoms of rapidly progressive dementia appear^[56]. Thus, autonomic seizures, delusion, and rapidly progressive dementia are the hallmarks of LE^[57]. Different mechanisms are linked to LE in the setting of CCA. The formation of new anti-neuronal antibodies is associated with limbic encephalitis. In some cases, autoantibodies are formed against intracellular antigens in the mesiotemporal region while others suggest that they are formed against surface antigens^[58]. Both hypermetabolism and hypometabolism of mesiotemporal lobe has also been reported after the onset of LE symptoms^[59].

Paraneoplastic cerebellar degeneration: Paraneoplastic cerebellar degeneration (PCD) typically presents in women with a sudden onset of ataxia progressively involving limbs and trunks, dysarthria, diplopia, and dysphagia that occurs in the background of a malignancy. It is rarely reported with CCA. In a case report by Bruhnding *et al.* it began with lower limbs, but gradually involved upper limbs as well. Sensation in legs was affected as well. Eventually, the patient was unable to stand anymore. Dysmetria ensued and interfered with self-sufficiency in feeding leading to a weight loss of 60 pounds over six months. Imaging revealed a 2 cm liver mass. Biopsy proved intrahepatic cholangiocarcinoma. Anti-Yo antibodies were also found positive in association with PCD^[60]. The underlying mechanism is not clearly understood. The

Table 1 Summary of literature on dermatological paraneoplastic syndromes in cholangiocarcinoma

Ref.	Year	Paraneoplastic syndrome	Mediator
Dermatologic			
Ravnborg <i>et al</i> ^[15]	1993	Acanthosis nigricans	TGF-alpha
Suchonwanit <i>et al</i> ^[19]	2018	Alopecia	T-lymphocytes
Antoniou <i>et al</i> ^[20]	2012	Alopecia	T-lymphocytes
Suh <i>et al</i> ^[23]	2016	Dermatomyositis	
Yasuda <i>et al</i> ^[26]	2018	Dermatomyositis	
Sotoodian <i>et al</i> ^[27]	2018	Disseminated superficial porokeratosis	p53
Cannavó <i>et al</i> ^[29]	2008	Disseminated superficial porokeratosis	p53
Chiyomaru <i>et al</i> ^[33]	2010	Necrolytic migratory erythema	
Tzovaras <i>et al</i> ^[36]	2007	Persistent erythema multiforme	
Shinojima <i>et al</i> ^[39]	2006	Sweet syndrome	G-CSF, IL-1, IL-6
Karabulut <i>et al</i> ^[43]	2006	Bazex syndrome	
Liau <i>et al</i> ^[47]	2016	Erythema gyratum	
Bar-Ilan <i>et al</i> ^[48]	2017	Pityriasis rubra pilaris	
Morgenthau <i>et al</i> ^[51]	2019	Sign of Leser- Trelat	EGF-alpha
Opneja <i>et al</i> ^[53]	2015	Subacute cutaneous lupus erythematosus	

TGF-alpha: Tissue growth factor-alpha; G-CSF: Granulocyte-colony stimulating factor; IL: Interleukin; EGF-alpha: Epidermal growth factor-alpha.

autoimmune nature of PCD is thought to be due to malignant cells expressing onconeural antigens that are otherwise found on neurons^[61]. Thus cross-reactivity leads to the development of PCD. However, no direct link between these antibodies and PCD has been developed yet.

Renal manifestations

Glomerulonephritis: Glomerulonephritis is defined as inflammation of small blood vessels inside the kidneys. Fibrillary glomerulonephritis (FGN) is a rare type of glomerulonephritis. A rare case of FGN is recently reported showing its association with iCCA. It presents as edema of lower limbs and face with uncontrolled hypertension. Nephrotic range proteinuria is evident (3 g/d) with 24 h-proteinuria of 0.74g/d mentioned by Normand in his study. Microscopic hematuria is also present. Complete remission of glomerulonephritis indicates the paraneoplastic nature of this disease^[62].

Hematological manifestations

Paraneoplastic vasculitis: Vasculitis is an inflammation of the wall of a blood vessel. Malignant diseases are both associated with vasculitis of arteries and veins. Paraneoplastic vasculitis constitutes less than 5% of all the forms of vasculitis^[63]. Vasculitis is more commonly associated with hematological malignancies than solid tumors^[64]. Small vessels are frequently linked to the paraneoplastic nature of vasculitis. The type of vasculitis associated with CCA is giant cell arteritis. It presents with one-month history of headache, scalp tenderness, pain, and stiffness in the neck, shoulder, and pelvic girdles. The resolution of symptoms right after removal of the tumor indicated the paraneoplastic nature of this vasculitis^[65].

Trousseau syndrome: Trousseau syndrome, also known as migratory thrombophlebitis, is an acquired abnormality of blood clotting. According to several reports, it is concluded that several clotting disorders are closely linked to internal organ malignancy^[66]. There have been 2 cases of trousseau syndrome with underlying isolated CCA, while others had either a hepatocellular CA or a lung adenocarcinoma along with CCA^[67]. Trousseau syndrome, in the setting of CCA, presents as weight loss, mild shortness of breath, right upper quadrant tenderness, and abnormal liver

function tests with raised alkaline phosphatase as evident from the studies of Jang and Blum^[68]. It is believed that tissue hypoxia leads to activation of the coagulation pathway and endothelial adhesion molecules^[69]. Low molecular weight heparin (LMWH) along with removal of the primary tumor has been found effective in its treatment^[70].

Anti-phospholipid antibody syndrome: Antiphospholipid antibody syndrome (APAS) is an autoimmune disorder that results in the formation of antibodies against phospholipids on platelets resulting in hypercoagulability. APAS has been linked to various solid organ malignancies but a few cases are found in association with CCA^[71]. APAS, in the setting of CCA, presents as unilateral leg pain, swelling, and tenderness. Moreover, lupus anticoagulant is raised with normal antinuclear antibodies as mentioned by Samadian in his case report^[72].

Paraneoplastic leukemoid reaction: Leukemoid reaction means an increase in the number of leukocytes (WBC's > 50000) particularly neutrophils, in reaction to any infection or carcinoma. Leukemoid reaction in the setting of CCA mimics a pyogenic liver abscess and clinically presents as pyrexia and leukocytosis^[73]. The fever is intermittent and remains there for at least a month along with weight loss, progressive generalized weakness, and a leukocyte count above 20000 (> 78% neutrophils) mentioned by Ham in his study related to leukemoid reaction^[74]. Only 2 cases of paraneoplastic leukemoid reaction have been reported with the primary cause of CCA^[75]. Treating CCA resulted in the resolution of leukemoid reaction.

Multisystemic manifestations

Polyarteritis nodosa: Polyarteritis nodosa (PAN) is a rare form of systemic necrotizing vasculitis that has heterogeneous forms of presentations^[76]. PAN that occurs due to underlying neoplasia is termed as paraneoplastic vasculitis. It may precede or follow the onset of neoplasia or it may also be evident on the recurrence of many malignant diseases, showing a strong link with carcinomas^[77]. PAN occurring in the setting of CCA is a rare finding but a study reports an association of CCA with PAN^[78]. Paraneoplastic vasculitis is mostly cutaneous but it can also affect internal organs^[79]. The earliest sign experienced in PAN is bilateral numbness in lower limbs followed by gradually increasing fever. After approximately 2 wk, an arthritis-like pathology is noticed with severe pain in bilateral lower limbs, ankles, metatarsal, and phalangeal joints. After a few months, skin manifestations appear which are necrosis and gangrene evident on distal phalanges. Lastly, gastrointestinal symptoms appear comprising of severe abdominal pain, nausea, and vomiting. The gradual sequence of these clinical symptoms has been mentioned by Hatzis in his case report. Digital ischemia is also a complication of paraneoplastic vasculitis^[80]. Paraneoplastic vasculitis such as PAN is highly associated with raise in titers of anti-neutrophilic cytoplasmic antibodies (ANCA)^[81]. Moreover, patients with raised ANCA and PAN also have raised CA 19-9, a tumor marker that rises in CCA and pancreatic carcinomas^[82]. Recent studies suggest immune dysregulation as the primary cause of paraneoplastic vasculitis. Another study states that it might be due to cross-reaction of tumor antigens, directly causing vascular damage or indirectly by releasing humoral agents like chemotactic factors. PAN is poorly responsive to steroids or other treatment modalities but the removal of the tumor or chemotherapy results in its resolution^[83].

Adult-onset still disease: Adult-onset Still disease (AOSD) is an adult version of juvenile idiopathic arthritis. It is caused by an altered immune response of the body against any foreign body or carcinoma. Previously, AOSD has been linked to various carcinomas like breast, esophageal, thyroid, and lung^[84]. This is the first case, reporting link of AOSD with CCA. Symptoms of AOSD occur before malignancy is diagnosed^[85]. AOSD linked to CCA clinically presents as high-grade fever and chills for 1 week along with other symptoms like sore throat, myalgia, pleuritic chest pain, cough, and pain in various joints as mentioned by the study of Raza^[85].

Humoral manifestations

Humoral hypercalcemia of malignancy: Parathyroid hormone-like hormone (PTHrH) is formed by proliferating bile duct epithelial cells in CCA that further interacts with various growth factors resulting in loops of uncontrolled proliferation^[86]. PTHrH derived from CCA cells is involved in causing humoral hypercalcemia of malignancy (HHM). HHM is evident in approximately 10%-20% of the patients with underlying malignancies^[87]. Almost 80%-90% of them are due to PTHrH. Clinical signs evident in a patient with HHM are changes in mental status, constipation, nausea, abdominal

discomfort, polydipsia, polyuria, and weakness. These are due to the increased concentration of calcium in the body^[88]. There is good evidence that proves hypercalcemia as a paraneoplastic manifestation of CCA. Patients showing symptoms of hypercalcemia with normal PTH levels and an increased PTHrH in the setting of CCA supports that it is paraneoplastic. PTHrH works with tumor growth factor-alpha and tumor necrosis factor-alpha to cause hypercalcemia that is HHM.

A summary of all the paraneoplastic syndromes associated with other systems is explained in [Table 2](#) at the bottom of the review.

Association with other neoplasms

Recent diagnostic studies have indicated the presence of certain serological markers in CCA which are raised in other malignancies too suggesting that CCA might be having a concomitant neoplasm that shares the same paraneoplastic syndrome. For example, erythematous skin rash, associated with CCA, has raised CA 19-9. This serological marker is also linked to certain malignancies like colorectal and pancreatic carcinomas^[53]. Another example is hypercalcemia. It is associated with CCA and also found to be raised in 10%-20% of malignancies like squamous cell carcinoma of the lung, head and neck, esophagus, and skin cancers^[89]. Further detailed studies might reveal the hidden facts of this involvement.

Early detection of occult malignancy

CCA has poor prognosis which has led to equalization of incidence and mortality rates^[90]. Although CCA has a poor prognosis, it has been reported that early screening and diagnosis can lead to decreased mortality rates. All the manifestations explained above showed how treating CCA resulted in resolution of symptoms. Thus any patient with the above paraneoplastic manifestation and clinical symptoms should always be screened for cholangiocarcinoma as this could lead to a better chance of survival.

CONCLUSION

CCA can cause a wide range of paraneoplastic syndromes. The exact mechanism linking them to CCA in most of these syndromes remains undetermined. Hence, no particular treatment modality can be recommended. The best management plan remains the removal of the underlying cause which in this case is CCA. The timely recognition of these syndromes and clinical suspicion can lead to early diagnosis of covert malignancies like CCA. The presence of symptoms can also predict the efficacy of treatment, relapse, or recurrence of the disease. Further studies are pertinent to understand the underlying mechanisms and targeted therapies for these paraneoplastic syndromes.

Table 2 Summary of literature on paraneoplastic syndromes in cholangiocarcinoma associated with other systems

Ref.	Year	Paraneoplastic syndrome	Mediator
Neurological			
Schmidt <i>et al</i> ^[57]	2016	Limbic encephalopathy	ANNA-1, Anti-Ma2, Anti-PCA-2/anti-Tr, Anti-VGKC
Bruhnding <i>et al</i> ^[60]	2016	Paraneoplastic cerebellar degeneration	Anti-Yo antibodies
Renal			
Normand <i>et al</i> ^[62]	2017	Glomerulonephritis	
Hematological			
Solans-Laque <i>et al</i> ^[65]	2008	Vasculitis	
Blum <i>et al</i> ^[67]	2016	Trousseau syndrome	
Jang <i>et al</i> ^[68]	2006	Trousseau syndrome	
Samadian <i>et al</i> ^[72]	1999	Anti-phospholipid Antibody Syndrome	
Ham <i>et al</i> ^[74]	2015	Paraneoplastic leukemoid reaction	
Multisystem			
Hatzis <i>et al</i> ^[78]	1998	Polyarteritis nodosa	p-ANCA
Raza <i>et al</i> ^[85]	2013	Adult-onset still disease	
Humoral			
Erdinc <i>et al</i> ^[88]	2019	Humoral hypercalcemia of malignancy	PTHrP

TGF- α : Tissue growth factor- α ; G-CSF: Granulocyte-colony stimulating factor; IL: Interleukin; EGF- α : Epidermal growth factor- α ; ANNA: Antineuronal nuclear antibody; PCA: Purkinje cell antibody; VGKC: Voltage-gated potassium channel; ANCA: Antineutrophil cytoplasmic antibody; PTHrP: Parathyroid hormone-related protein.

REFERENCES

- Bergquist A, von Seth E. Epidemiology of cholangiocarcinoma. *Best Pract Res Clin Gastroenterol* 2015; **29**: 221-232 [PMID: 25966423 DOI: 10.1016/j.bpg.2015.02.003]
- Blechacz B, Komuta M, Roskams T, Gores GJ. Clinical diagnosis and staging of cholangiocarcinoma. *Nat Rev Gastroenterol Hepatol* 2011; **8**: 512-522 [PMID: 21808282 DOI: 10.1038/nrgastro.2011.131]
- Razumilava N, Gores GJ. Cholangiocarcinoma. *Lancet* 2014; **383**: 2168-2179 [PMID: 24581682 DOI: 10.1016/S0140-6736(13)61903-0]
- Florio AA, Ferlay J, Znaor A, Ruggieri D, Alvarez CS, Laversanne M, Bray F, McGlynn KA, Petrick JL. Global trends in intrahepatic and extrahepatic cholangiocarcinoma incidence from 1993 to 2012. *Cancer* 2020; **126**: 2666-2678 [PMID: 32129902 DOI: 10.1002/cncr.32803]
- Saha SK, Zhu AX, Fuchs CS, Brooks GA. Forty-Year Trends in Cholangiocarcinoma Incidence in the U.S.: Intrahepatic Disease on the Rise. *Oncologist* 2016; **21**: 594-599 [PMID: 27000463 DOI: 10.1634/theoncologist.2015-0446]
- Kirstein MM, Vogel A. Epidemiology and Risk Factors of Cholangiocarcinoma. *Visc Med* 2016; **32**: 395-400 [PMID: 28229073 DOI: 10.1159/000453013]
- Everhart JE, Ruhl CE. Burden of digestive diseases in the United States Part III: Liver, biliary tract, and pancreas. *Gastroenterology* 2009; **136**: 1134-1144 [PMID: 19245868 DOI: 10.1053/j.gastro.2009.02.038]
- Darnell RB, Posner JB. Paraneoplastic syndromes involving the nervous system. *N Engl J Med* 2003; **349**: 1543-1554 [PMID: 14561798 DOI: 10.1056/NEJMr023009]
- Sökmen M, Demirsoy H, Ersoy O, Gökdemir G, Akbayir N, Karaca C, Ozdil K, Kesici B, Calişkan C, Yılmaz B. Paraneoplastic porphyria cutanea tarda associated with cholangiocarcinoma: case report. *Turk J Gastroenterol* 2007; **18**: 200-205 [PMID: 17891697]
- Foster MT, Moxon CA, Weir E, Sinha A. Dermal sinus tracts. *BMJ* 2019; **366**: 15202 [PMID: 31484633 DOI: 10.1136/bmj.15202]
- Meegada S, Eisen R, Coons G, Verma R. Intrahepatic Cholangiocarcinoma Associated with High Procalcitonin, Hypercalcemia, Polycythemia and Leukocytosis. *Cureus* 2020; **12**: e6587 [PMID: 32051799 DOI: 10.7759/cureus.6587]
- Lee JK, Chuang MJ, Lu CC, Hao LJ, Yang CY, Han TM, Lam HC. Parathyroid hormone and parathyroid hormone related protein assays in the investigation of hypercalcemic patients in hospital in a Chinese population. *J Endocrinol Invest* 1997; **20**: 404-409 [PMID: 9309539 DOI: 10.1007/BF03347992]
- Aizawa M, Koshiyama H, Inoue D, Fukunaga Y, Katakami H, Miki M, Mizuno Y. Postoperative aggravation of hypercalcemia-leukocytosis syndrome in a case of squamous cell type cholangiocarcinoma. *Intern Med* 1997; **36**: 232 [PMID: 9144021 DOI: 10.2169/internalmedicine.36.232]
- Davis JM, Sadasivan R, Dwyer T, Van Veldhuizen P. Case report: cholangiocarcinoma and hypercalcemia.

- Am J Med Sci* 1994; **307**: 350-352 [PMID: 8172229 DOI: 10.1097/0000441-199405000-00008]
- 15 **Ravnborg L**, Thomsen K. Acanthosis nigricans and bile duct malignancy. *Acta Derm Venereol* 1993; **73**: 378-379 [PMID: 7904407 DOI: 10.2340/0001555573378379]
 - 16 **Kozłowski LM**, Nigra TP. Esophageal acanthosis nigricans in association with adenocarcinoma from an unknown primary site. *J Am Acad Dermatol* 1992; **26**: 348-351 [PMID: 1569256 DOI: 10.1016/0190-9622(92)70053-j]
 - 17 **Cohen PR**, Kurzrock R. Malignancy-associated tripe palms. *J Am Acad Dermatol* 1992; **27**: 271-272 [PMID: 1430373 DOI: 10.1016/s0190-9622(08)80741-4]
 - 18 **Ellis DL**, Kafka SP, Chow JC, Nanney LB, Inman WH, McCadden ME, King LE Jr. Melanoma, growth factors, acanthosis nigricans, the sign of Leser-Trélat, and multiple acrochordons. A possible role for alpha-transforming growth factor in cutaneous paraneoplastic syndromes. *N Engl J Med* 1987; **317**: 1582-1587 [PMID: 2825016 DOI: 10.1056/NEJM198712173172506]
 - 19 **Suchonwanit P**, McMichael AJ. Alopecia in Association with Malignancy: A Review. *Am J Clin Dermatol* 2018; **19**: 853-865 [PMID: 30088232 DOI: 10.1007/s40257-018-0378-1]
 - 20 **Antoniou E**, Paraskeva P, Smyrnis A, Konstantopoulos K. Alopecia: a common paraneoplastic manifestation of cholangiocarcinoma in humans and animals. *BMJ Case Rep* 2012; **2012**: bcr2012006217 [PMID: 22717934 DOI: 10.1136/bcr-2012-006217]
 - 21 **El-Serag HB**, Mason AC. Rising incidence of hepatocellular carcinoma in the United States. *N Engl J Med* 1999; **340**: 745-750 [PMID: 10072408 DOI: 10.1056/NEJM199903113401001]
 - 22 **Knowles BP**, Corcoran NM, Usatoff V. Reading the signs: occult metastatic cholangiocarcinoma detected by full-body screening in dermatomyositis. *ANZ J Surg* 2007; **77**: 1026-1027 [PMID: 17931277 DOI: 10.1111/j.1445-2197.2007.04307.x]
 - 23 **Suh KJ**, Park JK, Cho S, Park H, Baek HW, Lee K, Lee DS, Lee KH. Dermatomyositis in a Patient with Cholangiocarcinoma Detected by an [(18)F]-Fluorodeoxyglucose Positron Emission Tomography-Computed Tomography Scan. *Cancer Res Treat* 2016; **48**: 848-852 [PMID: 25797574 DOI: 10.4143/crt.2014.310]
 - 24 **Al-Nahhas A**, Jawad AS. PET/CT imaging in inflammatory myopathies. *Ann N Y Acad Sci* 2011; **1228**: 39-45 [PMID: 21718321 DOI: 10.1111/j.1749-6632.2011.06016.x]
 - 25 **Kaji K**, Fujimoto M, Hasegawa M, Kondo M, Saito Y, Komura K, Matsushita T, Orito H, Hamaguchi Y, Yanaba K, Itoh M, Asano Y, Seishima M, Ogawa F, Sato S, Takehara K. Identification of a novel autoantibody reactive with 155 and 140 kDa nuclear proteins in patients with dermatomyositis: an association with malignancy. *Rheumatology (Oxford)* 2007; **46**: 25-28 [PMID: 16728436 DOI: 10.1093/rheumatology/ke1161]
 - 26 **Yasuda J**, Onda S, Shiozaki H, Gocho T, Shiba H, Yanaga K. A Successfully Treated Case of Intrahepatic Cholangiocarcinoma with Exacerbation of Dermatomyositis. *Case Rep Gastroenterol* 2018; **12**: 622-628 [PMID: 30483040 DOI: 10.1159/000493185]
 - 27 **Sotoodian B**, Mahmood MN, Salopek TG. Clinical and Dermoscopic Features of Pigmented Disseminated Superficial Actinic Porokeratosis: Case Report and Literature Review. *J Cutan Med Surg* 2018; **22**: 229-231 [PMID: 28922948 DOI: 10.1177/1203475417733465]
 - 28 **Lee HW**, Oh SH, Choi JC, Chang SE, Lee MW, Choi JH, Moon KC, Koh JK. Disseminated superficial porokeratosis in a patient with cholangiocarcinoma. *J Am Acad Dermatol* 2006; **54**: S56-S58 [PMID: 16428001 DOI: 10.1016/j.jaad.2005.07.035]
 - 29 **Cannavó SP**, Borgia F, Adamo B, Guarneri B. Simultaneous development and parallel course of disseminated superficial porokeratosis and ovarian cancer: Coincidental association or true paraneoplastic syndrome? *J Am Acad Dermatol* 2008; **58**: 657-660 [PMID: 18258333 DOI: 10.1016/j.jaad.2007.12.030]
 - 30 **de Roos MA**, de Bock GH, de Vries J, van der Veet B, Wesseling J. p53 overexpression is a predictor of local recurrence after treatment for both in situ and invasive ductal carcinoma of the breast. *J Surg Res* 2007; **140**: 109-114 [PMID: 17291532 DOI: 10.1016/j.jss.2006.10.045]
 - 31 **Shen CS**, Tabata K, Matsuki M, Goto T, Yokochi T, Yamanishi K. Premature apoptosis of keratinocytes and the dysregulation of keratinization in porokeratosis. *Br J Dermatol* 2002; **147**: 498-502 [PMID: 12207590 DOI: 10.1046/j.1365-2133.2002.04853.x]
 - 32 **Lee HJ**, Jang YH. Necrolytic migratory erythema. *CMAJ* 2019; **191**: E286 [PMID: 30858185 DOI: 10.1503/cmaj.180913]
 - 33 **Chiyomaru K**, Takai T, Ohashi A, Nishigori C. Necrolytic migratory erythema with cholangiocarcinoma: Pseudoglucagonoma syndrome. *Eur J Dermatol* 2010; **20**: 238-239 [PMID: 20133217 DOI: 10.1684/ejd.2010.0878]
 - 34 **Blechacz B**, Gores GJ. Cholangiocarcinoma: advances in pathogenesis, diagnosis, and treatment. *Hepatology* 2008; **48**: 308-321 [PMID: 18536057 DOI: 10.1002/hep.22310]
 - 35 **Pavlović MD**, Karadaglić DM, Kandolf LO, Mijusković ZP. Persistent erythema multiforme: a report of three cases. *J Eur Acad Dermatol Venereol* 2001; **15**: 54-58 [PMID: 11451326 DOI: 10.1046/j.1468-3083.2001.00185.x]
 - 36 **Tzovaras V**, Liberopoulos EN, Zioga A, Pavlidis N, Elisaf M. Persistent erythema multiforme in a patient with extrahepatic cholangiocarcinoma. *Oncology* 2007; **73**: 127-129 [PMID: 18337625 DOI: 10.1159/000121001]
 - 37 **Kurzrock R**, Cohen PR. Cutaneous paraneoplastic syndromes in solid tumors. *Am J Med* 1995; **99**: 662-671 [PMID: 7503090 DOI: 10.1016/s0002-9343(99)80254-x]
 - 38 **Cohen PR**, Holder WR, Tucker SB, Kono S, Kurzrock R. Sweet syndrome in patients with solid tumors. *Cancer* 1993; **72**: 2723-2731 [PMID: 8402496 DOI: 10.1002/1097-0142(19931101)72:9<2723::aid-cnrc2820720933>3.0.co;2-f]
 - 39 **Shinojima Y**, Toma Y, Terui T. Sweet syndrome associated with intrahepatic cholangiocarcinoma producing granulocyte colony-stimulating factor. *Br J Dermatol* 2006; **155**: 1103-1104 [PMID: 17034566 DOI: 10.1111/j.1365-2133.2006.07521.x]
 - 40 **Reuss-Borst MA**, Müller CA, Waller HD. The possible role of G-CSF in the pathogenesis of Sweet's syndrome. *Leuk Lymphoma* 1994; **15**: 261-264 [PMID: 7532508 DOI: 10.3109/10428199409049722]
 - 41 **Prevost-Blank PL**, Shwyder TA. Sweet's syndrome secondary to granulocyte colony-stimulating factor. *J*

- Am Acad Dermatol* 1996; **35**: 995-997 [PMID: 8959967 DOI: 10.1016/s0190-9622(96)90132-2]
- 42 **Sato K**, Terada K, Sugiyama T, Sugiyama T, Masuda H, Kakinuma H, Kato T. Granulocyte colony-stimulating factor produced by bladder carcinoma of a patient with leukemoid reaction did not affect proliferation of the tumor cells. *J Urol* 1994; **151**: 1687-1690 [PMID: 7514696 DOI: 10.1016/s0022-5347(17)35345-4]
- 43 **Karabulut AA**, Sahin S, Sahin M, Ekşioğlu M, Ustün H. Paraneoplastic acrokeratosis of Bazex (Bazex's syndrome): report of a female case associated with cholangiocarcinoma and review of the published work. *J Dermatol* 2006; **33**: 850-854 [PMID: 17169088 DOI: 10.1111/j.1346-8138.2006.00194.x]
- 44 **Obasi OE**, Garg SK. Bazex paraneoplastic acrokeratosis in prostate carcinoma. *Br J Dermatol* 1987; **117**: 647-651 [PMID: 3689683 DOI: 10.1111/j.1365-2133.1987.tb07498.x]
- 45 **Buxtorf K**, Hübscher E, Panizzon R. Bazex syndrome. *Dermatology* 2001; **202**: 350-352 [PMID: 11455158 DOI: 10.1159/000051680]
- 46 **Stone SP**, Buescher LS. Life-threatening paraneoplastic cutaneous syndromes. *Clin Dermatol* 2005; **23**: 301-306 [PMID: 15896545 DOI: 10.1016/j.clindermatol.2004.06.011]
- 47 **Liau MM**, Long V, Yang SS. Erythema gyratum repens: a paraneoplastic eruption. *BMJ Case Rep* 2016; **2016**: bcr2016214665 [PMID: 26951443 DOI: 10.1136/bcr-2016-214665]
- 48 **Bar-Ilan E**, Gat A, Sprecher E, Zeeli T. Paraneoplastic pityriasis rubra pilaris: case report and literature review. *Clin Exp Dermatol* 2017; **42**: 54-57 [PMID: 27896859 DOI: 10.1111/ced.13009]
- 49 **Pelosof LC**, Gerber DE. Paraneoplastic syndromes: an approach to diagnosis and treatment. *Mayo Clin Proc* 2010; **85**: 838-854 [PMID: 20810794 DOI: 10.4065/mcp.2010.0099]
- 50 **Mendes GB**, Zanetti G, Marchiori E. Leser-Trélat Sign Secondary to Thymic Carcinoma. *Arch Bronconeumol* 2018; **54**: 286-287 [PMID: 29103671 DOI: 10.1016/j.arbres.2017.08.020]
- 51 **Morgenthau A**, Almudaires A. Klatskin's cholangiocarcinoma presenting with the sign of Leser-Trélat. *BMJ Case Rep* 2019; **12**: e232507 [PMID: 31666258 DOI: 10.1136/bcr-2019-232507]
- 52 **Heaphy MR Jr**, Millns JL, Schroeter AL. The sign of Leser-Trélat in a case of adenocarcinoma of the lung. *J Am Acad Dermatol* 2000; **43**: 386-390 [PMID: 10901731 DOI: 10.1067/mjd.2000.104967]
- 53 **Opneja A**, Mahajan S, Kapoor S, Marur S, Yang SH, Manno R. Unusual Paraneoplastic Presentation of Cholangiocarcinoma. *Case Rep Med* 2015; **2015**: 806835 [PMID: 26495003 DOI: 10.1155/2015/806835]
- 54 **Trüeb RF**, Trüeb RM. Kutane Paraneoplasie als immunologisches Phänomen am Beispiel des paraneoplastischen subakuten kutanen Lupus erythematodes [Cutaneous paraneoplastic syndrome as an immunologic phenomenon exemplified by paraneoplastic subacute cutaneous lupus erythematosus]. *Praxis (Bern 1994)* 1999; **88**: 1803-1810 [PMID: 10584550]
- 55 **Mahawish K**, Teinert L, Cavanagh K, Brennan J. Limbic encephalitis. *BMJ Case Rep* 2014; **2014**: bcr2014204591 [PMID: 24891487 DOI: 10.1136/bcr-2014-204591]
- 56 **Rocamora R**, Becerra JL, Fossas P, Gomez M, Vivanco-Hidalgo RM, Mauri JA, Molins A. Pilocomotor seizures: an autonomic semiology of limbic encephalitis? *Seizure* 2014; **23**: 670-673 [PMID: 24890932 DOI: 10.1016/j.seizure.2014.04.013]
- 57 **Schmidt SL**, Schmidt JJ, Tolentino JC, Ferreira CG, de Almeida SA, Alvarenga RP, Simoes EN, Schmidt GJ, Canedo NH, Chimelli L. Cholangiocarcinoma associated with limbic encephalitis and early cerebral abnormalities detected by 2-deoxy-2-[fluorine-18]fluoro-D-glucose integrated with computed tomography-positron emission tomography: a case report. *J Med Case Rep* 2016; **10**: 200 [PMID: 27439460 DOI: 10.1186/s13256-016-0989-1]
- 58 **Hoffmann LA**, Jarius S, Pellkofer HL, Schueller M, Krumbholz M, Koenig F, Johannis W, la Fougere C, Newman T, Vincent A, Voltz R. Anti-Ma and anti-Ta associated paraneoplastic neurological syndromes: 22 newly diagnosed patients and review of previous cases. *J Neurol Neurosurg Psychiatry* 2008; **79**: 767-773 [PMID: 18223018 DOI: 10.1136/jnnp.2007.118588]
- 59 **Chatzikonstantinou A**, Szabo K, Ottomeyer C, Kern R, Hennerici MG. Successive affection of bilateral temporomesial structures in a case of non-paraneoplastic limbic encephalitis demonstrated by serial MRI and FDG-PET. *J Neurol* 2009; **256**: 1753-1755 [PMID: 19434439 DOI: 10.1007/s00415-009-5165-8]
- 60 **Bruhnding A**, Notch D, Beard A. Anti-Yo positive paraneoplastic cerebellar degeneration in the setting of cholangiocarcinoma. *J Clin Neurosci* 2017; **36**: 71-72 [PMID: 27816258 DOI: 10.1016/j.jocn.2016.10.034]
- 61 **Darnell RB**. Onconeural antigens and the paraneoplastic neurologic disorders: at the intersection of cancer, immunity, and the brain. *Proc Natl Acad Sci USA* 1996; **93**: 4529-4536 [PMID: 8643438 DOI: 10.1073/pnas.93.10.4529]
- 62 **Normand G**, Jolivot A, Rabeyrin M, Hervieu V, Valette PJ, Scoazec JY, Gougou JM, Juillard L, Dumortier J. Paraneoplastic fibrillary glomerulonephritis associated with intrahepatic cholangiocarcinoma: When diagnosis of a rare kidney disease leads to successful hepatic cancer treatment. *Clin Res Hepatol Gastroenterol* 2017; **41**: e8-e11 [PMID: 27542513 DOI: 10.1016/j.clinre.2016.07.002]
- 63 **García-Porrúa C**, González-Gay MA. Cutaneous vasculitis as a paraneoplastic syndrome in adults. *Arthritis Rheum* 1998; **41**: 1133-1135 [PMID: 9627026 DOI: 10.1002/1529-0131(199806)41:6<1133::AID-ART23>3.0.CO;2-S]
- 64 **Fortin PR**. Vasculitides associated with malignancy. *Curr Opin Rheumatol* 1996; **8**: 30-33 [PMID: 8867536 DOI: 10.1097/00002281-199601000-00005]
- 65 **Solans-Laqué R**, Bosch-Gil JA, Pérez-Bocanegra C, Selva-O'Callaghan A, Simeón-Aznar CP, Vilardell-Tarres M. Paraneoplastic vasculitis in patients with solid tumors: report of 15 cases. *J Rheumatol* 2008; **35**: 294-304 [PMID: 18085729]
- 66 **Tasi SH**, Juan CJ, Dai MS, Kao WY. Trousseau's syndrome related to adenocarcinoma of the colon and cholangiocarcinoma. *Eur J Neurol* 2004; **11**: 493-496 [PMID: 15257690 DOI: 10.1111/j.1468-1331.2004.00814.x]
- 67 **Blum MF**, Ma VY, Betbadal AM, Bonomo RA, Raju RR, Packer CD. Trousseau's Syndrome in Cholangiocarcinoma: The Risk of Making the Diagnosis. *Clin Med Res* 2016; **14**: 53-59 [PMID: 26847482 DOI: 10.3121/cm.2015.1304]
- 68 **Jang JW**, Yeo CD, Kim JD, Bae SH, Choi JY, Jung ES, Rha SE, Byun JY, Yoon SK. Trousseau's syndrome

- in association with cholangiocarcinoma: positive tests for coagulation factors and anticardiolipin antibody. *J Korean Med Sci* 2006; **21**: 155-159 [PMID: 16479083 DOI: 10.3346/jkms.2006.21.1.155]
- 69 **Varki A.** Trousseau's syndrome: multiple definitions and multiple mechanisms. *Blood* 2007; **110**: 1723-1729 [PMID: 17496204 DOI: 10.1182/blood-2006-10-053736]
- 70 **Walsh-McMonagle D,** Green D. Low-molecular-weight heparin in the management of Trousseau's syndrome. *Cancer* 1997; **80**: 649-655 [PMID: 9264346 DOI: 10.1002/(SICI)1097-0142(19970815)80:4<649::AID-CNCR1>3.0.CO;2-I]
- 71 **Mouas H,** Lortholary O, Eclache V, Leroux G, Casassus P, Guillemin L, Raphaël M. Antiphospholipid syndrome during acute monocytic leukaemia. *Eur J Haematol* 1994; **53**: 59-60 [PMID: 8062901 DOI: 10.1111/j.1600-0609.1994.tb00183.x]
- 72 **Samadian S,** Estcourt L. Recurrent thrombo-embolic episodes: the association of cholangiocarcinoma with antiphospholipid syndrome. *Postgrad Med J* 1999; **75**: 45-46 [PMID: 10396590 DOI: 10.1136/pgmj.75.879.45]
- 73 **Li C,** Li G, Miao R, Lu X, Zhong S, Sang X, Mao Y, Zhao H. Primary liver cancer presenting as pyogenic liver abscess: characteristics, diagnosis, and management. *J Surg Oncol* 2012; **105**: 687-691 [PMID: 21952992 DOI: 10.1002/jso.22103]
- 74 **Ham H,** Kim HY, Seo KJ, Lee SL, Kim CW. Cholangiocarcinoma with a paraneoplastic leukemoid reaction mimicking a pyogenic liver abscess. *Korean J Intern Med* 2015; **30**: 110-113 [PMID: 25589843 DOI: 10.3904/kjim.2015.30.1.110]
- 75 **Sohda T,** Shiga H, Nakane H, Watanabe H, Takeshita M, Sakisaka S. Cholangiocellular carcinoma that produced both granulocyte-colony-stimulating factor and parathyroid hormone-related protein. *Int J Clin Oncol* 2006; **11**: 246-249 [PMID: 16850133 DOI: 10.1007/s10147-006-0560-y]
- 76 **Karadag O,** Jayne DJ. Polyarteritis nodosa revisited: a review of historical approaches, subphenotypes and a research agenda. *Clin Exp Rheumatol* 2018; **36** Suppl 111: 135-142 [PMID: 29465365]
- 77 **Sánchez-Guerrero J,** Gutiérrez-Ureña S, Vidaller A, Reyes E, Iglesias A, Alarcón-Segovia D. Vasculitis as a paraneoplastic syndrome. Report of 11 cases and review of the literature. *J Rheumatol* 1990; **17**: 1458-1462 [PMID: 2273485]
- 78 **Hatzis GS,** Papachristodoulou A, Delladetsima IK, Moutsopoulos HM. Polyarteritis nodosa associated with cholangiocarcinoma. *Lupus* 1998; **7**: 301-306 [PMID: 9696133 DOI: 10.1191/096120398678920217]
- 79 **Maestri A,** Malacarne P, Santini A. Henoch-Schönlein syndrome associated with breast cancer. A case report. *Angiology* 1995; **46**: 625-627 [PMID: 7618766 DOI: 10.1177/000331979504600710]
- 80 **Naschitz JE,** Yeshurun D, Abrahamson J. Arterial occlusive disease in occult cancer. *Am Heart J* 1992; **124**: 738-745 [PMID: 1514502 DOI: 10.1016/0002-8703(92)90285-4]
- 81 **Navarro JF,** Quereda C, Rivera M, Navarro FJ, Ortuño J. Anti-neutrophil cytoplasmic antibody-associated paraneoplastic vasculitis. *Postgrad Med J* 1994; **70**: 373-375 [PMID: 8016013 DOI: 10.1136/pgmj.70.823.373]
- 82 **Steinberg W.** The clinical utility of the CA 19-9 tumor-associated antigen. *Am J Gastroenterol* 1990; **85**: 350-355 [PMID: 2183589]
- 83 **Kulp-Shorten CL,** Rhodes RH, Peterson H, Callen JP. Cutaneous vasculitis associated with pheochromocytoma. *Arthritis Rheum* 1990; **33**: 1852-1856 [PMID: 2261006 DOI: 10.1002/art.1780331215]
- 84 **Wu N,** Li Q, Gu CX, Ahmed T, Yao XP. Paraneoplastic syndrome mimicking adult-onset Still's disease caused by advanced lung cancer: a case report. *BMC Cancer* 2011; **11**: 487 [PMID: 22085873 DOI: 10.1186/1471-2407-11-487]
- 85 **Raza A,** Houk L, Yousaf W, Smiley D, Coberly L. Unusual para-neoplastic manifestation of cholangiocarcinoma. *J Gastrointest Cancer* 2013; **44**: 228-230 [PMID: 22903498 DOI: 10.1007/s12029-012-9430-1]
- 86 **Alvaro D,** Gigliozzi A, Attili AF. Regulation and deregulation of cholangiocyte proliferation. *J Hepatol* 2000; **33**: 333-340 [PMID: 10952254 DOI: 10.1016/s0168-8278(00)80377-3]
- 87 **Birkeland KI,** Gallefoss F, Olsson S, Haug E. Primary hyperparathyroidism or hypercalcaemia of malignancy? *Scand J Clin Lab Invest* 1992; **52**: 347-349 [PMID: 1439519 DOI: 10.1080/00365519209088368]
- 88 **Erdinc B,** Ramachandran P, Yadav R, Sahni S, Joseph G. Cholangiocarcinoma Presenting as Humoral Hypercalcemia of Malignancy: A Case Report and Literature Review. *Cureus* 2019; **11**: e6481 [PMID: 32025405 DOI: 10.7759/cureus.6481]
- 89 **O'Malley JT,** Schoppe C, Husain S, Grossman ME. Squamous Cell Carcinoma (Marjolin's Ulcer) Arising in a Sacral Decubitus Ulcer Resulting in Humoral Hypercalcemia of Malignancy. *Case Rep Med* 2014; **2014**: 715809 [PMID: 25197285 DOI: 10.1155/2014/715809]
- 90 **Khan SA,** Toledano MB, Taylor-Robinson SD. Epidemiology, risk factors, and pathogenesis of cholangiocarcinoma. *HPB (Oxford)* 2008; **10**: 77-82 [PMID: 18773060 DOI: 10.1080/13651820801992641]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

