

World Journal of *Gastroenterology*

World J Gastroenterol 2020 October 28; 26(40): 6111-6303



REVIEW

- 6111 Cirrhotic portal hypertension: From pathophysiology to novel therapeutics
Gunarathne LS, Rajapaksha H, Shackel N, Angus PW, Herath CB
- 6141 Role of gut microbiota *via* the gut-liver-brain axis in digestive diseases
Ding JH, Jin Z, Yang XX, Lou J, Shan WX, Hu YX, Du Q, Liao QS, Xie R, Xu JY

MINIREVIEWS

- 6163 Imaging-based algorithmic approach to gallbladder wall thickening
Gupta P, Marodia Y, Bansal A, Kalra N, Kumar-M P, Sharma V, Dutta U, Sandhu MS
- 6182 Overlooked risk for needle tract seeding following endoscopic ultrasound-guided minimally invasive tissue acquisition
Gao RY, Wu BH, Shen XY, Peng TL, Li DF, Wei C, Yu ZC, Luo MH, Xiong F, Wang LS, Yao J
- 6195 Prevalence, diagnosis, and treatment of primary hepatic gastrointestinal stromal tumors
Qian XH, Yan YC, Gao BQ, Wang WL

ORIGINAL ARTICLE**Basic Study**

- 6207 Prediction of clinically actionable genetic alterations from colorectal cancer histopathology images using deep learning
Jang HJ, Lee A, Kang J, Song IH, Lee SH
- 6224 New strain of *Pediococcus pentosaceus* alleviates ethanol-induced liver injury by modulating the gut microbiota and short-chain fatty acid metabolism
Jiang XW, Li YT, Ye JZ, Lv LX, Yang LY, Bian XY, Wu WR, Wu JJ, Shi D, Wang Q, Fang DQ, Wang KC, Wang QQ, Lu YM, Xie JJ, Li LJ

Retrospective Cohort Study

- 6241 Predicting cholecystocholedochal fistulas in patients with Mirizzi syndrome undergoing endoscopic retrograde cholangiopancreatography
Wu CH, Liu NJ, Yeh CN, Wang SY, Jan YY

Retrospective Study

- 6250 Novel endoscopic papillectomy for reducing postoperative adverse events (with videos)
Jiang L, Ling-Hu EQ, Chai NL, Li W, Cai FC, Li MY, Guo X, Meng JY, Wang XD, Tang P, Zhu J, Du H, Wang HB

Clinical Trials Study

- 6260 Pediatric bowel preparation: Sodium picosulfate, magnesium oxide, citric acid *vs* polyethylene glycol, a randomized trial
Cuffari C, Ciciora SL, Ando M, Boules M, Croffie JM

SYSTEMATIC REVIEWS

- 6270 Insufficient etiological workup of COVID-19-associated acute pancreatitis: A systematic review
Juhász MF, Ocskay K, Kiss S, Hegyi P, Párnicsky A

META-ANALYSIS

- 6279 Diagnostic efficacy of the Japan Narrow-band-imaging Expert Team and Pit pattern classifications for colorectal lesions: A meta-analysis
Zhang Y, Chen HY, Zhou XL, Pan WS, Zhou XX, Pan HH

CASE REPORT

- 6295 Compromised therapeutic value of pediatric liver transplantation in ethylmalonic encephalopathy: A case report
Zhou GP, Qu W, Zhu ZJ, Sun LY, Wei L, Zeng ZG, Liu Y

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Dr. Rakesh Kochhar is a Professor in and Head of the Department of Gastroenterology at the Postgraduate Institute of Medical Education and Research, Chandigarh, India. Having completed his MBBS from J N Medical College (Aligarh, India), he performed postgraduate training in Medicine and undertook a fellowship in Gastroenterology at the Postgraduate Institute of Medical Education and Research. He then joined his alma mater as faculty and became a professor in 2003. Widely travelled, he worked as a World Health Organization Fellow at Mayo Clinic (Rochester, USA) and St Bart's Hospital (London, UK). His ongoing research interests include corrosive injuries to the gastrointestinal tract, stricture dilatation, acute pancreatitis, and celiac disease. He has served as President of the Society of Gastrointestinal Endoscopy of India and is currently the President-elect of the Indian Society of Gastroenterology. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (*WJG*, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJG* mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2020 edition of Journal Citation Report® cites the 2019 impact factor (IF) for *WJG* as 3.665; IF without journal self cites: 3.534; 5-year IF: 4.048; Ranking: 35 among 88 journals in gastroenterology and hepatology; and Quartile category: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ji-Hong Liu; Production Department Director: Yun-Xiaojuan Wu; Editorial Office Director: Ze-Mao Gong.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, Subrata Ghosh

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

October 28, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Diagnostic efficacy of the Japan Narrow-band-imaging Expert Team and Pit pattern classifications for colorectal lesions: A meta-analysis

Yu Zhang, Hui-Yan Chen, Xiao-Lu Zhou, Wen-Sheng Pan, Xin-Xin Zhou, Hang-Hai Pan

ORCID number: Yu Zhang 0000-0003-1387-3962; Hui-Yan Chen 0000-0002-6483-5002; Xiao-Lu Zhou 0000-0002-6771-3027; Wen-Sheng Pan 0000-0002-2347-1695; Xin-Xin Zhou 0000-0003-0183-6400; Hang-Hai Pan 0000-0002-7366-8269.

Author contributions: Zhang Y contributed to the conception and design of the study, and critical revision; Pan HH and Zhou XX contributed to the acquisition of data and data analysis; Pan HH, Zhou XX, Pan WS, Chen HY and Zhou XL contributed to the interpretation of data; Pan HH and Zhou XX drafted the article; Pan WS, Chen HY and Zhou XL revised the manuscript; all authors contributed to the final approval of the manuscript.

Supported by the Natural Science Foundation of Zhejiang Province, No. LQ20H160061; and Medical Health Science and Technology Project of Zhejiang Provincial Health Commission, No. 2018255969

Conflict-of-interest statement: No potential conflicts of interest exist.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Yu Zhang, Wen-Sheng Pan, Hang-Hai Pan, Department of Gastroenterology, Zhejiang Provincial People's Hospital, People's Hospital of Hangzhou Medical College, Hangzhou 310014, Zhejiang Province, China

Hui-Yan Chen, School of Laboratory Medicine and Life Sciences, Wenzhou Medical University, Wenzhou 325035, Zhejiang Province, China

Xiao-Lu Zhou, Department of Medical College, The Medical College of Qingdao University, Qingdao 266071, Shandong Province, China

Xin-Xin Zhou, Department of Gastroenterology, The First Affiliated Hospital, Zhejiang University, Hangzhou 310003, Zhejiang Province, China

Corresponding author: Hang-Hai Pan, MD, Attending Doctor, Department of Gastroenterology, Zhejiang Provincial People's Hospital, People's Hospital of Hangzhou Medical College, No. 158 Shangtang Road, Hangzhou 310014, Zhejiang Province, China. panhanghai@163.com

Abstract

BACKGROUND

Pit pattern classification using magnifying chromoendoscopy is the established method for diagnosing colorectal lesions. The Japan Narrow-band-imaging (NBI) Expert Team (JNET) classification is a novel NBI magnifying endoscopic classification that focuses on the vessel, and surface patterns.

AIM

To determine the diagnostic efficacy of each category of the JNET and Pit pattern classifications for colorectal lesions.

METHODS

A systematic literature search was performed using PubMed, Embase, the Cochrane Library, and Web of Science databases. The pooled sensitivity, specificity, diagnostic odds ratio, and area under the summary receiver operating characteristic curve of each category of the JNET and Pit pattern classifications were calculated.

RESULTS

A total of 19227 colorectal lesions in 31 studies were included. The diagnostic performance of the JNET classification was equivalent to the Pit pattern classification in each corresponding category. The pooled sensitivity, specificity,

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: June 21, 2020

Peer-review started: June 21, 2020

First decision: August 21, 2020

Revised: August 26, 2020

Accepted: September 23, 2020

Article in press: September 23, 2020

Published online: October 28, 2020

P-Reviewer: Chiba H

S-Editor: Yan JP

L-Editor: Webster JR

P-Editor: Liu JH



and area under the curve (AUC) for each category of the JNET classification were as follows: 0.73 (95%CI: 0.55-0.85), 0.99 (95%CI: 0.97-1.00), and 0.97 (95%CI: 0.95-0.98), respectively, for Type 1; 0.88 (95%CI: 0.78-0.94), 0.72 (95%CI: 0.64-0.79), and 0.84 (95%CI: 0.81-0.87), respectively, for Type 2A; 0.56 (95%CI: 0.47-0.64), 0.91 (95%CI: 0.79-0.96), and 0.72 (95%CI: 0.68-0.76), respectively, for Type 2B; 0.51 (95%CI: 0.42-0.61), 1.00 (95%CI: 1.00-1.00), and 0.90 (95%CI: 0.87-0.93), respectively, for Type 3.

CONCLUSION

This meta-analysis suggests that the diagnostic efficacy of the JNET classification may be equivalent to that of the Pit pattern classification. However, due to its simpler and clearer clinical application, the JNET classification should be promoted for the classification of colorectal lesions, and to guide the treatment strategy.

Key Words: Colorectal neoplasms; Colonoscopy; Chromoendoscopy; Japan Narrow-band-imaging Expert Team; Pit pattern; Meta-analysis

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This systematic review and meta-analysis examined the diagnostic efficacy of each category of the Japan Narrow-band imaging Expert Team (JNET) and Pit pattern classifications for colorectal lesions. The pooled analyses of 19227 colorectal lesions from 31 studies were performed. The sensitivity, specificity, diagnostic odds ratio, and area under the summary receiver operating characteristic curve of each category of the JNET and Pit pattern classifications were calculated. The results suggest that the diagnostic efficacy of the JNET classification may be equivalent to that of the Pit pattern classification. However, due to its simpler and clearer clinical application, the JNET classification should be promoted for classification of colorectal lesions, and to guide the treatment strategy.

Citation: Zhang Y, Chen HY, Zhou XL, Pan WS, Zhou XX, Pan HH. Diagnostic efficacy of the Japan Narrow-band-imaging Expert Team and Pit pattern classifications for colorectal lesions: A meta-analysis. *World J Gastroenterol* 2020; 26(40): 6279-6294

URL: <https://www.wjgnet.com/1007-9327/full/v26/i40/6279.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v26.i40.6279>

INTRODUCTION

Colorectal cancer (CRC) is one of the most common malignant tumors in the world^[1]. Accurate identification and treatment of early CRC have significantly reduced its incidence and mortality^[2]. However, early CRC is generally asymptomatic, and colonoscopy allows the direct visual inspection of the intestinal tract and same-session detection, biopsy, and subsequent removal of lesions. Endoscopic evaluation of colorectal lesions is important to guide the selection of an appropriate treatment. The ideal endoscopic management of colorectal lesions involves two steps. First, any mucosal lesions need to be detected. Second, the lesions need to be characterized based on histological characteristics assessed *via* endoscopic evaluation; which forms the basis of the endoscopic judgment on whether the lesions need to be resected. In general, there is a consensus that low-grade dysplasias (LGDs), high-grade dysplasias (HGDs), and superficial submucosal invasive (SM-s) carcinomas are considered appropriate for endoscopic resection, while observation is recommended for hyperplastic polyps (HPs), in contrast, surgery is recommended for deep submucosal invasive (SM-d) carcinomas^[3].

Kudo *et al*^[4] first proposed the "Pit patterns" (the opening shape of a colorectal crypt) based on the classification of colorectal lesions *via* magnifying endoscopy, and indigo carmine dye contrast^[4]. Pit pattern diagnosis (Type I-V) is clinically significant as it can differentiate between neoplasia and non-neoplasia, characterize the degree of histological atypia in a tumor, and reveal the invasion depth of early carcinoma.

Usually, standard magnification requires the use of chromoagents (*e.g.*, indigo carmine, crystal violet, or methylene blue) to clarify the pit structures in these diagnostic procedures. However, with the emergence of electronic staining endoscopic equipment, chemical staining endoscopy has gradually been replaced. In addition, simpler and more convenient procedures are desirable for magnifying procedures. It has been suggested that narrow-band-imaging (NBI) technology is as effective as chromoendoscopy in differentiating the gross type of colorectal lesions^[5]. The Colon Tumor NBI Interest Group proposed the NBI International Colorectal Endoscopic (NICE) classification in 2010, which is based on non-magnifying NBI observations^[6]. However, the NICE classification cannot distinguish between benign adenoma and superficial mucosal carcinoma, therefore, it plays a limited role in guiding endoscopic treatment strategies. As more endoscopic devices are being equipped with a magnifying function, NBI combined with magnifying endoscopy is increasingly being used, which further improves the diagnostic efficiency, and plays an important role in estimating the invasion depth of the lesion. In order to better guide the endoscopic treatment strategy, in 2014, the Japan NBI Expert Team (JNET) proposed the JNET classification as a universal NBI magnifying endoscopic classification^[7]. The JNET classification focuses on vessel and surface patterns to diagnose colorectal lesions as Types 1, 2A and B, 3.

Recently, several studies have proposed that the JNET classification of colorectal lesions *via* NBI magnifying endoscopy is a useful and objective tool for differentiating the gross type of colorectal lesions. However, differences in the diagnostic performance of the JNET classification have not been reported according to each gross type. Although several investigators now accept the application of the JNET classification for colorectal lesions, to what extent we can trust the results of the JNET classification, and whether the Pit pattern classification can be replaced by the JNET classification are aspects that remain unclear.

Few studies have compared the diagnostic efficacy of the JNET and Pit pattern classifications for each gross type of lesion, and determined the correlation between endoscopic features and pathological findings, and, to our knowledge, there are no meta-analyses on this topic. Accordingly, we performed a systematic review and meta-analysis to analyze the data on existing magnifying endoscopy trials using the JNET classification and the Pit pattern classification for characterization of colorectal lesions, and, to obtain a statistically convincing conclusion on the diagnostic accuracy, and practicability of these two comparable methods.

MATERIALS AND METHODS

Literature search

We performed a systematic literature search of articles in PubMed, Web of Science, Embase and the Cochrane Library (January 1990 to May 2020) containing quantitative data, and manually searched the reference lists of retrieved articles. The following search terms were used: "Japan Narrow-Band Imaging Expert Team", "Japan NBI Expert Team", "JNET", "Pit pattern", "Kudo's classification", "Colorectal Neoplasm", "Neoplasm, Colorectal", "Colorectal Carcinoma", "Carcinoma, Colorectal", "Carcinomas, Colorectal", "Colorectal Carcinomas", "Colorectal Cancer", "Cancer, Colorectal", "Cancers, Colorectal", "Colorectal Cancers", "Colorectal Tumors", "Colorectal Tumor", "Tumor, Colorectal", "Tumors, Colorectal", "Neoplasms, Colorectal", "Colon polyps", "Colorectal polyps", and "Colorectal lesions". The queries used are displayed in the supplementary materials. All similar possible word variations were also searched. The attained records were retrieved and managed with EndNote X 9.0 (Bld 10136, Thomson Reuters, New York, NY, United States).

Study selection

Studies were included when all of the following conditions were met: (1) Studies in which all participants received the JNET or Pit pattern classification for colorectal lesions diagnosed *via* endoscopy; (2) Histological diagnosis was chosen as the gold standard; and (3) Studies in which sufficient data were reported to calculate true positive (TP), false positive (FP), false negative (FN), and true negative (TN) results. In addition, we included only the latest-published article when the same population was reported in more than one article. However, conference papers and duplicate published studies that fulfilled the above two criteria were excluded. This meta-analysis was performed in compliance with the preferred reporting items for systematic reviews and meta-analyses statement (PRISMA)^[8].

Data extraction

Data from the included studies were extracted and cross-checked by 2 authors independently (Zhang Y and Pan HH). If there was a discrepancy in their opinions, it was discussed with other authors to achieve a consistent result. The extracted data included the name of the first author, year of publication, demographics of the population, type of endoscope, number of included patients, number of colorectal lesions examined, design of the study, and the type of classification for colorectal lesions. The number of TP, FP, TN, and FN results for the JNET, and Pit pattern classifications were the main statistics extracted from the studies. We computed sensitivity [TP/(TP + FN)] and specificity [TN/(TN + FP)] for each technique separately.

Quality assessment

The quality of the included studies was independently assessed by 2 authors independently (Zhou XX and Chen HY) using the quality assessment of diagnostic accuracy studies II (QUADAS-II) tool^[9].

Statistical analysis

The original data from each study (TP, FP, TN and FN) were integrated through meta-analysis, and the pooled sensitivity, specificity, positive likelihood ratio, and diagnostic odds ratio (DOR) of both the JNET and Pit pattern classifications were calculated by the DerSimonian Laird random effects model^[10]. The heterogeneity of pooled sensitivity and specificity was calculated using the I^2 statistic, and a high degree of heterogeneity was set at $I^2 > 50\%$ ^[11]. Mose's constant linear model was used to perform the summary receiver operating characteristic curve^[12]. Cochrane's Q test was used to evaluate the accuracy of the JNET and Pit pattern classifications in the diagnosis of colorectal lesions. When heterogeneity was present, the Spearman correlation coefficient, and the P value or heterogeneity ratio caused by the threshold effect were calculated. Meta-regression was conducted to explore the existing source of heterogeneity. Publication bias was assessed using the Deeks' Funnel Plot Asymmetry Test. The t -test was used to compare the statistical significance of the area under the curve (AUC) and pooled sensitivity, with the significance set at $P < 0.05$. The statistical software used for the diagnostic accuracy test was Stat 15.1. Revman5.3 was used to evaluate the quality of the included studies.

RESULTS**Study selection**

A total of 1146 articles were initially searched (351 in Web of Science, 241 in PubMed, 44 in Cochrane Library, and 510 in Embase), and 31 studies^[5,13-42] with a total of 14674 patients were ultimately included in this meta-analysis. 1114 studies were excluded of which 306 studies were duplicate references, 442 studies were excluded based on title and abstract, and 70 studies were excluded after full-text review. A detailed PRISMA flow diagram is shown in [Figure 1](#).

Description of studies and qualitative analysis

All studies used the JNET or Pit pattern classifications as the diagnostic criteria for colorectal lesions examined *via* endoscopy. The JNET classified colorectal lesions into four categories: Type 1 is a hyperplastic polyp (HP)/sessile serrated lesion (SSL), Type 2A is a LGD/adenoma, Type 2B is a HGD/M-SM-s and Type 3 is a SM-d^[5]. The Pit pattern classification has five corresponding categories: Type I is normal mucosa, Type II is hyperplasia/SSL, Types III_L/IV are LGD/adenomas, Types III_S/V_{L-L}, and V_{H-H}/V_N are associated with HGD/M-SM-s lesions, and SM-d lesions, respectively^[13]. The detailed interpretation criteria are shown in [Table 1](#). Our study was restricted to only cross-sectional outcomes such as sensitivity and specificity, and the screening tests were compared to the reference standard (histopathological diagnosis). General information on the included studies is presented in [Table 2](#). Of the 31 studies^[5,13-42], 11 were retrospective^[5,13-22], and 20 were prospective^[23-42]. Seven studies^[14-20] used the JNET classification alone, 21 studies^[22-42] used the Pit pattern classification alone, and only 3 studies^[5,13,21] used both the JNET and Pit pattern classifications in the same population. Thirteen studies^[5,13-16,18-21,24,28,35,41] used narrow-band imaging magnifying endoscopy (NBI-ME), 13 studies used magnifying chromoendoscopy^[22,25-27,29,30,32-35,37,38,42], and the other studies used BLI-magnification^[17,23] ($n = 2$), NBI^[31,39] ($n = 2$), and

Table 1 Detailed tentative criteria for interpretation of the Japan Narrow-band-imaging Expert Team and Pit pattern classifications compared with histologic diagnosis

Interpretation	JNET	Pit pattern	Histological diagnosis	Therapeutic strategy
Non-neoplastic		I	Normal mucosa	
	Type 1	II	Hyperplastic/SSL	Follow-up observation
Neoplastic	Type 2A	III _L + IV	Adenoma/LGD	Endoscopic resection (polypectomy/EMR)
	Type 2B	III _S + V _{L-L}	HGD/M-SM-s cancer	Endoscopic resection (ESD)
	Type 3	V _N + V _{L-H}	SM-d cancer	Surgery

JNET: Japan Narrow-band-imaging Expert Team; EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection.

chromoendoscopy^[36,40] ($n = 2$). The QUADAS-II quality assessment for each study is presented in [Figure 2](#).

Diagnostic efficacy

In total, 19227 colorectal lesions in 14674 patients were identified in the 31 included studies. [Table 3](#) summarizes the pooled sensitivity, specificity, DOR, and AUC in each category of the JNET and Pit pattern classifications corresponding to the pathological results.

Ten studies^[5,13-21] involving 13479 colorectal lesions reported the diagnostic efficacy of the JNET classification. The pooled values for each category were as follows. Type 1 (non-neoplastic): Sensitivity, 0.73 (95%CI: 0.55-0.85); specificity, 0.99 (95%CI: 0.97-1.00) ([Figure 3A](#)); DOR, 245 (95%CI: 64-936); and AUC, 0.97 (95%CI: 0.95-0.98) ([Figure 4A](#)). Type 2A: sensitivity, 0.88 (95%CI: 0.78-0.94); specificity, 0.72 (95%CI: 0.64-0.79) ([Figure 8A](#)); DOR, 19 (95%CI: 11-33); and AUC, 0.84 (95%CI: 0.81-0.87) ([Figure 4B](#)). Type 2B: sensitivity, 0.56 (95%CI: 0.47-0.64); specificity, 0.91 (95%CI: 0.79-0.96) ([Figure 5A](#)); DOR, 13 (95%CI: 7-24); and AUC, 0.72 (95%CI: 0.68-0.76) ([Figure 4C](#)); Type 3: sensitivity, 0.51 (95%CI: 0.42-0.61); specificity, 1.00 (95%CI: 1.00-1.00) ([Figure 6A](#)); DOR, 801 (95%CI: 267-2398); and AUC, 0.90 (95%CI: 0.87-0.93) ([Figure 4D](#)).

Twenty-one studies^[5,13,21-42] involving 6150 colorectal lesions reported the diagnostic value of the Pit pattern classification. The pooled values for each category were as follows. Types I + II (non-neoplastic): sensitivity, 0.86 (95%CI: 0.81-0.90); specificity, 0.94 (95%CI: 0.90-0.96) ([Figure 7](#)); DOR, 88 (95%CI: 48-156); and AUC, 0.95 (95%CI: 0.93-0.97) ([Figure 9A](#)). Types III_L + IV: sensitivity, 0.80 (95%CI: 0.67-0.89); specificity, 0.80 (95%CI: 0.74-0.86) ([Figure 8B](#)); DOR, 17 (95%CI: 8-34); and AUC, 0.87 (95%CI: 0.83-0.89) ([Figure 9B](#)). Types III_S + V_{L-L}: sensitivity, 0.45 (95%CI: 0.23-0.69); specificity, 0.88 (95%CI: 0.75-0.94) ([Figure 5B](#)); DOR, 6 (95%CI: 1-26); and AUC, 0.79 (95%CI: 0.75-0.82) ([Figure 9C](#)). Types V_N + V_{L-H}: sensitivity, 0.73 (95%CI: 0.55-0.85); specificity, 0.99 (95%CI: 0.98-1.00) ([Figure 6B](#)); DOR, 449 (95%CI: 93-2182); and AUC, 0.98 (95%CI: 0.97-0.99) ([Figure 9D](#)).

The Student's *t*-test was used to compare the sensitivities, specificities, and AUCs of each corresponding category of these two classifications. A statistically significant difference was found in sensitivity between JNET Type 3, and Pit pattern Types V_N + V_{L-H} ($P < 0.05$), however, no significant difference was found in specificity, and AUC. The results showed that no significant differences was found between the remaining categories among sensitivities, specificities, and AUCs: JNET Type 1 (non-neoplastic) *vs* Pit pattern Types I + II (non-neoplastic), JNET Type 2A *vs* Pit pattern Types III_L + IV, and JNET Type 2B *vs* Pit pattern Types III_S + V_{L-L}.

Heterogeneity analysis

Significant heterogeneity existed among the included studies. Estimation of Spearman's correlation coefficient (coef) and the *P*-value for each category of the JNET and Pit pattern classifications are shown in [Table 4](#). Significant differences were noted for JNET Type 1 (non-neoplastic) (coef. = 0.14, $P = 0.02$), JNET Type 3 (coef. = -0.17, $P = 0.03$), and Pit pattern Types I + II (non-neoplastic) (coef. = -0.12, $P = 0.02$), while for the remaining subtypes of these two classifications, there were no significant differences. These results indicated the threshold effect existed in JNET Types 1, 3, and Pit pattern Types I + II. Additionally, for the non-threshold effect, we performed meta-regression analysis, including the population in the study (Asian or non-Asian), design of the

Table 2 General characteristics of the included studies

Ref.	Year	Country	Study design	Patients (n)	Lesions (n)	Type of endoscopy	Type of classification	Category
Shibagaki <i>et al</i> ^[13]	2020	Japan	Retrospective	432	718	A-NBI-ME	JNET/Pit pattern	1/2A/2B/3; I, II, III/IV, V _L , V _{L-H} , V _N
Kobayashi <i>et al</i> ^[5]	2019	Japan	Retrospective	750	1402	NBI-ME	JNET/Pit pattern	1/2A/2B/3; I, II, III _L , III _S , IV, IV _H , V _{L-L} , V _{L-H} , V _N
Sumimoto <i>et al</i> ^[14]	2017	Japan	Retrospective	1901	2933	NBI-ME	JNET	1/2A/2B/3
Murano <i>et al</i> ^[15]	2019	Japan	Retrospective	680	1472	NBI-ME	JNET	1/2A/2B/3
Komeda <i>et al</i> ^[16]	2017	Japan	Retrospective	199	199	NBI-ME	JNET	1/2A/2B/3
Suzuki <i>et al</i> ^[17]	2019	Japan	Retrospective	145	185	BLI-ME	JNET	1/2A/2B/3
Hirata <i>et al</i> ^[18]	2019	Japan	Retrospective	6138	6138	NBI-ME	JNET	1/2A/2B/3
Sakamoto <i>et al</i> ^[19]	2019	Japan	Retrospective	246	246	NBI-ME	JNET	1/2A/2B/3
Gonai <i>et al</i> ^[20]	2020	Japan	Retrospective	153	169	NBI-ME	JNET	2A/2B/3
Kawasaki <i>et al</i> ^[21]	2019	Japan	Retrospective	17	17	NBI-ME	JNET/Pit pattern	1/2A/2B/3; III, IV, V _{L-L} , V _{L-H} , V _N
Nakano <i>et al</i> ^[23]	2017	Japan	Prospective	506	799	BLI-ME	Pit pattern	II, III _L , III _S /IV, V _{L-L} , V _{L-H} /V _N
Mirosław <i>et al</i> ^[24]	2015	Poland	Prospective	270	386	WL, NBI-ME	Pit pattern	I/II/III _L /III _S /IV/V
Su <i>et al</i> ^[25]	2004	China	Prospective	230	270	Magnifying chromoendoscopy	Pit pattern	I/II/III _S /III _L /IV/V
Kiesslich <i>et al</i> ^[26]	2001	Germany	Prospective	100	283	Magnifying chromoendoscopy	Pit pattern	I/II/III _S /III _L /IV/V
Kawaguti <i>et al</i> ^[27]	2019	Brazil	Prospective	121	123	Magnifying chromoendoscopy	Pit pattern	II/III _S /III _L /IV/V
East <i>et al</i> ^[28]	2008	United Kingdom	Prospective	62	116	NBI-ME	Pit pattern	I/II/III _S /III _L /IV/V
Liu <i>et al</i> ^[29]	2003	China	Prospective	948	954	Magnifying chromoendoscopy	Pit pattern	I/II/III _L /III _S /IV/V
Kato <i>et al</i> ^[30]	2006	Japan	Prospective	180	210	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Van den Broek <i>et al</i> ^[31]	2011	Netherlands	Prospective	48	153	NBI	Pit pattern	Neoplastic/non-neoplastic
Togashi <i>et al</i> ^[32]	2009	Japan	Prospective	50	107	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Kiesslich <i>et al</i> ^[33]	2003	Germany	Prospective	84	118	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Hurlstone <i>et al</i> ^[34]	2005	United Kingdom	Prospective	350	288	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Tischendorf <i>et al</i> ^[35]	2007	Germany	Prospective	52	100	NBI-ME	Pit pattern	Neoplastic/non-neoplastic
Tischendorf <i>et al</i> ^[35]	2007	Germany	Prospective	47	100	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Su <i>et al</i> ^[36]	2006	China	Prospective	79	110	Chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Dos Santos <i>et al</i> ^[37]	2012	Brazil	Prospective	69	120	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Dos Santos <i>et al</i> ^[38]	2010	Brazil	Prospective	72	137	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Rogart <i>et al</i> ^[39]	2008	United States	Prospective	131	265	NBI	Pit pattern	Neoplastic/non-neoplastic
Apel <i>et al</i> ^[40]	2006	Germany	Prospective	158	273	Chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic

Van den Broek <i>et al</i> ^[41]	2008	Netherlands	Prospective	50	98	NBI-ME	Pit pattern	Neoplastic/non-neoplastic
Chiu <i>et al</i> ^[42]	2007	China	Prospective	133	180	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic
Liu <i>et al</i> ^[22]	2008	China	Retrospective	223	451	Magnifying chromoendoscopy	Pit pattern	Neoplastic/non-neoplastic

JNET: Japan Narrow-band-imaging Expert Team; NBI: Narrow band imaging; NBI-ME: NBI magnifying endoscopy; WL: White light; BLI: Blue laser imaging; A-NBIME: NBIME with acetic acid enhancement.

Table 3 Summary of the results of each category for the Japan Narrow-band-imaging Expert Team classification and the Pit pattern classification corresponding to histological diagnosis in the included studies

JNET	Sensitivity	Specificity	DOR	AUC	Pit pattern	Sensitivity	Specificity	DOR	AUC
1	0.73 [0.55, 0.85]	0.99 [0.97, 1.00]	245 [64, 936]	0.97 [0.95, 0.98]	II	0.76 [0.62, 0.86]	0.96 [0.88, 0.98]	68 [15, 309]	0.92 [0.90, 0.94]
2A	0.88 [0.78, 0.94]	0.72 [0.64, 0.79]	19 [11, 33]	0.84 [0.81, 0.87]	III _L + IV	0.80 [0.67, 0.89]	0.80 [0.74, 0.86]	17 [8, 34]	0.87 [0.83, 0.89]
2B	0.56 [0.47, 0.64]	0.91 [0.79, 0.96]	13 [7, 24]	0.72 [0.68, 0.76]	III _S + V _{L-L}	0.45 [0.23, 0.69]	0.88 [0.75, 0.94]	6 [1, 26]	0.79 [0.75, 0.82]
3	0.51 [0.42, 0.61]	1.00 [1.00, 1.00]	801 [267, 2398]	0.90 [0.87, 0.93]	V _N + V _{I-H}	0.73 [0.55, 0.85]	0.99 [0.98, 1.00]	449 [93, 2182]	0.98 [0.97, 0.99]
Non-neoplastic	0.73 [0.55, 0.85]	0.99 [0.97, 1.00]	245 [64, 936]	0.97 [0.95, 0.98]	Non-neoplastic	0.86 [0.81, 0.90]	0.94 [0.90, 0.96]	88 [48, 156]	0.95 [0.93, 0.97]

JNET: Japan Narrow-band-imaging Expert Team; NBI: Narrow-band-imaging; DOR: Diagnostic odds ratio; AUC: Area under the curve.

Table 4 Summary of the results of Spearman's correlation coefficient of each category for the Japan Narrow-band-imaging Expert Team classification and the Pit pattern classification corresponding to histological diagnosis in the included studies

JNET (type)	Coef	P value	Pit pattern (type)	Coef	P value
			Non-neoplastic	-0.12	0.02
Type 1 (Non-neoplastic)	0.14	0.02	II	0.45	0.20
Type 2A	-0.70	0.49	III _L + IV	-0.24	0.06
Type 2B	-1.00	1.00	III _S + V _{L-L}	0.41	0.17
Type 3	-0.17	0.03	V _N + V _{I-H}	1.00	1.00

JNET: Japan Narrow-band-imaging Expert Team.

study (retrospective or prospective), patient sample size (≥ 100 or < 100), QUADAS-2 score (≥ 7 or < 7), publication year (before, or after 2014), as well as the type of endoscopy as covariates. The sources of potential heterogeneity in the sensitivity and specificity were detected by univariate regression analysis, and the results are shown in Table 5. Deeks' Funnel Plot Asymmetry Test revealed no publication bias in each category of these two classifications (Table 6).

DISCUSSION

To best of our knowledge, no meta-analysis has been reported that systematically compared the diagnostic efficacy of detailed histologic characteristics and interobserver diagnostic agreements between the JNET and Pit pattern classifications. The present meta-analysis compared and evaluated the diagnostic outcomes of each category of these two classifications corresponding to the histological diagnosis. Our results revealed that the diagnostic performance of the JNET classification is

Table 5 Summary of the results of meta-regression analysis of each category for the Japan Narrow-band-imaging Expert Team classification and the Pit pattern classification corresponding to histological diagnosis in the included studies

Classification	Type	P value						
		Prodesign	Sampsize	Qscore	Endoscopy	Category	Country	Pubyear
JNET	Type 1	0.26		0.63	0.06			
	Type 2A		0.88	0.58	0.76	0.00		
	Type 2B							
	Type 3		0.57	0.12	0.19			
Pit pattern	II	0.52		0.00	0.04		0.35	0.00
	III _L + IV	0.63	0.77	0.41	0.33	0.77	0.00	0.09
	III _S + V _{L-L}	0.35	0.35	0.66	0.35	0.35	0.21	0.02
	V _N + V _{L-H}							
	Non-neoplastic	0.49	0.60	0.04	0.52	0.05	0.03	0.00

JNET: Japan Narrow-band-imaging Expert Team.

Table 6 Summary of the results of Deek’s test for publication bias of each category for the Japan Narrow-band-imaging Expert Team classification and the Pit pattern classification corresponding to histological diagnosis in the included studies

JNET (type)	n of study	Pubbias (P value)	Pit pattern (type)	No. of study	Pubbias (P value)
Type 1	8	0.26	II	5	0.54
Type 2A	10	0.62	III _L + IV	10	0.41
Type 2B	9	0.52	III _S + V _{L-L}	6	0.09
Type 3	9	0.50	V _N + V _{L-H}	8	0.91
Non-neoplastic	8	0.26	Non-neoplastic	23	0.13

JNET: Japan Narrow-band-imaging Expert Team.

equivalent to the Pit pattern classification in each corresponding category. We also proposed a treatment strategy for colorectal lesions using the JNET classification with its confidence level.

Endoscopic diagnosis *via* magnification colonoscopy has been reported to provide high diagnostic accuracy and improve the prognosis of colorectal lesions. The Pit pattern classification is the most frequently used criteria for the accurate diagnosis of colorectal neoplasms. To date, several trials^[5,13-21] have evaluated the efficacy of the JNET classification, which provides useful criteria for optical-histologic diagnoses of colorectal lesions. Whereas, JNET Types 1 and 2A correspond to Pit pattern Types I + II, and Type III_L + IV, respectively; JNET Type 2B and 3 correspond to Pit pattern Types III_S + V_{L-L} and V_{L-H} + V_N, respectively. However, there are some differences between the JNET classification and Pit pattern classification. The Pit pattern classification is only based on the surface structure of lesions, while the JNET classification is based on the surface structure combined with the microvascular structure of lesions. In addition, the JNET classification is more concise in terms of guiding appropriate treatment strategies. According to our proposed treatment strategy, most JNET Type 1 lesions are HPs, which generally do not require resection and could be followed up by endoscopy. However, whether using the JNET classification or Pit pattern classification, it is difficult to distinguish between HP and SSL with endoscopy. Endoscopic resection is also recommended if the colorectal lesion is large or tends to enlarge obviously with endoscopy follow-up. Type 2A lesions could be resected by polypectomy or endoscopic mucosal resection (EMR). Type 2B lesions should be resected en bloc by endoscopic submucosal dissection (ESD) to obtain a precise histologic diagnosis concerning the invasion depth and determine endoscopic curability. If the lesion is relatively small, EMR is also recommended. Lastly, surgical resection is recommended for Type 3 lesions. However, few




Figure 1 Study identification, inclusion and exclusion for meta-analysis.

studies^[5,13-21] have been published that have used the JNET classification for treatment of colorectal lesions in practice.

The JNET Type 1 is considered to have as high diagnostic efficacy as the Pit pattern Type II, moreover, no significant difference was found between these categories in this meta-analysis, which implies that the magnification technology could accurately distinguish non-neoplastic from neoplastic lesions and guide clinicians to formulate the appropriate treatment.

In this study, JNET Type 2B had a relatively low pooled sensitivity and AUC of 0.56 and 0.72, respectively. Interestingly, similar results were also obtained for Pit pattern Types III_s + V_{L-L}, with a sensitivity of 0.45 and AUC of 0.79. Through data analysis, it was found that, in these two categories, the evaluation regarding the dysplasia of some lesions was too low, or the depth of invasion was too shallow. Firstly, one reason might be that a large lesion often contained several pathological features, for example, a lesion contained two or more histological features of HGD, SM-s carcinoma, and SM-d carcinoma at the same time. Most of the surface structures showed the features of HGD, or SM-s carcinoma, while, only a focal, or the deep part of these lesions showed the features of SM-d carcinoma; therefore, the lesions were identified as Type 2B. Secondly, due to the large size, or the special location of the lesions, the endoscopist might be unable to observe the full picture of these lesions, which might be one reason for the low sensitivity. In addition, due to the spontaneous or contact bleeding of some large lesions, the blood might be attached to the surface of the lesions and then affect the judgment of pathological Types. Therefore, for large lesions, special location lesions and pedicled lesions, endoscopists are required to observe carefully, to obtain the whole picture of the lesions, and make a comprehensive judgement, which might be helpful in improving sensitivity.

Generally, SM-d carcinomas correspond to JNET Type 3. Our meta-analysis showed that JNET Type 3 has a lower sensitivity than that of Pit pattern Type V_N + V_{L-H}. Some polypoid advanced lesions could have a slightly less irregular NBI appearance than Type 3 because the surface microvillous structure persisted, which might be classified into JNET Type 2B. Thus, it has been proposed that Type 2B can be divided into two subtypes, Type 2B-low, and Type 2B-high^[14]. The classification subtypes help to optimize the choice of treatment strategies, which also indicates that the JNET classification may need to be updated and optimized by experts to further improve the sensitivity of diagnosis. In addition, two studies^[5,14] suggested that the endoscopist needed to perform an additional Pit pattern diagnosis using chromoendoscopy to differentiate Type 2B from Type 3, which might help to improve sensitivity; however, this requires further validation. In terms of specificity, compared with the Pit pattern




Figure 2 Quality assessment of the included studies.




Figure 3 Forest plots of pooled sensitivity and specificity. A: Japan Narrow-band-imaging Expert Team type 1; B: Pit pattern II.

classification, the overall JNET types were slightly higher, which was attributed to the evaluation of vessel structure by NBI-ME.

There are several limitations to this meta-analysis. First, the high degree of statistical heterogeneity with a high I^2 value could not be avoided. The quality of endoscopic images, type of endoscopy, size of the population, year of publication, and experience of the endoscopists (expert or non-expert) possibly affected the heterogeneity of the included studies. For the Pit pattern classification, chemical staining magnifications were used in most of the included studies, but several studies used electronic staining magnification, and a few studies also used non-magnifying technology, which might have affected the results. Second, previous studies of the JNET classification were all retrospective single-center studies and the included populations were all Japanese. This indicates a potential need for large-scale prospective multi-center validation




Figure 4 Summary receiver operating characteristic of the Japan Narrow-band-imaging Expert Team classification. To diagnose colorectal lesions with the corresponding 95% confidence region. A: Type 1; B: Type 2A; C: Type 2B and Type 3. SROC: Summary receiver operating characteristic.




Figure 5 Forest plots of pooled sensitivity and specificity. A: Japan Narrow-band-imaging Expert Team type 2B; B: Pit pattern III_S + V_{LL}.

studies of the JNET classification in the future. Additionally, it is better to compare the JNET classification with the Pit pattern classification in the same endoscopic and histopathologic center.

CONCLUSION

In conclusion, this meta-analysis has shown that the diagnostic efficacy of the JNET




Figure 6 Forest plots of pooled sensitivity and specificity. A: Japan Narrow-band-imaging Expert Team type 3; B: Pit pattern V_N + V_{I-H}.




Figure 7 Forest plots of pooled sensitivity and specificity of non-neoplastic lesions by Pit pattern.

classification is equivalent to that of the Pit pattern classification as both classifications are divided into four major categories according to similar histopathology. The sensitivity of JNET Type 2B can be further improved by differentiating subtypes and combining it with the Pit pattern classification. Due to its simpler and clearer application, it is easier to guide the choice of treatment strategy, which suggests that we can promote the application of the JNET classification for colorectal lesions in the clinic. However, future prospective multi-center studies with a uniform endoscopic and histopathology protocol are required to validate our results.




Figure 8 Forest plots of pooled sensitivity and specificity. A: Japan Narrow-band-imaging Expert Team type 2A; B: Pit pattern III_L + IV.




Figure 9 Summary receiver operating characteristic of Pit pattern classification. To diagnose colorectal lesions with the corresponding 95% confidence region. A: II; B: III_L + IV; C: III_S + V_{HL} and V_N + V_{HH}. SROC: Summary receiver operating characteristic.

ARTICLE HIGHLIGHTS

Research background

The Pit pattern classification using magnifying chromoendoscopy is the established and traditional choice for diagnosing colorectal lesions. The Japan Narrow-band imaging (NBI) Expert Team (JNET) classification is a novel NBI magnifying endoscopic classification, which focuses on vessel and surface patterns to diagnose colorectal lesions. However, the comparative diagnostic efficacy of the JNET classification is inconclusive.

Research motivation

Several studies have proposed that the JNET classification of colorectal lesions *via* NBI magnifying endoscopy is a useful and objective tool for differentiating the gross type of colorectal lesions. However, to what extent we can trust the results of the JNET classification, and whether the Pit pattern classification can be replaced by the JNET classification are aspects that remain unclear.

Research objectives

To perform a systematic review and meta-analysis on the diagnostic efficacy of each category of the JNET and Pit pattern classifications for colorectal lesions.

Research methods

A systematic literature search was performed using PubMed, Embase, the Cochrane Library and Web of Science. The pooled sensitivity, specificity, diagnostic odds ratio, and area under the curve (AUC) of each category of the JNET and Pit pattern classifications were calculated.

Research results

A total of 19227 colorectal lesions in 31 studies were included. The diagnostic performance of the JNET classification was equivalent to the Pit pattern classification in each corresponding category. The pooled sensitivity, specificity, and AUC for each category of the JNET classification were as follows: 0.73 (95%CI: 0.55-0.85), 0.99 (95%CI: 0.97-1.00), and 0.97 (95%CI: 0.95-0.98), respectively, for Type 1; 0.88 (95%CI: 0.78-0.94), 0.72 (95%CI: 0.64-0.79), and 0.84 (95%CI: 0.81-0.87), respectively, for Type 2A; 0.56 (95%CI: 0.47-0.64), 0.91 (95%CI: 0.79-0.96), and 0.72 (95%CI: 0.68-0.76), respectively, for Type 2B; 0.51 (95%CI: 0.42-0.61), 1.00 (95%CI: 1.00-1.00), and 0.90 (95%CI: 0.87-0.93), respectively, for Type 3.

Research conclusions

Our study found that the diagnostic efficacy of the JNET classification is equivalent to that of the Pit pattern classification as both classifications are divided into four major categories according to similar histopathology.

Research perspectives

The JNET classification is easier for guiding the choice of treatment strategy, which suggests that we can promote the application of the JNET classification for colorectal lesions in the clinic. However, future prospective multi-center studies with uniform endoscopic and histopathology protocol are required to validate our results.

REFERENCES

- 1 **Bray F**, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; **68**: 394-424 [PMID: 30207593 DOI: 10.3322/caac.21492]
- 2 **Lin JS**, Piper MA, Perdue LA, Rutter CM, Webber EM, O'Connor E, Smith N, Whitlock EP. Screening for Colorectal Cancer: Updated Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA* 2016; **315**: 2576-2594 [PMID: 27305422 DOI: 10.1001/jama.2016.3332]
- 3 **Tanaka S**, Kashida H, Saito Y, Yahagi N, Yamano H, Saito S, Hisabe T, Yao T, Watanabe M, Yoshida M, Kudo SE, Tsuruta O, Sugihara KI, Watanabe T, Saitoh Y, Igarashi M, Toyonaga T, Ajioka Y, Ichinose M, Matsui T, Sugita A, Sugano K, Fujimoto K, Tajiri H. JGES guidelines for colorectal endoscopic submucosal dissection/endoscopic mucosal resection. *Dig Endosc* 2015; **27**: 417-434 [PMID: 25652022 DOI: 10.1111/den.12456]
- 4 **Kudo S**, Tamura S, Nakajima T, Yamano H, Kusaka H, Watanabe H. Diagnosis of colorectal tumorous lesions by magnifying endoscopy. *Gastrointest Endosc* 1996; **44**: 8-14 [PMID: 8836710 DOI: 10.1016/S0016-5107(96)00008-0]

- 10.1016/s0016-5107(96)70222-5]
- 5 **Kobayashi S**, Yamada M, Takamaru H, Sakamoto T, Matsuda T, Sekine S, Igarashi Y, Saito Y. Diagnostic yield of the Japan NBI Expert Team (JNET) classification for endoscopic diagnosis of superficial colorectal neoplasms in a large-scale clinical practice database. *United European Gastroenterol J* 2019; **7**: 914-923 [PMID: 31428416 DOI: 10.1177/2050640619845987]
 - 6 **Tanaka S**, Sano Y. Aim to unify the narrow band imaging (NBI) magnifying classification for colorectal tumors: current status in Japan from a summary of the consensus symposium in the 79th Annual Meeting of the Japan Gastroenterological Endoscopy Society. *Dig Endosc* 2011; **23** Suppl 1: 131-139 [PMID: 21535219 DOI: 10.1111/j.1443-1661.2011.01106.x]
 - 7 **Sano Y**, Tanaka S, Kudo SE, Saito S, Matsuda T, Wada Y, Fujii T, Ikematsu H, Uraoka T, Kobayashi N, Nakamura H, Hotta K, Horimatsu T, Sakamoto N, Fu KI, Tsuruta O, Kawano H, Kashida H, Takeuchi Y, Machida H, Kusaka T, Yoshida N, Hirata I, Terai T, Yamano HO, Kaneko K, Nakajima T, Sakamoto T, Yamaguchi Y, Tamai N, Nakano N, Hayashi N, Oka S, Iwatate M, Ishikawa H, Murakami Y, Yoshida S, Saito Y. Narrow-band imaging (NBI) magnifying endoscopic classification of colorectal tumors proposed by the Japan NBI Expert Team. *Dig Endosc* 2016; **28**: 526-533 [PMID: 26927367 DOI: 10.1111/den.12644]
 - 8 **Liberati A**, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 2009; **62**: e1-34 [PMID: 19631507 DOI: 10.1016/j.jclinepi.2009.06.006]
 - 9 **Whiting PF**, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, Leeflang MM, Sterne JA, Bossuyt PM; QUADAS-2 Group. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011; **155**: 529-536 [PMID: 22007046 DOI: 10.7326/0003-4819-155-8-201110180-00009]
 - 10 **DerSimonian R**, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**: 177-188 [PMID: 3802833 DOI: 10.1016/0197-2456(86)90046-2]
 - 11 **Higgins JPT**, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch V. Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019). Cochrane, 2019. Available from: <https://training.cochrane.org/cochrane-handbook-systematic-reviews-interventions#how-to-access>
 - 12 **Moses LE**, Shapiro D, Littenberg B. Combining independent studies of a diagnostic test into a summary ROC curve: data-analytic approaches and some additional considerations. *Stat Med* 1993; **12**: 1293-1316 [PMID: 8210827 DOI: 10.1002/sim.4780121403]
 - 13 **Shibagaki K**, Ishimura N, Yuki T, Taniguchi H, Aimi M, Kobayashi K, Kotani S, Yazaki T, Yamashita N, Tamagawa Y, Mishiro T, Ishihara S, Yasuda A, Kinshita Y. Magnification endoscopy in combination with acetic acid enhancement and narrow-band imaging for the accurate diagnosis of colonic neoplasms. *Endosc Int Open* 2020; **8**: E488-E497 [PMID: 32258370 DOI: 10.1055/a-1068-2056]
 - 14 **Sumimoto K**, Tanaka S, Shigita K, Hayashi N, Hirano D, Tamaru Y, Ninomiya Y, Oka S, Arihiro K, Shimamoto F, Yoshihara M, Chayama K. Diagnostic performance of Japan NBI Expert Team classification for differentiation among noninvasive, superficially invasive, and deeply invasive colorectal neoplasia. *Gastrointest Endosc* 2017; **86**: 700-709 [PMID: 28257790 DOI: 10.1016/j.gie.2017.02.018]
 - 15 **Murano T**, Ikematsu H, Shinmura K, Ito R, Minamide T, Sato D, Yamamoto Y, Takashima K, Kadota T, Yoda Y, Hori K, Oono Y, Yano T. Endoscopic prediction of advanced histology in colorectal lesions sized <10 mm using the Japan Narrow-band imaging Expert Team classification. *Dig Endosc* 2019; **32**: 785-790 [PMID: 31630442 DOI: 10.1111/den.13566]
 - 16 **Komeda Y**, Kashida H, Sakurai T, Asakuma Y, Tribonias G, Nagai T, Kono M, Minaga K, Takenaka M, Arizumi T, Hagiwara S, Matsui S, Watanabe T, Nishida N, Chikugo T, Chiba Y, Kudo M. Magnifying Narrow Band Imaging (NBI) for the Diagnosis of Localized Colorectal Lesions Using the Japan NBI Expert Team (JNET) Classification. *Oncology* 2017; **93** Suppl 1: 49-54 [PMID: 29258091 DOI: 10.1159/000481230]
 - 17 **Suzuki H**, Yamamura T, Nakamura M, Hsu CM, Su MY, Chen TH, Chiu CT, Hirooka Y, Goto H. An International Study on the Diagnostic Accuracy of the Japan Narrow-Band Imaging Expert Team Classification for Colorectal Polyps Observed with Blue Laser Imaging. *Digestion* 2020; **101**: 339-346 [PMID: 30982036 DOI: 10.1159/000499856]
 - 18 **Hirata D**, Kashida H, Iwatate M, Tochio T, Teramoto A, Sano Y, Kudo M. Effective use of the Japan Narrow Band Imaging Expert Team classification based on diagnostic performance and confidence level. *World J Clin Cases* 2019; **7**: 2658-2665 [PMID: 31616682 DOI: 10.12998/wjcc.v7.i18.2658]
 - 19 **Sakamoto T**, Takamaru H, Sekiguchi M, Yamada M, Matsuda T, Saito Y. Reliability of Japan Narrow-Band Imaging Expert Team Classification for the Diagnosis of Colorectal Neoplasms: A Pilot Study. *Digestion* 2019; **1-6** [PMID: 31394529 DOI: 10.1159/000501960]
 - 20 **Gonai T**, Kawasaki K, Nakamura S, Yanai S, Akasaka R, Sato K, Toya Y, Asakura K, Urushikubo J, Fujita Y, Eizuka M, Uesugi N, Sugai T, Matsumoto T. Microvascular density under magnifying narrow-band imaging endoscopy in colorectal epithelial neoplasms. *Intest Res* 2020; **18**: 107-114 [PMID: 31671929 DOI: 10.5217/ir.2019.00061]
 - 21 **Kawasaki K**, Nakamura S, Esaki M, Kurahara K, Eizuka M, Nuki Y, Kochi S, Fujiwara M, Oshiro Y, Sugai T, Matsumoto T. Clinical usefulness of magnifying colonoscopy for the diagnosis of ulcerative colitis-associated neoplasia. *Dig Endosc* 2019; **31** Suppl 1: 36-42 [PMID: 30994234 DOI: 10.1111/den.13382]
 - 22 **Liu YX**, Huang LY, Bian XP, Cui J, Xu N, Wu CR. Fuji Intelligent Chromo Endoscopy and staining technique for the diagnosis of colon tumor. *Chin Med J (Engl)* 2008; **121**: 977-982 [PMID: 18706244 DOI: 10.3238/arztebl.2008.0421b]
 - 23 **Nakano A**, Hirooka Y, Yamamura T, Watanabe O, Nakamura M, Funasaka K, Ohno E, Kawashima H, Miyahara R, Goto H. Comparison of the diagnostic ability of blue laser imaging magnification versus pit pattern analysis for colorectal polyps. *Endosc Int Open* 2017; **5**: E224-E231 [PMID: 28367494 DOI: 10.1055/s-0043-102400]
 - 24 **Szura M**, Pasternak A, Bucki K, Urbańczyk K, Matyja A. Two-stage optical system for colorectal polyp assessments. *Surg Endosc* 2016; **30**: 204-214 [PMID: 25840894 DOI: 10.1007/s00464-015-4186-x]

- 25 **Su MY**, Ho YP, Chen PC, Chiu CT, Wu CS, Hsu CM, Tung SY. Magnifying endoscopy with indigo carmine contrast for differential diagnosis of neoplastic and nonneoplastic colonic polyps. *Dig Dis Sci* 2004; **49**: 1123-1127 [PMID: [15387332](#) DOI: [10.1023/b:ddas.0000037798.55845.f7](#)]
- 26 **Kiesslich R**, von Bergh M, Hahn M, Hermann G, Jung M. Chromoendoscopy with indigocarmine improves the detection of adenomatous and nonadenomatous lesions in the colon. *Endoscopy* 2001; **33**: 1001-1006 [PMID: [11740641](#) DOI: [10.1055/s-2001-18932](#)]
- 27 **Kawaguti FS**, Franco MC, Martins BC, Segateli V, Marques CFS, Nahas CSR, Pinto RA, Safatle-Ribeiro AV, Ribeiro-Junior U, Nahas SC, Maluf-Filho F. Role of Magnification Chromoendoscopy in the Management of Colorectal Neoplastic Lesions Suspicious for Submucosal Invasion. *Dis Colon Rectum* 2019; **62**: 422-428 [PMID: [30730457](#) DOI: [10.1097/DCR.0000000000001343](#)]
- 28 **East JE**, Suzuki N, Bassett P, Stavrinidis M, Thomas HJ, Guenther T, Tekkis PP, Saunders BP. Narrow band imaging with magnification for the characterization of small and diminutive colonic polyps: pit pattern and vascular pattern intensity. *Endoscopy* 2008; **40**: 811-817 [PMID: [18828077](#) DOI: [10.1055/s-2008-1077586](#)]
- 29 **Liu HH**, Kudo SE, Juch JP. Pit pattern analysis by magnifying chromoendoscopy for the diagnosis of colorectal polyps. *J Formos Med Assoc* 2003; **102**: 178-182 [PMID: [12783135](#) DOI: [10.1007/s11263-006-7834-8](#)]
- 30 **Kato S**, Fu KI, Sano Y, Fujii T, Saito Y, Matsuda T, Koba I, Yoshida S, Fujimori T. Magnifying colonoscopy as a non-biopsy technique for differential diagnosis of non-neoplastic and neoplastic lesions. *World J Gastroenterol* 2006; **12**: 1416-1420 [PMID: [16552812](#) DOI: [10.3748/wjg.v12.i9.1416](#)]
- 31 **van den Broek FJ**, Fockens P, van Eeden S, Stokkers PC, Ponsioen CY, Reitsma JB, Dekker E. Narrow-band imaging versus high-definition endoscopy for the diagnosis of neoplasia in ulcerative colitis. *Endoscopy* 2011; **43**: 108-115 [PMID: [21165822](#) DOI: [10.1055/s-0030-1255956](#)]
- 32 **Togashi K**, Osawa H, Koinuma K, Hayashi Y, Miyata T, Sunada K, Nokubi M, Horie H, Yamamoto H. A comparison of conventional endoscopy, chromoendoscopy, and the optimal-band imaging system for the differentiation of neoplastic and non-neoplastic colonic polyps. *Gastrointest Endosc* 2009; **69**: 734-741 [PMID: [19251019](#) DOI: [10.1016/j.gie.2008.10.063](#)]
- 33 **Kiesslich R**, Fritsch J, Holtmann M, Koehler HH, Stolte M, Kanzler S, Nafe B, Jung M, Galle PR, Neurath MF. Methylene blue-aided chromoendoscopy for the detection of intraepithelial neoplasia and colon cancer in ulcerative colitis. *Gastroenterology* 2003; **124**: 880-888 [PMID: [12671882](#) DOI: [10.1053/gast.2003.50146](#)]
- 34 **Hurlstone DP**, Sanders DS, Lobo AJ, McAlindon ME, Cross SS. Indigo carmine-assisted high-magnification chromoscopic colonoscopy for the detection and characterisation of intraepithelial neoplasia in ulcerative colitis: a prospective evaluation. *Endoscopy* 2005; **37**: 1186-1192 [PMID: [16329015](#) DOI: [10.1055/s-2005-921032](#)]
- 35 **Tischendorf JJ**, Wasmuth HE, Koch A, Hecker H, Trautwein C, Winograd R. Value of magnifying chromoendoscopy and narrow band imaging (NBI) in classifying colorectal polyps: a prospective controlled study. *Endoscopy* 2007; **39**: 1092-1096 [PMID: [18072061](#) DOI: [10.1055/s-2007-966781](#)]
- 36 **Su MY**, Hsu CM, Ho YP, Chen PC, Lin CJ, Chiu CT. Comparative study of conventional colonoscopy, chromoendoscopy, and narrow-band imaging systems in differential diagnosis of neoplastic and nonneoplastic colonic polyps. *Am J Gastroenterol* 2006; **101**: 2711-2716 [PMID: [17227517](#) DOI: [10.1111/j.1572-0241.2006.00932.x](#)]
- 37 **Dos Santos CE**, Malaman D, Lopes CV, Pereira-Lima JC, Parada AA. Digital chromoendoscopy for diagnosis of diminutive colorectal lesions. *Diagn Ther Endosc* 2012; **2012**: 279521 [PMID: [23082070](#) DOI: [10.1155/2012/279521](#)]
- 38 **dos Santos CE**, Lima JC, Lopes CV, Malaman D, Salomão AD, Garcia AC, Teixeira CR. Computerized virtual chromoendoscopy versus indigo carmine chromoendoscopy combined with magnification for diagnosis of small colorectal lesions: a randomized and prospective study. *Eur J Gastroenterol Hepatol* 2010; **22**: 1364-1371 [PMID: [20453654](#) DOI: [10.1097/MEG.0b013e32833a5d63](#)]
- 39 **Rogart JN**, Jain D, Siddiqui UD, Oren T, Lim J, Jamidar P, Aslanian H. Narrow-band imaging without high magnification to differentiate polyps during real-time colonoscopy: improvement with experience. *Gastrointest Endosc* 2008; **68**: 1136-1145 [PMID: [18691708](#) DOI: [10.1016/j.gie.2008.04.035](#)]
- 40 **Apel D**, Jakobs R, Schilling D, Weickert U, Teichmann J, Bohrer MH, Riemann JF. Accuracy of high-resolution chromoendoscopy in prediction of histologic findings in diminutive lesions of the rectosigmoid. *Gastrointest Endosc* 2006; **63**: 824-828 [PMID: [16650546](#) DOI: [10.1016/j.gie.2005.09.013](#)]
- 41 **van den Broek FJ**, Fockens P, van Eeden S, Reitsma JB, Hardwick JC, Stokkers PC, Dekker E. Endoscopic tri-modal imaging for surveillance in ulcerative colitis: randomised comparison of high-resolution endoscopy and autofluorescence imaging for neoplasia detection; and evaluation of narrow-band imaging for classification of lesions. *Gut* 2008; **57**: 1083-1089 [PMID: [18367559](#) DOI: [10.1136/gut.2007.144097](#)]
- 42 **Chiu HM**, Chang CY, Chen CC, Lee YC, Wu MS, Lin JT, Shun CT, Wang HP. A prospective comparative study of narrow-band imaging, chromoendoscopy, and conventional colonoscopy in the diagnosis of colorectal neoplasia. *Gut* 2007; **56**: 373-379 [PMID: [17005766](#) DOI: [10.1136/gut.2006.099614](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

