

1
Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 57870

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Clinical characteristics of perineal endometriosis: A case series

Yan Liang, Duo Zhang, Ling Jiang, Yuan Liu, Jian Zhang

Abstract

BACKGROUND

The prevalence of perineal endometriosis (PEM) is low among women with endometriosis (EM) treated by surgery. It manifests as hard or cystic nodules with pain in the perineal wounds and surrounding areas. Implantation theory is regarded as the main pathogenesis of PEM. There are few clinical researches studying the incidence and clinical characteristics of PEM. This study aims to summarize the clinical data of 14 PEM cases and analyze the factors that may be related to the incubation period and

Match Overview

Match 1 of 1

1

Internet 19 words
crawled on 20-Dec-2019
f6publishing.blob.core.windows.net

<1%



ALL

IMAGES

VIDEOS

6,010,000 Results

Any time ▾

[The clinical characteristics and surgical approach of scar ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6087557>

Scar **endometriosis** can be clinically misdiagnosed as hernia, lipoma, or hematoma. Here we retrospectively analyzed the **clinical** aspects of scar **endometriosis** and surgical approach in **14** patients from a tertiary hospital, who were treated by surgery, between 2012 and 2017. The mean age was 32.71 ± 8.61 years (range: 19–45).

Cited by: 6

Author: Faik Tatli, Orhan Gozeneli, Hacer Uyanik...

Publish Year: 2017

[Abdominal and perineal scar endometriosis: Retrospective ...](#)

<https://www.sciencedirect.com/science/article/pii/S0301211520304292>

Sep 01, 2020 · We confirmed 26 women with AWE from 860 cases with **endometriosis** (3,0 %) and **14** patients with **perineal endometriosis** (1,6 %). The mean age of Group I was $36,5 \pm 3,4$, in contrast with Group II in which mean age was $32,5 \pm 2,9$ respectively (). Table 1 shows the outcomes of baseline pathological and surgical **characteristics** of patients with AWE and **perineum scar endometriosis**.

Author: Michail Matalliotakis, Michail Matalliot...

Publish Year: 2020

[Perineal Endometriosis: A Rarity](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6314099>

A **clinical** diagnosis of episiotomy scar **endometriosis** was considered. On transperineal ultrasonography, a focal hypoechoic lesion with small cystic spaces was noted in the subcutaneous plane of the right **perineal** region [Figure 1], measuring $3.0 \text{ cm} \times 2.8 \text{ cm} \times 3.2 \text{ cm}$, and a possible diagnosis of **perineal endometriosis** stated.

Author: Reddy Ravikanth

Publish Year: 2018

[Abdominal and perineal scar endometriosis: Retrospective ...](#)

<https://www.sciencedirect.com/science/article/abs/pii/S0301211520304292>

All cases with **perineal scar endometriosis** were multiparous and delivered vaginally with episiotomy. 92,8 % of patients presented with cyclical pain and swelling. 3 cases suffered from **perineal endometriosis** combined with pelvic **endometriosis**. There was a recurrence of **perineal endometriosis** in 2 women (14,2 %).

Author: Michail Matalliotakis, Michail Matalliot...

Publish Year: 2020



ALL

IMAGES

VIDEOS

226,000 Results

Any time ▾

Characteristics and long-term outcomes of perineal ...

<https://pubmed.ncbi.nlm.nih.gov/32502046>

Patients were divided into the gonadotropins releasing hormone (GnRH) agonist group and non-GnRH agonist group. The main clinical symptom was **vulvar painful swellings related to menstrual cycles**. Thirty-three patients' lesions (94.29%) were on the episiotomy scar while 1 ...

Author: Yu Liu, Ruyu Pi, Hong Luo, Wei Wang, Xi... **Publish Year:** 2020

Perineal groove in female infants: A case series and ...

<https://onlinelibrary.wiley.com/doi/10.1111/pde.13311>

Oct 11, 2017 · A retrospective study of female infants diagnosed with a **perineal** groove was performed, demographic **characteristics** and **clinical** features were analyzed, and a photographic review was conducted. Results. Five patients with **perineal** groove were observed in our clinic in 2015-16. The mean age was 14 months.

Cited by: 8 **Author:** Maria Garcia-Palacios, Roberto Mendez-Ga...
Publish Year: 2017

The clinical characteristics and surgical approach of scar ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6087557>

Nov 14, 2017 · Clinically, scar endometriosis is **difficult to diagnose**, but **several symptoms commonly occur, including cyclic pain during menstruation [15] and swelling in the pelvic region**. Similarly, most of our patients had been referred to the hospital due to their complaints of ...

Cited by: 6 **Author:** Faik Tatli, Orhan Gozeneli, Hacer Uyanikogl...
Publish Year: 2017

Abdominal and perineal scar endometriosis: Retrospective ...

<https://www.sciencedirect.com/science/article/pii/S0301211520304292>

Sep 01, 2020 · All cases with perineal scar endometriosis were **multiparous and delivered vaginally with an episiotomy**. The majority of women (92,8 %) presented with complaints of cyclical pain and swelling. Finally, 3 patients suffered from perineal combined with pelvic endometriosis. There was a recurrence of perineal endometriosis in 2 women (14,2 %). Comment

Author: Michail Matalliotakis, Michail Matalliotas... **Publish Year:** 2020

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

234,000 Results

Any time ▼

[Characteristics and long-term outcomes of perineal ...](#)

<https://pubmed.ncbi.nlm.nih.gov/32502046>

Patients were divided into the gonadotropins releasing hormone (GnRH) agonist group and non-GnRH agonist group. The main clinical symptom was **vulvar painful swellings related to menstrual cycles**. Thirty-three patients' lesions (94.29%) were on the episiotomy scar while 1 ...

Author: Yu Liu, Ruyu Pi, Hong Luo, Wei Wang, Xi... **Publish Year:** 2020

[Perineal Endometriosis in an Episiotomy Scar: Case Report ...](#)

<https://journals.sagepub.com/doi/abs/10.5301/JE.2012.9410>

Jul 17, 2012 · Perineal endometriosis is a rare disease characterized by the **presence of ectopic endometrial stroma and glands in the perineum**. Most commonly observed in the episiotomy scar after normal vaginal deliveries, perineal endometriosis is not often considered in the differential diagnosis of perineal masses.

Cited by: 1 **Author:** Teresa Tam, Stella Huang

Publish Year: 2012

[The clinical characteristics and surgical approach of scar ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6087557>

Nov 14, 2017 · Clinically, scar endometriosis is **difficult to diagnose**, but **several symptoms commonly occur, including cyclic pain during menstruation [15] and swelling in the pelvic region**. Similarly, most of our patients had been referred to the hospital due to their complaints of ...

Cited by: 6 **Author:** Faik Tatli, Orhan Gozeneli, Hacer Uyanikogl...

Publish Year: 2017

[The clinical characteristics and surgical approach of ...](#)

<https://www.researchgate.net/publication/322131203...>

Request PDF | The **clinical characteristics** and surgical approach of abdominal wall **endometriosis: A case series** of 14 women | Abdominal wall **endometriosis** (AWE) is a rare form of **endometriosis** ...