

# World Journal of *Gastrointestinal Endoscopy*

*World J Gastrointest Endosc* 2020 December 16; 12(12): 504-559



**REVIEW**

- 504 Assessment of disease activity in inflammatory bowel diseases: Non-invasive biomarkers and endoscopic scores

*Rodrigues BL, Mazzaro MC, Nagasako CK, Ayrizono MLS, Fagundes JJ, Leal RF*

**MINIREVIEWS**

- 521 Recent advances in third space or intramural endoscopy

*Shanbhag AB, Thota PN, Sanaka MR*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 532 Efficacy and safety of peroral endoscopic myotomy after prior sleeve gastrectomy and gastric bypass surgery

*Kolb JM, Jonas D, Funari MP, Hammad H, Menard-Katcher P, Wagh MS*

- 542 Crohn's strictures open with anti-mycobacterial antibiotic therapy: A retrospective review

*Collyer R, Clancy A, Agrawal G, Borody TJ*

**CASE REPORT**

- 555 Common bile duct lesions - how cholangioscopy helps rule out intraductal papillary neoplasms of the bile duct: A case report

*Cocca S, Grande G, Reggiani Bonetti L, Magistri P, Di Sandro S, Di Benedetto F, Conigliaro R, Bertani H*

**ABOUT COVER**

Editorial board member of *World Journal of Gastrointestinal Endoscopy*, Dr. José Celso Ardeng is Distinguished Professor at the University of São Paulo, Department of Surgery and Anatomy, Hospital das Clínicas, Faculty of Medicine of Ribeirão Preto (HCFMRP-US), and the Federal University of São Paulo (Department of Diagnostic Imaging at Escola Paulista de Medicina) in Brazil. He completed his postgraduate studies in gastrointestinal surgery at the Santa Casa de Misericórdia de São Paulo, receiving his Master's degree (1999) and PhD (2002). Later, he obtained the title of free-Professor of Medicine (2009) at the University of São Paulo. His research interests involve the application of evidence-based medicine in digestive diseases, particularly studying the effects of advanced treatment methods. His professional accomplishments also include being Former President of the São Paulo State of Brazilian Society of Digestive Endoscopy (2008/2010) and the São Paulo Society of Gastroenterology (2016/2018). (L-Editor: Filipodia)

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Endoscopy* (WJGE, *World J Gastrointest Endosc*) is to provide scholars and readers from various fields of gastrointestinal endoscopy with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGE mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal endoscopy and covering a wide range of topics including capsule endoscopy, colonoscopy, double-balloon enteroscopy, duodenoscopy, endoscopic retrograde cholangiopancreatography, endosonography, esophagoscopy, gastrointestinal endoscopy, gastroscopy, laparoscopy, natural orifice endoscopic surgery, proctoscopy, and sigmoidoscopy.

**INDEXING/ABSTRACTING**

The WJGE is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, China National Knowledge Infrastructure (CNKI), and Superstar Journals Database.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Li-Li Wang; Production Department Director: Yun-Xiao Jian Wu; Editorial Office Director: Jia-Ping Yan.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Endoscopy*

**ISSN**

ISSN 1948-5190 (online)

**LAUNCH DATE**

October 15, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Anastasios Koulaouzidis, Bing Hu, Sang Chul Lee

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5190/editorialboard.htm>

**PUBLICATION DATE**

December 16, 2020

**COPYRIGHT**

© 2020 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Common bile duct lesions - how cholangioscopy helps rule out intraductal papillary neoplasms of the bile duct: A case report

Silvia Cocca, Giuseppe Grande, Luca Reggiani Bonetti, Paolo Magistri, Stefano Di Sandro, Fabrizio Di Benedetto, Rita Conigliaro, Helga Bertani

**ORCID number:** Silvia Cocca 0000-0002-0642-8054; Giuseppe Grande 0000-0003-3907-9740; Luca Reggiani Bonetti 0000-0001-9132-4374; Paolo Magistri 0000-0001-8326-069X; Stefano Di Sandro 0000-0002-5217-8885; Fabrizio Di Benedetto 0000-0002-0693-7430; Rita Conigliaro 0000-0003-3463-7862; Helga Bertani 0000-0003-0818-7167.

**Author contributions:** Bertani H, Grande G, Cocca S, Di Benedetto F and Di Sandro S treated the patient as endoscopists and surgeons, reviewed the literature and contributed to manuscript drafting; Magistri P reviewed the literature and contributed to manuscript drafting; Reggiani Bonetti L performed the histopathological analyses and interpretation and contributed to manuscript drafting; Conigliaro R, Di Benedetto F and Bertani H were responsible for revision of the manuscript for important intellectual content; all authors issued final approval of the version to be submitted.

### Informed consent statement:

Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest.

**Silvia Cocca, Giuseppe Grande, Rita Conigliaro, Helga Bertani**, Endoscopy Unit, Azienda Ospedaliero-Universitaria di Modena, Modena 41121, MO, Italy

**Luca Reggiani Bonetti**, Department of Pathologic Anatomy, University of Modena and Reggio Emilia, Modena 41124, MO, Italy

**Paolo Magistri, Stefano Di Sandro, Fabrizio Di Benedetto**, Hepato-pancreato-biliary Surgery and Liver Transplantation Unit, University of Modena and Reggio Emilia, Modena 41124, MO, Italy

**Corresponding author:** Silvia Cocca, MD, Doctor, Endoscopy Unit, Azienda Ospedaliero-Universitaria di Modena, Via del Pozzo, Modena 41121, MO, Italy. [cocca.silvia@aou.mo.it](mailto:cocca.silvia@aou.mo.it)

## Abstract

### BACKGROUND

Intraductal papillary neoplasm of the bile duct (IPNB) is a rare variant of bile duct tumors, characterized by an exophytic growth exhibiting a papillary mass within the bile duct lumen and it can be localized anywhere along the biliary tree, with morphological variations and occasional invasion.

### CASE SUMMARY

We present a patient with obstructive jaundice who was diagnosed with IPNB using cholangioscopy during endoscopic retrograde cholangio-pancreatography. Using the SpyGlass DS II technology, we were able to define tumor extension and obtain targeted Spy-byte biopsies. After multidisciplinary evaluation, the patient was scheduled for surgical resection of the tumor, which was radically removed.

### CONCLUSION

Cholangioscopy appears to be crucial for the rapid and clear diagnosis of lesions in the bile duct to achieve radical surgical resection.

**Key Words:** Intraductal papillary neoplasm; Cholangioscopy; Bile duct; Surgical resection; SpyGlass; Case report

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

**CARE Checklist (2016) statement:**

The authors have read the CARE Checklist, and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Specialty type:** Gastroenterology and hepatology

**Country/Territory of origin:** Italy

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): C  
Grade D (Fair): D  
Grade E (Poor): 0

**Received:** August 18, 2020

**Peer-review started:** August 18, 2020

**First decision:** September 18, 2020

**Revised:** September 28, 2020

**Accepted:** November 4, 2020

**Article in press:** November 4, 2020

**Published online:** December 16, 2020

**P-Reviewer:** Rerknimitr R

**S-Editor:** Chen XF

**L-Editor:** Webster JR

**P-Editor:** Liu JH



**Core Tip:** Intraductal papillary neoplasm of the bile duct (IPNB) is a rare variant of bile duct tumors, characterized by an exophytic growth exhibiting a papillary mass within the bile duct lumen and it can be localized anywhere along the biliary tree, with morphological variations and occasional invasion. We present a patient with obstructive jaundice who was diagnosed with IPNB using cholangioscopy during endoscopic retrograde cholangio-pancreatography. Using the SpyGlass DS II technology we were able to define tumor extension and to obtain targeted Spy-byte biopsies. The patient underwent successful surgical resection of the tumor.

**Citation:** Cocca S, Grande G, Reggiani Bonetti L, Magistri P, Di Sandro S, Di Benedetto F, Conigliaro R, Bertani H. Common bile duct lesions - how cholangioscopy helps rule out intraductal papillary neoplasms of the bile duct: A case report. *World J Gastrointest Endosc* 2020; 12(12): 555-559

**URL:** <https://www.wjgnet.com/1948-5190/full/v12/i12/555.htm>

**DOI:** <https://dx.doi.org/10.4253/wjge.v12.i12.555>

## INTRODUCTION

Intraductal papillary neoplasm of the bile duct (IPNB) is a rare variant of bile duct tumors, characterized by an exophytic growth exhibiting a papillary mass within the bile duct lumen and it can be localized anywhere along the biliary tree, with morphological variations and occasional invasion. IPNB can be classified into extra-hepatic IPNB and intra-hepatic IPNB. Surgical resection is the first-line treatment and the prognosis of this tumor is better than that of cholangiocarcinoma<sup>[1,2]</sup>.

## CASE PRESENTATION

### Chief complaints

A 72-year-old male patient was admitted to our hospital due to obstructive jaundice and abdominal pain.

### History of present illness

He was a pacemaker carrier and had a history of significant cardiovascular comorbidities.

### History of past illness

No significant past illnesses, apart from cardiovascular disorders.

### Personal and family history

No family history of gastrointestinal tumors.

### Physical examination

On physical examination there was tenderness in the right upper quadrant of the abdomen. Temperature was normal and blood pressure was 100/60 mmHg.


### Laboratory examinations

Blood tests showed high direct bilirubin levels (6.5 mg/dL) and altered liver function tests.

### Imaging examinations

Computed tomography scan revealed a 3 cm hyperdense mass within the common hepatic duct (CHD). Cholangiography during endoscopic retrograde cholangio-pancreatography (ERCP) confirmed the presence of segmental CHD dilation without clear filling defects in the distal tract (Figure 1).





**Figure 1** Cholangiography findings during endoscopic retrograde cholangio-pancreatography of the intraductal papillary neoplasm of the bile duct.

## FINAL DIAGNOSIS

Peroral cholangioscopy (POC), using the SpyGlass DS II technology (Boston Scientific Ltd., Ireland) was performed. After extensive irrigation due to the presence of secretions, an intraductal papillary mass, characterized by papillary projections with fine vascular cores protruding into the lumen was visualized (Figure 2). The mass involved the CHD and was 1 cm from the hilar bifurcation (Klatskin Tumor type I), extending 1 cm up to the cystic insertion in the CHD. Targeted Spy-byte biopsies were obtained, and the results of pathologic examination were consistent with IPNB.

## TREATMENT

After multidisciplinary evaluation, the patient was scheduled for CHD surgical resection, cholecystectomy and hepatico-jejunostomy.

## OUTCOME AND FOLLOW-UP

Final pathology showed IPNB within different grades of dysplasia and invasive cancer involving 3% of the lesion, which was radically resected (R0 margins) (Figure 3). The post-operative course was uneventful and a good prognosis is expected at this stage.

## DISCUSSION

POC is an essential tool in the diagnosis and management of undetermined biliary strictures or filling defects, and the accuracy of the direct visual impression during cholangioscopy seems to be greater than that with ERCP. Moreover, POC allows exact localization of the tumor, which is crucial for surgical planning in order to perform a radical resection<sup>[3-6]</sup>.

## CONCLUSION

The SpyGlass technology appears to be crucial for the rapid diagnosis and a clear evaluation of the extension of lesions such as IPNB, which can easily be misdiagnosed due to their mucin production and flat appearance.




Figure 2 Image of the intraductal papillary neoplasm of the bile duct visualized using the SpyGlass DS II technology.




Figure 3 Hematoxylin and eosin staining results. A: Whole section of the intraductal papillary neoplasm of the bile duct with villous silhouette and pseudo-infiltration of the stroma - arrow (1.25 ×); B and C: Foci of marked cytological atypia in papillomatous background architecture (20 ×).

## REFERENCES

- 1 Nakanuma Y, Kakuda Y, Uesaka K. Characterization of Intraductal Papillary Neoplasm of the Bile Duct with Respect to the Histopathologic Similarities to Pancreatic Intraductal Papillary Mucinous Neoplasm. *Gut Liver* 2019; **13**: 617-627 [PMID: [30982236](#) DOI: [10.5009/gnl18476](#)]
- 2 Lendvai G, Szekerczés T, Illyés I, Dóra R, Kontsek E, Gógl A, Kiss A, Werling K, Kovalszky I, Schaff Z, Borka K. Cholangiocarcinoma: Classification, Histopathology and Molecular Carcinogenesis. *Pathol Oncol Res* 2020; **26**: 3-15 [PMID: [30448973](#) DOI: [10.1007/s12253-018-0491-8](#)]
- 3 Bill JG, Chatterjee D, Mullady DK. Using peroral cholangioscopy to diagnose an intraductal papillary neoplasm of the bile duct. *VideoGIE* 2020; **5**: 68-71 [PMID: [32051913](#) DOI: [10.1007/s12253-018-0491-8](#)]

[10.1016/j.vgie.2019.11.008](https://doi.org/10.1016/j.vgie.2019.11.008)]

- 4 **Korrapati P**, Ciolino J, Wani S, Shah J, Watson R, Muthusamy VR, Klapman J, Komanduri S. The efficacy of peroral cholangioscopy for difficult bile duct stones and indeterminate strictures: a systematic review and meta-analysis. *Endosc Int Open* 2016; **4**: E263-E275 [PMID: [27004242](https://pubmed.ncbi.nlm.nih.gov/27004242/) DOI: [10.1055/s-0042-100194](https://doi.org/10.1055/s-0042-100194)]
- 5 **Nanashima A**, Imamura N, Hiyoshi M, Hamada T, Yano K, Wada T, Kawakami H, Ban T, Kubota Y, Sato Y, Harada K. Planned limited resection of the extrahepatic bile duct in a case of intraductal papillary neoplasm of the bile duct based on preoperative examinations. *Clin J Gastroenterol* 2020; **13**: 233-239 [PMID: [31598905](https://pubmed.ncbi.nlm.nih.gov/31598905/) DOI: [10.1007/s12328-019-01049-8](https://doi.org/10.1007/s12328-019-01049-8)]
- 6 **Ogawa T**, Ito K, Koshita S, Kanno Y, Masu K, Kusunose H, Sakai T, Murabayashi T, Hasegawa S, Noda Y. Usefulness of cholangioscopic-guided mapping biopsy using SpyGlass DS for preoperative evaluation of extrahepatic cholangiocarcinoma: a pilot study. *Endosc Int Open* 2018; **6**: E199-E204 [PMID: [29399618](https://pubmed.ncbi.nlm.nih.gov/29399618/) DOI: [10.1055/s-0043-117949](https://doi.org/10.1055/s-0043-117949)]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

