



Name of the journal: World Journal of Hepatology

Manuscript type: ORIGINAL ARTICLE

Retrospective study

Early tacrolimus exposure does not impact long-term outcomes after liver transplantation

Gastaca M *et al.* Early tacrolimus exposure after liver transplantation

Mikel Gastaca, Patricia Ruiz, Javier Bustamante, Lorea Martinez-Indart, Alberto

Match Overview

1	Crossref 159 words Gastaca, Mikel, Andrés Valdivieso, Javier Bustamante, José R. Fernández, Patricia Ruiz, Alberto Ventoso, Milagro	3%
2	Crossref 33 words Luis A Ruiz, Leyre Serrano, Pedro P España, Lorea Martinez-Indart et al. "Factors influencing long-term survival ..."	1%
3	Crossref 30 words Alberto Lué, Elena Martinez, Mercedes Navarro, Viviana Laredo et al. "Donor Age Predicts Calcineurin Inhibitor ..."	1%
4	Internet 19 words crawled on 24-Jan-2020 ascopubs.org	<1%
5	Crossref 17 words Rodríguez-Perálvarez, M., G. Germani, T. Darius, J. Lerut, E. Tsochatzis, and A. K. Burroughs. "Tacrolimus Trou"	<1%
6	Internet 16 words crawled on 03-Mar-2016 ccsenet.org	<1%
	Internet 15 words	<1%

激活 Windows
转到“设置”以激活 Windows。

Early tacrolimus exposure does not impact long-term outcomes aft



ALL IMAGES VIDEOS

920,000 Results Any time ▾

Early tacrolimus exposure after liver transplantation ...

<https://www.sciencedirect.com/science/article/pii/S0168827812007507>

Feb 01, 2013 · This high exposure to tacrolimus, especially early after LT, leads to adverse effects, most importantly a two-fold increase in early renal impairment , , , which is independently associated with chronic renal impairment and death . A lower early tacrolimus TC than currently recommended should reduce short and long-term adverse effects, but could result in more moderate/severe histological ACR and graft loss.

Cited by: 83 Author: Manuel Rodríguez-Perálvarez, Manuel Rodr...
Publish Year: 2013

Tacrolimus Trough Concentrations After Liver ...

<https://journals.lww.com/transplantjournal/...>

A systematic review and meta-analysis of 32 randomized trials comparing different targets of tacrolimus TC found a linear correlation between mean TC of tacrolimus within the first month and renal impairment rates at 1 year after LT. 2 Indeed, patients receiving mean tacrolimus TC > 10 ng/mL showed twice higher renal impairment rates at 12 months. 2 Mean tacrolimus TCs > 10 ng/mL were also associated with ...

Tacrolimus Exposure After Liver Transplantation in ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/ajt.12216>

Apr 26, 2013 · To the Editor: . We read with great interest the randomized controlled trial (RCT) by De Simone et al. 1, which evaluated the combination of tacrolimus and everolimus after liver transplantation (LT).They demonstrated that glomerular filtration rate decreased less at 12 months with everolimus and reduced tacrolimus, compared to a conventional tacrolimus based regimen, while rejection rates ...

Cited by: 7 Author: M Rodríguez-Perálvarez, M Rodríguez-Perál...
Publish Year: 2013

PEOPLE ALSO ASK

Can tacrolimus improve kidney transplant outcomes? ▾

Can tacrolimus be taken before or after a transplant? ▾

What are the adverse effects of immunosuppression? ▾

What is the pharmacokinetics of tacrolimus in liver transplant recipients?



ALL

IMAGES

VIDEOS

Add the Give with Bing extension >

778,000 Results

Any time ▾

Open links in new tab



Early tacrolimus exposure after liver transplantation ...

<https://www.sciencedirect.com/science/article/pii/S0168827812007507>

Feb 01, 2013 · This high exposure to tacrolimus, especially early after LT, leads to adverse effects, most importantly a two-fold increase in early renal impairment , , , which is independently associated with chronic renal impairment and death . A lower early tacrolimus TC than currently recommended should reduce short and long-term adverse effects, but could result in more moderate/severe histological ACR and **graft loss**.

Cited by: 82

Author: Manuel Rodríguez-Perálvarez, Giaco...

Publish Year: 2013

Early tacrolimus exposure after liver transplantation ...

<https://www.researchgate.net/publication/231612717...>

Background & aims: Liver transplant (LT) patients might be overimmunosuppressed as recommendations for **tacrolimus trough concentrations** (TC) within **4-6 weeks** after liver transplantation are set too...

PEOPLE ALSO ASK

Tacrolimus

Common brand names: Protopic, Prograf

Tacrolimus is used with other medications to prevent rejection of a kidney transplant.

Precautions



Side Effects



Interactions



Warnings

Brand names: Prograf · Astagraf XL · Envarsus XR · Protopic +**Drug classes:** Calcineurin Inhibitor Immunosuppressant · Immune Suppressant**Pregnancy category:** Consult a doctor**Availability:** Prescription needed**May treat:** Transplant rejection · 激活 Windows

转到“设置”以激活 Windows。

People also search for

802,000 Results Any time ▾

Early tacrolimus exposure after liver transplantation ...

<https://www.sciencedirect.com/science/article/pii/S0168827812007507>

Feb 01, 2013 · This high **exposure** to **tacrolimus**, especially **early after LT**, leads to adverse effects, most importantly a two-fold increase in **early renal** impairment , , , which is independently associated with chronic renal impairment and death . A lower **early tacrolimus TC** than currently recommended should reduce short and **long-term** adverse effects, but could result in more moderate/severe histological ...

Cited by: 85 **Author:** Manuel Rodríguez-Perálvarez, Giacomo Ge...

Publish Year: 2013

Early tacrolimus exposure after liver transplantation ...

<https://www.researchgate.net/publication/231612717...>

Early tacrolimus exposure did not influence chronic rejection (p = 0.58), or chronic renal impairment (p = 0.25). **Conclusions:** During the first 2 weeks after LT, tacrolimus TC between 7 and 10...

PEOPLE ALSO ASK

Is tacrolimus better than cyclosporine? ▾

Can tacrolimus be taken before or after a transplant? ▾

What are the adverse effects of immunosuppression? ▾

What are the side effects of tacrolimus? ▾

Feedback

Tacrolimus Exposure After Liver Transplantation in ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/ajt.12216>

To the Editor: . We read with great interest the randomized controlled trial (RCT) by De Simone et al. 1, which evaluated the combination of **tacrolimus** and everolimus **after liver transplantation** (LT).They demonstrated that glomerular filtration rate decreased less at 12 months with everolimus and reduced **tacrolimus**, compared to a conventional **tacrolimus** based regimen, while rejection rates ...

Cited by: 7 **Author:** M Rodríguez-Perálvarez, M Rodríguez-Perál...

Publish Year: 2013

Tacrolimus

Common brand names: Protopic, Prograf

Tacrolimus is used with other medications to prevent rejection of a kidney transplant.

Precautions

 Side Effects
  Interactions
  Warnings

Brand names: Prograf · Astagraf XL · Envarsus XR · Protopic · Hecoria +

Drug classes: Calcineurin Inhibitor Immunosuppressant · Immune Suppressant

Pregnancy category: Consult a doctor

Availability: Prescription needed

May treat: Transplant rejection · Lichen Planus

People also search for

[Mycophenolic acid](#)

[cyclosporine](#)

[sirolimus](#)

[azathioprine](#)

[pimecrolimus](#)

[See more ▾](#)

Data from First Databank and others.[Learn more](#)

Consult a medical professional for advice

[Suggest an edit](#)