

**Name of Journal:** *World Journal of Clinical Oncology*

**Manuscript NO:** 58306

**Manuscript Type:** CASE REPORT

**Preoperative rectal tumor embolization as an adjunctive tool for bloodless abdominoperineal excision: A case report.**

Feitosa MR *et al.* Preoperative rectal tumor embolization.

Marley Ribeiro Feitosa, Lucas Fernandes de Freitas, Antonio Balestrim Filho, Guilherme S Nakiri, Daniel Giansante Abud, Ligia Magnani Landell, Mariângela Ottoboni Brunaldi, Jose JR da Rocha, Omar Feres, Rogério Serafim Parra

## Match Overview

1	<b>Internet</b> 36 words crawled on 22-Feb-2020 <a href="http://www.labome.org">www.labome.org</a>	2%
2	<b>Crossref</b> 36 words LIMING WANG, YASUMITSU HIRANO, GREGORY HEN G, TOSHIMASA ISHII <i>et al.</i> "Prognostic Utility of Apical ...	2%
3	<b>Crossref</b> 36 words Christina Wu, Terence M Williams, Ryan Robb, Amy Webb <i>et al.</i> "Phase I Trial of Trametinib with Neoadjuvant Ch...	2%
4	<b>Internet</b> 34 words crawled on 23-Mar-2020 <a href="http://www.scielo.br">www.scielo.br</a>	2%
5	<b>Internet</b> 33 words crawled on 25-Jul-2020 <a href="http://jcancer.org">jcancer.org</a>	2%
6	<b>Internet</b> 31 words crawled on 17-Jan-2020 <a href="http://cancercommun.biomedcentral.com">cancercommun.biomedcentral.com</a>	2%



Preoperative rectal tumor embolization as an adjunctive tool for blo



ALL

IMAGES

VIDEOS

30,500 Results

Any time ▾

## Preoperative versus Postoperative Chemoradiotherapy for ...

<https://www.nejm.org/doi/full/10.1056/NEJMoa040694>

Among patients with tumors judged by the surgeon to require an abdominoperineal excision, the rate of sphincter-preserving surgery was more than doubled after preoperative chemoradiotherapy ...

Cited by: 6353

Author: Rolf Sauer, Heinz Becker, Werner Hohen...

Publish Year: 2004

## Uncommon Colorectal Neoplasms: Retrorectal Tumors

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3311502>

Patients with locally recurrent and metastatic chordoma respond well to chemotherapeutic agents such as cetuximab and gefitinab. 40, 41, 42 Transcatheter arterial embolization (TAE) of tumors including chordomas has been performed as an adjunct to successful resection. TAE can significantly decrease intraoperative blood loss, make the surgical ...

### PEOPLE ALSO ASK

Can chemo be used to treat Stage IV rectal cancer? ▾

What is the best treatment for a rectal tumor? ▾

What is the procedure for rectal cancer? ▾

Can a polyp be removed during colonoscopy? ▾

Feedback

## Large hemangiopericytoma of the pelvis—towards a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4551704>

Case presentation. A 70-year-old female patient was presented with lower abdominal pain and constipation at our hospital. A computed tomography (CT) was performed and revealed a highly vascularized solid mass measuring 9 × 14 × 19 cm, congestion of the kidneys, and compression of the rectum (Fig 1). The highly vascularisation raised concerns of high blood loss during surgery, and ...

Cited by: 2

Author: Mohammad Fard-Aghaie, Gregor A. Stavr...

Publish Year: 2015

国内版

国际版



Preoperative rectal tumor embolization as an adjunctive tool for blo



ALL

IMAGES

VIDEOS

57,800 Results

Any time ▾

## Preoperative versus Postoperative Chemoradiotherapy for ...

<https://www.nejm.org/doi/10.1056/NEJMoa040694>

Among patients with **tumors** judged by the surgeon to require an **abdominoperineal excision**, the rate of sphincter-preserving surgery was more than doubled after **preoperative** chemoradiotherapy ...

Cited by: 6665

Author: Rolf Sauer, Heinz Becker, Werner Hohenber...

Publish Year: 2004

## Uncommon Colorectal Neoplasms: Retrorectal Tumors

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3311502>

Patients with locally recurrent and metastatic chordoma respond well to chemotherapeutic agents such as cetuximab and gefitinab. 40, 41, 42 Transcatheter arterial **embolization** (TAE) of **tumors** including chordomas has been performed as an **adjunct** to successful resection. TAE can significantly decrease intraoperative **blood** loss, make the surgical ...

PEOPLE ALSO ASK

What is the procedure for rectal skin removal?





[ALL](#)
[IMAGES](#)
[VIDEOS](#)
[MAPS](#)
[NEWS](#)
[SHOPPING](#)

30,400 Results Any time ▾

## Preoperative versus Postoperative Chemoradiotherapy for ...

<https://www.nejm.org/doi/10.1056/NEJMoa040694>

Among patients with **tumors** judged by the surgeon to require an **abdominoperineal excision**, the rate of sphincter-preserving surgery was more than doubled after **preoperative** chemoradiotherapy ...

**Cited by:** 7080      **Author:** Rolf Sauer, Heinz Becker, Werner Hohen...  
**Publish Year:** 2004

## Utility of preoperative 3-D simulation of laparoscopic ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/ases.12476>

Mar 12, 2018 · The records of 10 consecutive patients with advanced low **rectal cancer** (below the peritoneal reflection) who underwent laparoscopic unilateral LPLD following total mesorectal **excision**, **tumor-specific mesorectal excision**, or total proctocolectomy after **preoperative** 3-D simulation were reviewed retrospectively.

**Author:** Hisanaga Horie, Koji Koinuma, Homar...      **Publish Year:** 2018

## Large hemangiopericytoma of the pelvis—towards a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4551704>

**Case** presentation. A 70-year-old female patient was presented with lower abdominal pain and constipation at our hospital. A computed tomography (CT) was performed and revealed a highly vascularized solid mass measuring 9 × 14 × 19 cm, congestion of the kidneys, and compression of the **rectum** (Fig 1). The highly vascularisation raised concerns of high **blood** loss during surgery, and ...

**Cited by:** 3      **Author:** Mohammad Fard-Aghaie, Gregor A. Stav...  
**Publish Year:** 2015

## Association of Coloproctology of Great Britain & Ireland ...

<https://onlinelibrary.wiley.com/doi/10.1111/codi.13704>

Jun 20, 2017 · **Abdominoperineal excision** should not be regarded as an inferior operation to low anterior resection in the management of low **rectal cancer** on the basis of quality of life alone (How et al., 2012). Bowel function after a low anterior resection is often problematic and many patients have urgency and frequency, partly attributable to loss of the ...

**Cited by:** 52      **Author:** Simon Gollins, Brendan Moran, Richard