



1829-7197-58437_Auto_Edited.docx

Quotes Excluded
Bibliography Excluded2%
SIMILAR

² Name of Journal: *World Journal of Gastrointestinal Pharmacology and Therapeutics*

Manuscript NO: 58437

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

¹ Clinical features and relative factors of constipation in a cohort of Chinese patients with parkinson's disease

¹ Constipation in Chinese patients with parkinson's disease

Baihua Sun, Tao Wang, Nianying LI, Qiong Wu, Jin Qiao

Match Overview

1	Internet 27 words crawled on 24-Sep-2020 www.hindawi.com	1%
2	Internet 17 words crawled on 01-Nov-2019 f6publishing.blob.core.windows.net	<1%
3	Internet 15 words crawled on 09-Dec-2018 id.scribd.com	<1%
4	Internet 14 words crawled on 07-Oct-2020 pesquisa.bvsalud.org	<1%
5	Crossref 13 words Tetsuya UEKI, Keiko NAGAI, Yasuko MIZUKAMI, Akihiro TAKAHASHI et al. "Cross-sectional Study on Relationsh...	<1%
6	Crossref 13 words "Poster Presentations", Movement Disorders, 2012	<1%



Clinical features and relative factors of constipation in a cohort of CI



ALL

IMAGES

VIDEOS

2,680,000 Results

Any time ▾

Parkinson disease with constipation: clinical features and ...

<https://www.nature.com/articles/s41598-017-16790-8>

Jan 12, 2018 · In this study, 61.4% PD patients had constipation, in accordance with previous studies reporting 50% to 80% of constipation in PD patients 9,10, thus, constipation is a ...

Cited by: 20

Author: Qiu-Jin Yu, Shu-Yang Yu, Li-Jun Zuo, Ten...

Publish Year: 2018

Parkinson's disease in China: a forty-year growing track ...

<https://translationalneurodegeneration.biomedcentral.com/...> ▾

Jul 31, 2019 · About 54.10% of PD patients suffered from constipation, and 47.59% of them reported constipation before onset of the motor symptoms . A cross-sectional survey in Chinese PD patients in Hoehn-Yahr stage I-III revealed that 29.9% patients suffered from PD-related pain, associated with dyskinesias, and/or depression .

Cited by: 8

Author: Gen Li, Jianfang Ma, Shishuang Cui, Yixi ...

Publish Year: 2019

Parkinson's Disease Constipation: Causes & Treatments

<https://www.webmd.com/parkinsons-disease/guide/parkinsons-constipation> ▾

Mar 20, 2002 · Constipation often affects those with Parkinson's disease. It occurs when bowel movements become difficult or less frequent. The normal length of time between bowel movements (also known as "stools ...

PEOPLE ALSO ASK

What causes constipation in Parkinson s disease? ▾

Does Parkinson s disease affect bowel movements? ▾

Why do I have constipation with Parkinson's? ▾

Feedback

Management of constipation in patients with Parkinson's ...

<https://www.nature.com/articles/s41531-018-0042-8>

Clinical features and relative factors of constipation in a cohort of C



ALL

IMAGES

VIDEOS

2,740,000 Results

Any time ▼

[Parkinson's disease in China: a forty-year growing track ...](https://translationalneurodegeneration.biomedcentral.com/...)

<https://translationalneurodegeneration.biomedcentral.com/...> ▼

Jul 31, 2019 · About 54.10% of PD **patients** suffered from **constipation**, and 47.59% of them reported **constipation** before onset of the motor symptoms . A cross-sectional survey in **Chinese PD patients** in Hoehn-Yahr stage I-III revealed that 29.9% **patients** suffered from PD-related pain, associated with dyskinesias, and/or depression .

Cited by: 8**Author:** Gen Li, Jianfang Ma, Shishuang Cui, Yixi He...**Publish Year:** 2019

[Constipation: an emerging risk factor for Parkinson's disease?](https://www.researchgate.net/publication/305622568...)

<https://www.researchgate.net/publication/305622568...>

Constipation is the most prominent and disabling manifestation of lower gastrointestinal (GI) dysfunction in **Parkinson's disease** (PD). The prevalence of **constipation** in PD **patients** ranges from 24 ...

[Table of Contents | Parkinson's Disease | Hindawi](https://www.hindawi.com/journals/pd/contents)

<https://www.hindawi.com/journals/pd/contents> ▼

Oct 10, 2020 · **Parkinson's Disease**-Special Issue - Volume 2020 - Article ID 8692509 - Research Article; Analysis of Motor Complication and Relative Factors in a Cohort of Chinese Patients with Parkinson's ...

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

2,880,000 Results

Any time ▾

Constipation preceding Parkinson's disease: a systematic ...

<https://jnnp.bmj.com/content/87/7/710> ▾

Jul 01, 2016 · Introduction. **Parkinson's disease** (PD) is the second most common neurodegenerative disorder with a prevalence of approximately 0.4%—a figure which is expected to double by 2040.¹ PD is diagnosed when motor **features** such as tremor, bradykinesia and rigidity become overt, by which time approximately 50% of neurons within the substantia nigra remain.² Over the past two decades, a ...

Cited by: 96

Author: Kerala L Adams-Carr, Jonathan P Bestwi...

Publish Year: 2016

Constipation: an emerging risk factor for Parkinson's disease?

<https://www.researchgate.net/publication/305622568...>

Constipation is the most prominent and disabling manifestation of lower gastrointestinal (GI) dysfunction in **Parkinson's disease** (PD). The prevalence of **constipation** in PD **patients** ranges from 24 ...

A survey of subjective constipation in Parkinson's disease ...

<https://bmcneurol.biomedcentral.com/articles/10.1186/s12883-018-1034-3> ▾

Mar 15, 2018 · **Constipation** is one of the most frequent non-motor symptoms (NMS) in **Parkinson's disease** (PD) and the prevalence of **constipation** in PD **patients** varies among different studies. We designed this study to survey the prevalence and **clinical** characteristics of subjective **constipation** and the appearance chronology between the emergence of **constipation** and onset of motor symptoms in PD **patients** ...

Cited by: 8

Author: Jing Gan, Ying Wan, Junjie Shi, Mingzhu ...

Publish Year: 2018

(PDF) Management of constipation in Parkinson's disease

<https://www.researchgate.net/publication/270002789...>

Constipation is prevalent in **Parkinson's disease** (PD), 1-3 distressing to many **patients**, and sometimes leads to serious complications such as intestinal pseudo-obstruction, volvulus, and acute ...

Constipation and risk of Parkinson's disease: A Danish ...

Parkinson's Disease

Medical Condition



A chronic and progressive movement disorder.

- Common (More than 200,000 cases per year in US)
- Often requires lab test or imaging
- Treatments can help manage condition, no known cure
- Can last several years or be lifelong

Parkinson's disease is caused by a loss of nerve cells in substantia nigra of the brain. This leads to a reduction in a chemical called dopamine in the brain. Many different symptoms are associated with Parkinson's disease and the more common symptoms are slowness in movement and muscle stiffness. Currently there is no cure for Parkinson's disease. But treatments are available to help relieve the symptoms and maintain quality of life. These treatments include: - Supportive therapies – such as physiotherapy - Medication - Surgery