

World Journal of *Diabetes*

World J Diabetes 2021 March 15; 12(3): 198-305



Contents**Monthly Volume 12 Number 3 March 15, 2021****OPINION REVIEW**

- 198** Diabetes and COVID-19: Diseases of racial, social and glucose intolerance
Chowdhury TA

EVIDENCE REVIEW

- 206** Progress on haptoglobin and metabolic diseases
Wan BN, Zhou SG, Wang M, Zhang X, Ji G

REVIEW

- 215** Bidirectional link between diabetes mellitus and coronavirus disease 2019 leading to cardiovascular disease: A narrative review
Viswanathan V, Puvvula A, Jamthikar AD, Saba L, Johri AM, Kotsis V, Khanna NN, Dhanjil SK, Majhail M, Misra DP, Agarwal V, Kitas GD, Sharma AM, Kolluri R, Naidu S, Suri JS
- 238** Anti- and non-tumor necrosis factor- α -targeted therapies effects on insulin resistance in rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis
Wang CR, Tsai HW

ORIGINAL ARTICLE**Basic Study**

- 261** Causal effect of education on type 2 diabetes: A network Mendelian randomization study
Liao LZ, Chen ZC, Li WD, Zhuang XD, Liao XX

Clinical Trials Study

- 278** Altered spontaneous brain activity in patients with diabetic optic neuropathy: A resting-state functional magnetic resonance imaging study using regional homogeneity
Guo GY, Zhang LJ, Li B, Liang RB, Ge QM, Shu HY, Li QY, Pan YC, Pei CG, Shao Y

Randomized Controlled Trial

- 292** Effectiveness of cognitive behavior therapy for sleep disturbance and glycemic control in persons with type 2 diabetes mellitus: A community-based randomized controlled trial in China
Zhang HZ, Zhang P, Chang GQ, Xiang QY, Cao H, Zhou JY, Dong ZM, Qiao C, Xu CR, Qin Y, Lou PA

Contents

Monthly Volume 12 Number 3 March 15, 2021

ABOUT COVER

Mustafa Altay, MD, Professor, Chief, Department of Endocrinology and Metabolism, Health Sciences University Turkey, Keçiören Training and Research Hospital, Kuşcağız, Ankara 54600, Turkey. mustafa.altay@sbu.edu.tr

AIMS AND SCOPE

The primary aim of *World Journal of Diabetes* (WJD, *World J Diabetes*) is to provide scholars and readers from various fields of diabetes with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of diabetes and covering a wide range of topics including risk factors for diabetes, diabetes complications, experimental diabetes mellitus, type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes, diabetic angiopathies, diabetic cardiomyopathies, diabetic coma, diabetic ketoacidosis, diabetic nephropathies, diabetic neuropathies, Donohue syndrome, fetal macrosomia, and prediabetic state.

INDEXING/ABSTRACTING

The WJD is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for WJD as 3.247; IF without journal self cites: 3.222; Ranking: 70 among 143 journals in endocrinology and metabolism; and Quartile category: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yu-Jie Ma; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Diabetes

ISSN

ISSN 1948-9358 (online)

LAUNCH DATE

June 15, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Timothy Koch

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9358/editorialboard.htm>

PUBLICATION DATE

March 15, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

© 2021 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

E-mail: bpgoffice@wjgnet.com <https://www.wjgnet.com>



Submit a Manuscript: <https://www.f6publishing.com>*World J Diabetes* 2021 March 15; 12(3): 261-277DOI: [10.4239/wjd.v12.i3.261](https://doi.org/10.4239/wjd.v12.i3.261)

ISSN 1948-9358 (online)

ORIGINAL ARTICLE

Basic Study**Causal effect of education on type 2 diabetes: A network Mendelian randomization study**

Li-Zhen Liao, Zhi-Chong Chen, Wei-Dong Li, Xiao-Dong Zhuang, Xin-Xue Liao

ORCID number: Li-Zhen Liao [0000-0001-5807-2339](#); Zhi-Chong Chen [0000-0002-8691-6399](#); Wei-Dong Li [0000-0003-1430-9749](#); Xiao-Dong Zhuang [0000-0001-6508-8507](#); Xin-Xue Liao [0000-0001-7631-1866](#).

Author contributions: Liao LZ and Chen ZC conceived the study and contributed equally to this study; Li WD completed the analyses; Zhuang XD led the writing; Liao XX supervised the study.

Supported by the National Natural Science Foundation of China, No. 81701378.

Institutional review board statement: Not applicable.

Conflict-of-interest statement: The authors declare that they have no conflict of interest to disclose.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially,

Li-Zhen Liao, Wei-Dong Li, Department of Health, Guangdong Pharmaceutical University, Guangzhou 510275, Guangdong Province, China

Li-Zhen Liao, Wei-Dong Li, Guangdong Provincial Key Laboratory of Pharmaceutical Bioactive Substances, Guangdong Pharmaceutical University, Guangzhou 510006, Guangdong Province, China

Zhi-Chong Chen, Department of Cardiology, The Sixth Affiliated Hospital of Sun Yat-Sen University, Guangzhou 510080, Guangdong Province, China

Xiao-Dong Zhuang, Xin-Xue Liao, Department of Cardiology, The First Affiliated Hospital of Sun Yat-Sen University, Guangzhou 510080, Guangdong Province, China

Corresponding author: Xin-Xue Liao, PhD, Chief Physician, Department of Cardiology, The First Affiliated Hospital of Sun Yat-Sen University, No. 58 Zhongshan 2nd Road, Yuexiu District, Guangzhou 510080, Guangdong Province, China. liaoxinx@mail.sysu.edu.cn

Abstract

BACKGROUND

The causality between education and type 2 diabetes (T2DM) remains unclear.

AIM

To identify the causality between education and T2DM and the potential metabolic risk factors [coronary heart disease (CHD), total cholesterol, low-density lipoprotein, triglycerides (TG), body mass index (BMI), waist circumference (WC), waist-to-hip ratio (WHR), fasting insulin, fasting glucose, and glycated hemoglobin] from summarized genome-wide association study (GWAS) data used a network Mendelian randomization (MR).

METHODS

Two-sample MR and network MR were performed to obtain the causality between education-T2DM, education-mediator, and mediator-T2DM. Summary statistics from the Social Science Genetic Association Consortium (discovery data) and Neale Lab consortium (replication data) were used for education and DIAGRAMplusMetabochip for T2DM.

RESULTS

The odds ratio for T2DM was 0.392 (95%CI: 0.263-0.583) per standard deviation

and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http://creativecommons.org/License_s/by-nc/4.0/

Manuscript source: Unsolicited manuscript

Specialty type: Genetics and heredity

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

Received: July 27, 2020

Peer-review started: July 27, 2020

First decision: November 4, 2020

Revised: November 10, 2020

Accepted: December 23, 2020

Article in press: December 23, 2020

Published online: March 15, 2021

P-Reviewer: Karatza AA

S-Editor: Zhang H

L-Editor: Wang TQ

P-Editor: Yuan YY



increase (3.6 years) in education by the inverse variance weighted method, without heterogeneity or horizontal pleiotropy. Education was genetically associated with CHD, TG, BMI, WC, and WHR in the discovery phase, yet only the results for CHD, BMI, and WC were replicated in the replication data. Moreover, BMI was genetically associated with T2DM.

CONCLUSION

Short education was found to be associated with an increased T2DM risk. BMI might serve as a potential mediator between them.

Key Words: Mendelian randomization; Education; Type 2 diabetes mellitus; Genome-wide association study; Coronary heart disease; Body mass index

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Genetically predicted education was negatively causally associated with type 2 diabetes (T2DM). The odds ratio for T2DM was 0.392 (95%CI: 0.263-0.583) per standard deviation increase (3.6 years) in education. Body mass index might serve as a potential mediator.

Citation: Liao LZ, Chen ZC, Li WD, Zhuang XD, Liao XX. Causal effect of education on type 2 diabetes: A network Mendelian randomization study. *World J Diabetes* 2021; 12(3): 261-277

URL: <https://www.wjgnet.com/1948-9358/full/v12/i3/261.htm>

DOI: <https://dx.doi.org/10.4239/wjd.v12.i3.261>

INTRODUCTION

Type 2 diabetes mellitus (T2DM), affecting approximately 415 million people, has become a major global health problem. This number will rapidly increase to 642 million by 2040^[1]. Environmental, genetic, and metabolic risk factors all contribute to T2DM^[2]. Among these, controlling the modifiable risk factors to decrease the T2DM morbidity, turns into a key public health priority^[3]. A previous systematic review and meta-analysis reported that diabetes self-management education can reduce all-cause mortality risk in T2DM patients^[4], indicating that education may help T2DM management. However, the causality between primary education (years of schooling) and T2DM remains unclear. If the causal association exists, the potential pathways involved in the association from education to T2DM have not yet been studied.

Mendelian randomization (MR) serves as a strategy for assessing the causality between disease and common risk factors^[5]. Since the genetically determined risk factors occur before the disease onset, MR can avoid the potential bias of reverse causation in retrospective studies. The risk of confounding is also reduced^[5]. In this study, the causality between education and T2DM was analyzed and potential metabolic risk factors were explored from summarized genome-wide association study (GWAS) data. The potential metabolic risk factors included coronary heart disease (CHD), total cholesterol (TC), low-density lipoprotein (LDL), triglycerides (TG), body mass index (BMI), waist circumference (WC), waist-to-hip ratio (WHR), fasting insulin, fasting glucose, and glycated hemoglobin (HbA1c).

MATERIALS AND METHODS

GWAS data summary

Summary data from array-based analysis for single nucleotide polymorphism (SNP) was included. We selected all SNPs associated with education at genome-wide significance ($P < 5 \times 10^{-8}$) in the available GWAS (Supplementary Table 1). The metrics of SNP quality were requested as follows. There was strong evidence of between-study heterogeneity in the SNP-trait association ($P \leq 0.005$). The imputation quality metric (info or r^2) was ≤ 0.90 , and there was Hardy-Weinberg disequilibrium ($P \leq$

0.001). Summary statistics from the Social Science Genetic Association Consortium (SSGAC)^[6] and Neale Lab consortium (<http://www.nealelab.is/blog/2017/9/11/details-and-considerations-of-the-uk-biobank-gwas>) were used for education. Because several large summary statistics from different consortia were available for the same biomarker, we used the earliest and largest summary data (SSGAC) as the discovery data set and the most recent data (Neale Lab) as the replication data set. DIAGRAM plus Metabochip consortium data were used for T2DM^[7], CARDIoGRAM plus C4D consortium data were utilized for CHD^[8], GLGC consortia data were used for TC^[9], LDL^[9], and TG^[9], Genetic Investigation of Anthropometric Traits consortium data were used for BMI^[10], WC^[11], and WHR^[12], and Mapping and Geographic Information Consortium data were used for fasting insulin^[13], fasting glucose^[14], and HbA1c^[15]. The datasets are presented in Table 1.

Two-sample MR and causality evaluation

Our MR study was conducted on the MR-Base platform online (<http://www.mrbase.org>)^[16]. We explored the associations as follows: (1) Causality: The conventional MR approach [inverse variance weighted (IVW)] method, MR Egger method, and weighted median method were used: (a) Causality between genetically determined education and T2DM; (b) Causality between education and metabolic risk factors (CHD, TG, LDL, TC, BMI, WC, WHR, fasting insulin, fasting glucose, and HbA1c); and (c) Causality between metabolic risk factors and T2DM; (2) Heterogeneity: We conducted heterogeneity tests in MR analyses using IVW and MR Egger; (3) Horizontal pleiotropy: The MR egger intercept was assessed; (4) Leave-one-out analysis; and (5) Funnel plots.

Network MR for “education-mediator-T2DM” analyses

Two-sample MR and network MR were performed to obtain the causality between education-T2DM, education-mediator, and mediator-T2DM^[18]. A network MR analysis consisted of three two-sample MR tests: (1) The causality between education and T2DM; (2) The causality between education and the potential mediators; and (3) The causality between the potential mediators and T2DM.

We could conclude that the specific metabolic risk factor might serve as a mediator between education and T2DM if the causality was estimated in all three steps.

Statistical analysis

The statistical tests were two-sided. The statistical test for the MR analyses was considered statistically significant at $P < 0.05$. All of the calculations were conducted using Stata (College Station, TX, United States) and R language.

RESULTS

Causality between genetically determined education and T2DM

Characteristics of the SNPs are shown in Supplementary Table 1. We used the SSGAC consortia for education to explore the causal associations between education and T2DM. In the IVW method, the odds ratio [OR (95%CI)] for T2DM was 0.392 (0.263-0.583) per standard deviation increase (3.6 years) in education (Table 2 and Figure 1A). Results were consistent in weighted median method (OR: 0.406, 95%CI: 0.246-0.672; $P = 0.000$) (Table 2). Both IVW and MR Egger estimates indicated no heterogeneity amongst these 17 SNPs ($P = 0.163$ and $P = 0.124$, respectively) (Table 2). There was no directional horizontal pleiotropy (MR egger intercept $P = 0.979$) (Table 2). In a leave-one-out analysis, no single instrument was strongly driving the overall effect of education on T2DM (Figure 1C). Besides, there was no funnel plot asymmetry (Figure 1D). Both the leave-one-out analysis and funnel plot further suggested that no SNPs exhibited horizontal pleiotropy.

To sum up, the genetically predicted education was negatively causally associated with T2DM.

Causality between education and metabolic risk factors

The causality between education and metabolic biomarkers, including CHD, TG, LDL, TC, BMI, WC, WHR, fasting insulin, fasting glucose, and HbA1c, is shown in Table 3. Education was causally associated with CHD, TG, BMI, WC, and WHR in the discovery phase. Yet in the replication data, only the results for CHD, BMI, and WC were duplicated.

Table 1 Details of studies and datasets used for analyses

Exposure/Outcome	Cases, n	Controls, n	Sample size	PubMed ID	First author	Consortium	Year	Units
Education (discovery data)	NA	NA	293723	27225129	Okbay	SSGAC	2016	SD (yr)
Education 2 (replication data)	NA	NA	226899	UKB-a:505	Neale	Neale Lab	2017	NA
T2DM	34840	114981	77312	22885922	Morris	DIAGRAM plus Metabochip	2012	Log odds
CHD	60801	123504	184305	26343387	Nikpay	CARDIoGRAM plus C4D	2015	Log odds
TG	NA	NA	108514	24097068	Willer	GLGC	2013	SD (mg/dL)
LDL	NA	NA	99073	24097068	Willer	GLGC	2013	SD (mg/dL)
TC	NA	NA	108363	24097068	Willer	GLGC	2013	SD (mg/dL)
BMI	NA	NA	152893	25673413	Locke	GIANT	2015	SD (kg/m^2)
WC	NA	NA	232101	25673412	Shungin	GIANT	2015	SD (cm)
WHR	NA	NA	60586	23754948	Randall	GIANT	2013	SD (ratio)
Fasting insulin	NA	NA	38238	20081858	Dupuis	MAGIC	2010	Log pmol/L
Fasting glucose	NA	NA	58074	22581228	Manning	MAGIC	2012	mmol/L
HbA1c	NA	NA	46368	20858683	Soranzo	MAGIC	2010	%

BMI: Body mass index; CHD: Coronary heart disease; GIANT: Genetic Investigation of Anthropometric Traits; HbA1c: Glycated hemoglobin; LDL: Low-density lipoprotein; MAGIC: Mapping and Geographic Information Consortium; NA: Not available; SSGAC: Social Science Genetic Association Consortium; TC: Total cholesterol; T2DM: Type 2 diabetes mellitus; TG: Triglycerides; WC: Waist circumference; WHR: Waist-to-hip ratio.

Table 2 Causal associations between genetically determined education and type 2 diabetes mellitus

Trait	Method	nSNP	OR	95%CI	P value	Heterogeneity P	MR Egger intercept P
Education-T2DM	MR Egger	17	0.381	0.047–3.093	0.381	0.124	0.979
	Weighted median	17	0.406	0.246–0.672	0.000		
	Inverse variance weighted	17	0.392	0.263–0.583	0.000	0.163	

MR: Mendelian randomization; OR: Odds ratio; nSNP: Number of single-nucleotide polymorphisms; T2DM: Type 2 diabetes mellitus.

Causality between metabolic risk factors and T2DM

Based on the above results, CHD, BMI, and WC might serve as potential mediators between education and T2DM. Thus, we further evaluated whether these three potential mediators were associated with T2DM using MR analyses. Only BMI (but not CHD or WC) was positively associated with T2DM (Table 4). Hence, BMI might serve as a potential mediator between education and T2DM.

DISCUSSION

Diabetes has rapidly become a global epidemic and a significant public health concern. Identifying the high-risk populations and addressing the risk factors for diabetes might prove to be an effective diabetes prevention strategy. Traditional risk factors for T2DM include obesity, diet, physical activity, socioeconomic status, etc.^[19], some of which are modifiable through behavioral or pharmacological intervention (e.g., obesity, diet, and physical activity). It is of great importance to determine whether traditional risk factors have a causal role in T2DM or are merely bystanders.

Previous studies have reported that education might help T2DM management. A prospective, randomized, single-center study revealed that pharmacotherapeutic education of patients with T2DM could significantly improve 30-d post-discharge

Table 3 Causal association between genetically determined education and metabolic risk factors

Trait	Method	nSNP	Beta	SE	P value	Heterogeneity P	MR Egger intercept P
Discovery (education-metabolic risk factors)							
CHD	MR Egger	71	0.145	0.417	0.729	0.123	0.263
	Weighted median	71	-0.238	0.107	0.026		
	Inverse variance weighted	71	-0.318	0.079	0.000	0.117	
TG	MR Egger	57	-0.183	0.263	0.489	0.112	0.937
	Weighted median	57	-0.192	0.057	0.001		
	Inverse variance weighted	57	-0.204	0.043	0.000	0.130	
LDL	MR Egger	57	-0.274	0.275	0.324	0.316	0.415
	Weighted median	57	0.007	0.063	0.909		
	Inverse variance weighted	57	-0.051	0.045	0.261	0.326	
TC	MR Egger	57	-0.087	0.277	0.755	0.197	0.838
	Weighted median	57	-0.018	0.064	0.775		
	Inverse variance weighted	57	-0.031	0.045	0.500	0.223	
BMI	MR Egger	59	0.013	0.353	0.970	0.000	0.415
	Weighted median	59	-0.209	0.048	0.000		
	Inverse variance weighted	59	-0.272	0.058	0.000	0.000	
WC	MR Egger	59	-0.002	0.404	0.996	0.000	0.429
	Weighted median	59	-0.269	0.056	0.000		
	Inverse variance weighted	59	-0.319	0.066	0.000	0.000	
WHR	MR Egger	58	-0.743	0.461	0.113	0.028	0.346
	Weighted median	58	-0.298	0.097	0.002		
	Inverse variance weighted	58	-0.311	0.076	0.000	0.027	
Fasting insulin	MR Egger	58	-0.269	0.215	0.215	0.010	0.312
	Weighted median	58	-0.039	0.048	0.419		
	Inverse variance weighted	58	-0.053	0.035	0.129	0.014	
Fasting glucose	MR Egger	58	-0.185	0.172	0.287	0.167	0.398
	Weighted median	58	-0.044	0.039	0.257		
	Inverse variance weighted	58	-0.041	0.028	0.152	0.172	
HbA1c	MR Egger	58	0.004077	0.1779	0.9818	0.5674	0.889
	Weighted median	58	-0.009323	0.04094	0.8199		
	Inverse variance weighted	58	-0.02056	0.02908	0.4796	0.6038	
Replication (education-metabolic risk factors)							
CHD	MR Egger	20	-0.985	0.772	0.218	0.196	0.537
	Weighted median	20	-0.474	0.209	0.024		
	Inverse variance	20	-0.511	0.164	0.002	0.222	

weighted							
TG	MR Egger	13	0.099	1.076	0.928	0.000	0.702
	Weighted median	13	-0.161	0.134	0.229		
	Inverse variance weighted	13	-0.319	0.163	0.051	0.000	
BMI	MR Egger	14	0.830	0.582	0.179	0.079	0.071
	Weighted median	14	-0.381	0.106	0.000		
	Inverse variance weighted	14	-0.308	0.095	0.001	0.018	
WC	MR Egger	14	0.669	0.637	0.314	0.179	0.169
	Weighted median	14	-0.246	0.113	0.030		
	Inverse variance weighted	14	-0.252	0.096	0.009	0.118	
WHR	MR Egger	15	0.235	1.375	0.867	0.010	0.661
	Weighted median	15	-0.511	0.222	0.021		
	Inverse variance weighted	15	-0.375	0.213	0.079	0.014	

BMI: Body mass index; CHD: Coronary heart disease; HbA1c: Glycated hemoglobin; LDL: Low-density lipoprotein; MR: Mendelian randomization; nSNP: Number of Single-nucleotide polymorphisms; TC: Total cholesterol; T2DM: Type 2 diabetes mellitus; TG: Triglycerides; WC: Waist circumference; WHR: Waist-to-hip ratio.



Table 4 Causal association between genetically determined metabolic risk factors and type 2 diabetes mellitus


Trait	Method	nSNP	OR	95%CI	P value	Heterogeneity P	MR Egger intercept P
CHD-T2DM	MR Egger	28	0.947	0.697 - 1.288	0.733	0.000	0.815
	Weighted median	28	1.111	0.996 - 1.239	0.059		
	Inverse variance weighted	28	1.073	0.954 - 1.207	0.243	0.000	
BMI-T2DM	MR Egger	72	3.370	1.328 - 8.556	0.013	0.000	0.240
	Weighted median	72	2.622	2.164 - 3.178	0.000		
	Inverse variance weighted	72	2.046	1.374 - 3.048	0.000	0.000	
WC-T2DM	MR Egger	39	11.670	1.590 - 85.654	0.021	0.000	0.049
	Weighted median	39	2.371	1.788 - 3.145	0.000		
	Inverse variance weighted	39	1.607	0.879 - 2.938	0.123	0.000	

BMI: Body mass index; CHD: Coronary heart disease; MR: Mendelian randomization; OR: Odds ratio; nSNP: Number of single-nucleotide polymorphisms; T2DM: Type 2 diabetes mellitus; WC: Waist circumference.

medication adherence without a significant reduction in adverse clinical outcomes^[20]. A systematic review and meta-analysis also demonstrated that educational interventions improved medication adherence among adult patients diagnosed with diabetes^[20]. Moreover, structured education had a positive impact on glucose control and hypoglycemia in T2DM^[21]. However, there has been no direct study exploring the causality between original education and T2DM.

Interestingly, previous findings indicated a causal association between low educational attainment and increased risk of smoking^[22]. Also, observational studies suggested an association between smoking and risk of T2DM^[24,25]. Based on the above reviews, we made a bold assumption that education might be negatively causally associated with T2DM. As we expected, our MR results revealed that the OR (95%CI)

A**B**

C

MR leave-one-out sensitivity analysis for education on T2DM


D

Figure 1 Mendelian randomization study of the effect of education on type 2 diabetes mellitus. A: Mendelian randomization estimate for education on type 2 diabetes mellitus (T2DM); B: Effect sizes of the single-nucleotide polymorphism (SNP)-education associations [x-axis, SD (3.6 years) units] and the SNP-T2DM associations (y-axis, log odds of T2DM); C: Leave-one-out sensitivity analysis; D: Funnel plot of the causality between education and T2DM. MR: Mendelian randomization; OR: Odds ratio.

for T2DM was 0.392 (0.263-0.583) per standard deviation increase (3.6 years) in education, indicating that the genetically predicted education was negatively causally associated with T2DM. Since education was shown to be a protective factor against T2DM, and it was modifiable, longer education years among the population are recommended for T2DM prevention.

As there was no full description of the underlying mechanisms possibly connecting education to T2DM, we investigated this relationship. As for the possible mediators from education to T2DM, ten representative modifiable metabolic risk factors were chosen for further MR analysis. Our results indicated that education was causally associated with CHD, TG, BMI, WC, and WHR in the discovery phase. Yet, in the replication data set, only the results for CHD, BMI, and WC were duplicated. Therefore, there were negative causal associations between genetically determined education and CHD, BMI, and WC, which were considered to be the potential mediators between education and T2DM.

A previous study revealed that genetic predisposition towards 3.6 years of additional education was associated with a one-third lower risk of CHD, indicating that low education was a causal risk factor in the development of CHD^[26], which was in accord with our finding. Another MR analysis also suggested that there might be a negative causal effect of education on BMI^[27]. Our results provide more evidence supporting that more education years might help the public better control BMI, which is beneficial for blood glucose regulation and T2DM prevention. There was no direct causal study exploring the effect of education on WC. A randomized clinical trial revealed that nutrition therapy and a multimedia diabetes education program positively impacted achieving metabolic control goals in T2DM, including HbA1C, glucose decrease, TG, and weight loss. Yet, the WC change was still not statistically different^[28]. Another study reported that diet-related and lifestyle-related school-based education could reduce central adiposity in pre-teenagers^[29]. We analyzed whether the different effects of education on WC might be due to various educational programs or diverse populations used in the studies.

Since our MR investigation indicated that CHD, BMI, and WC might be potential mediators from less education years to increased risk of T2DM, we further evaluated whether these three potential mediators were associated with T2DM by MR analysis. Only BMI was positively associated with T2DM. In summary, BMI might serve as a mediator between education and T2DM.

BMI is a well-described risk factor for T2DM^[30], and several large-scale MR studies have addressed its positive causal association with T2DM^[31-33]. Our results also demonstrated that higher BMI could lead to higher risk of T2MD. Combined with the effects of education on BMI and T2MD, for most developing countries where the majority of population receives short education years, we recommend longer education time and a BMI control program for public T2DM prevention.

CONCLUSION

Short education was found to be associated with an increased T2DM risk. BMI might serve as a potential mediator between them.

ARTICLE HIGHLIGHTS

Research background

The causality between education and type 2 diabetes mellitus (T2DM) remains unclear.

Research motivation

In this study, a network Mendelian randomization (MR) framework was applied to determine the causality between education and T2DM from summarized genome-wide association study data.

Research objectives

We used a network MR to identify the causality between education and T2DM and the potential metabolic risk factors [coronary heart disease (CHD), total cholesterol, low-density lipoprotein, triglycerides, body mass index (BMI), waist circumference (WC), waist-to-hip ratio, fasting insulin, fasting glucose, and glycated hemoglobin] from

summarized genome-wide association study data.

Research methods

Two-sample MR and network MR were performed to obtain the causality between education-T2DM, education-mediator, and mediator-T2DM. Summary statistics from the Social Science Genetic Association Consortium (discovery data) and Neale Lab consortium (replication data) were used for education. DIAGRAM plus Metabochip consortium data were utilized for T2DM.

Research results

In the IVW method, the odds ratio (95%CI) for T2DM was 0.392 (0.263-0.583) per standard deviation increase (3.6 years) in education, without heterogeneity or horizontal pleiotropy. Education was genetically associated with CHD, triglycerides, BMI, WC, and waist-to-hip ratio in the discovery phase, yet only the results for CHD, BMI, and WC were confirmed in the replication data. Moreover, BMI was positively associated with T2DM.

Research conclusions

Short education was found to be associated with increased T2DM risk. BMI might serve as a potential mediator between them.

Research perspectives

For most developing countries, the majority of the population receive short education years. Longer education time is recommended as is a BMI control program, for public T2DM prevention.

REFERENCES

- 1 **Dall TM**, Narayan KM, Gillespie KB, Gallo PD, Blanchard TD, Solcan M, O'Grady M, Quick WW. Detecting type 2 diabetes and prediabetes among asymptomatic adults in the United States: modeling American Diabetes Association versus US Preventive Services Task Force diabetes screening guidelines. *Popul Health Metr* 2014; **12**: 12 [PMID: 24904239 DOI: 10.1186/1478-7954-12-12]
- 2 **Fletcher B**, Gulanick M, Lamendola C. Risk factors for type 2 diabetes mellitus. *J Cardiovasc Nurs* 2002; **16**: 17-23 [PMID: 11800065 DOI: 10.1097/00005082-200201000-00003]
- 3 **Vitale M**, Masulli M, Calabrese I, Rivelles AA, Bonora E, Signorini S, Perriello G, Squatrito S, Buzzetti R, Sartore G, Babini AC, Gregori G, Giordano C, Clemente G, Grioni S, Dolce P, Riccardi G, Vaccaro O; TOSCA. IT Study Group. Impact of a Mediterranean Dietary Pattern and Its Components on Cardiovascular Risk Factors, Glucose Control, and Body Weight in People with Type 2 Diabetes: A Real-Life Study. *Nutrients* 2018; **10** [PMID: 30103444 DOI: 10.3390/nu10081067]
- 4 **He X**, Li J, Wang B, Yao Q, Li L, Song R, Shi X, Zhang JA. Diabetes self-management education reduces risk of all-cause mortality in type 2 diabetes patients: a systematic review and meta-analysis. *Endocrine* 2017; **55**: 712-731 [PMID: 27837440 DOI: 10.1007/s12020-016-1168-2]
- 5 **Emdin CA**, Khera AV, Kathiresan S. Mendelian Randomization. *JAMA* 2017; **318**: 1925-1926 [PMID: 29164242 DOI: 10.1001/jama.2017.17219]
- 6 **Okbay A**, Beauchamp JP, Fontana MA, Lee JJ, Pers TH, Rietveld CA, Turley P, Chen GB, Emilsson V, Meddins SF, Oskarsson S, Pickrell JK, Thom K, Timshel P, de Vlaming R, Abdellaoui A, Ahluwalia TS, Bacelis J, Baumbach C, Bjornsdottir G, Brandsma JH, Pina Concas M, Derringer J, Furlotte NA, Galesloot TE, Giroto G, Gupta R, Hall LM, Harris SE, Hofer E, Horikoshi M, Huffman JE, Kaasik K, Kalafati IP, Karlsson R, Kong A, Lahti J, van der Lee SJ, deLeeuw C, Lind PA, Lindgren KO, Liu T, Mangino M, Marten J, Mihailov E, Miller MB, van der Most PJ, Oldmeadow C, Payton A, Pervjakova N, Peyrot WJ, Qian Y, Raitakari O, Rueedi R, Salvi E, Schmidt B, Schraut KE, Shi J, Smith AV, Poot RA, St Pourcain B, Teumer A, Thorleifsson G, Verweij N, Vuckovic D, Wellmann J, Westra HJ, Yang J, Zhao W, Zhu Z, Alizadeh BZ, Amin N, Bakshi A, Baumeister SE, Biino G, Bønnelykke K, Boyle PA, Campbell H, Cappuccio FP, Davies G, De Neve JE, Deloukas P, Demuth I, Ding J, Eibich P, Eisele L, Eklund N, Evans DM, Faul JD, Feitosa MF, Forstner AJ, Gandin I, Gunnarsson B, Halldórrsson BV, Harris TB, Heath AC, Hocking LJ, Holliday EG, Homuth G, Horan MA, Hottenga JJ, de Jager PL, Joshi PK, Jugessur A, Kaakinen MA, Kähönen M, Kanoni S, Keltigangas-Järvinen L, Kiemeney LA, Kolcic I, Koskinen S, Kraja AT, Kroh M, Kutalik Z, Latvala A, Launer LJ, Lebreton MP, Levinson DF, Lichtenstein P, Lichtner P, Liewald DC; LifeLines Cohort Study, Loukola A, Madden PA, Mägi R, Mäki-Opas T, Marioni RE, Marques-Vidal P, Meddins GA, McMahon G, Meisinger C, Meitinger T, Milaneschi Y, Milani L, Montgomery GW, Myhre R, Nelson CP, Nyholt DR, Ollier WE, Palotie A, Paternoster L, Pedersen NL, Petrovic KE, Porteous DJ, Räikkönen K, Ring SM, Robino A, Rostapshova O, Rudan I, Rustichini A, Salomaa V, Sanders AR, Sarin AP, Schmidt H, Scott RJ, Smith BH, Smith JA, Staessen JA, Steinhagen-Thiessen E, Strauch K, Terracciano A, Tobin MD, Ulivi S, Vaccargiu S, Quaye L, van Rooij FJ, Venturini C, Vinkhuyzen

- AA, Völker U, Völzke H, Vonk JM, Vozzi D, Waage J, Ware EB, Willemsen G, Attia JR, Bennett DA, Berger K, Bertram L, Bisgaard H, Boomsma DI, Borecki IB, Bültmann U, Chabris CF, Cucca F, Cusi D, Dearly IJ, Dedoussis GV, van Duijn CM, Eriksson JG, Franke B, Franke L, Gasparini P, Gejman PV, Gieger C, Grabe HJ, Gratton J, Groenen PJ, Gudnason V, van der Harst P, Hayward C, Hinds DA, Hoffmann W, Hyppönen E, Iacono WG, Jacobsson B, Järvelin MR, Jöckel KH, Kaprio J, Kardia SL, Lehtimäki T, Lehrer SF, Magnusson PK, Martin NG, McGuire M, Metspalu A, Pendleton N, Penninx BW, Perola M, Pirastu N, Pirastu M, Polasek O, Posthuma D, Power C, Province MA, Samani NJ, Schlessinger D, Schmidt R, Sørensen TI, Spector TD, Stefansson K, Thorsteinsdottir U, Thurik AR, Timpton NJ, Tiemeier H, Tung JY, Uitterlinden AG, Vitart V, Vollenweider P, Weir DR, Wilson JF, Wright AF, Conley DC, Krueger RF, Davey Smith G, Hofman A, Laibson DI, Medland SE, Meyer MN, Yang J, Johannesson M, Visscher PM, Esko T, Koellinger PD, Cesarini D, Benjamin DJ. Genome-wide association study identifies 74 loci associated with educational attainment. *Nature* 2016; **533**: 539-542 [PMID: 27225129 DOI: 10.1038/nature17671]
- 7 **Morris AP**, Voight BF, Teslovich TM, Ferreira T, Segre AV, Steinhordsdottir V, Strawbridge RJ, Khan H, Grallert H, Mahajan A, Prokopenko I, Kang HM, Dina C, Esko T, Fraser RM, Kanoni S, Kumar A, Lagou V, Langenberg C, Luan J, Lindgren CM, Müller-Nurasyid M, Pechlivanis S, Rayner NW, Scott LJ, Wiltshire S, Yengo L, Kinnunen L, Rossin EJ, Raychaudhuri S, Johnson AD, Dimas AS, Loos RJ, Vedantam S, Chen H, Florez JC, Fox C, Liu CT, Rybin D, Couper DJ, Kao WH, Li M, Cornelis MC, Kraft P, Sun Q, van Dam RM, Stringham HM, Chines PS, Fischer K, Fontanillas P, Holmen OL, Hunt SE, Jackson AU, Kong A, Lawrence R, Meyer J, Perry JR, Platou CG, Potter S, Rehnberg E, Robertson N, Sivapalaratnam S, Stančáková A, Stirrups K, Thorleifsson G, Tikkanen E, Wood AR, Almgren P, Atalay M, Benediktsson R, Bonnycastle LL, Burtt N, Carey J, Charpentier G, Crenshaw AT, Doney AS, Dorkhan M, Edkins S, Emilsson V, Eury E, Forseen T, Gertow K, Gigante B, Grant GB, Groves CJ, Guiducci C, Herder C, Hreidarsson AB, Hui J, James A, Jonsson A, Rathmann W, Klopp N, Kravic J, Krjutškov K, Langford C, Leander K, Lindholm E, Lobbens S, Männistö S, Mirza G, Mühlleisen TW, Musk B, Parkin M, Rallidis L, Saramies J, Sennblad B, Shah S, Sigurðsson G, Silveira A, Steinbach G, Thorand B, Trakalo J, Veglia F, Wennauer R, Winckler W, Zabaneh D, Campbell H, van Duijn C, Uitterlinden AG, Hofman A, Sijbrands E, Abecasis GR, Owen KR, Zeggini E, Trip MD, Forouhi NG, Syvänen AC, Eriksson JG, Peltonen L, Nöthen MM, Balkau B, Palmer CN, Lyssenko V, Tuomi T, Isomaa B, Hunter DJ, Qi L; Wellcome Trust Case Control Consortium; Meta-Analyses of Glucose and Insulin-related traits Consortium (MAGIC) Investigators; Genetic Investigation of ANthropometric Traits (GIANT) Consortium; Asian Genetic Epidemiology Network-Type 2 Diabetes (AGEN-T2D) Consortium; South Asian Type 2 Diabetes (SAT2D) Consortium, Shuldiner AR, Roden M, Barroso I, Wilsgaard T, Beilby J, Hovingh K, Price JF, Wilson JF, Rauramaa R, Lakka TA, Lind L, Dedoussis G, Njølstad I, Pedersen NL, Khaw KT, Wareham NJ, Keinanen-Kiukaanniemi SM, Saaristo TE, Korpi-Hyövälte E, Saltevo J, Laakso M, Kuusisto J, Metspalu A, Collins FS, Mohlke KL, Bergman RN, Tuomilehto J, Boehm BO, Gieger C, Hveem K, Cauchi S, Froguel P, Baldassarre D, Tremoli E, Humphries SE, Saleheen D, Danesh J, Ingelsson E, Ripatti S, Salomaa V, Erbel R, Jöckel KH, Moebus S, Peters A, Illig T, de Faire U, Hamsten A, Morris AD, Donnelly PJ, Frayling TM, Hattersley AT, Boerwinkle E, Melander O, Kathiresan S, Nilsson PM, Deloukas P, Thorsteinsdottir U, Groop LC, Stefansson K, Hu F, Pankow JS, Dupuis J, Meigs JB, Altshuler D, Boehnke M, McCarthy MI; DIAbetes Genetics Replication And Meta-analysis (DIAGRAM) Consortium. Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat Genet* 2012; **44**: 981-990 [PMID: 22885922 DOI: 10.1038/ng.2383]
- 8 **Nikpay M**, Goel A, Won HH, Hall LM, Willenborg C, Kanoni S, Saleheen D, Kyriakou T, Nelson CP, Hopewell JC, Webb TR, Zeng L, Dehghan A, Alver M, Armasu SM, Auro K, Bjonnes A, Chasman DI, Chen S, Ford I, Franceschini N, Gieger C, Grace C, Gustafsson S, Huang J, Hwang SJ, Kim YK, Kleber ME, Lau KW, Lu X, Lu Y, Lyytikäinen LP, Mihailov E, Morrison AC, Pervjakova N, Qu L, Rose LM, Salfati E, Saxena R, Scholz M, Smith AV, Tikkanen E, Uitterlinden A, Yang X, Zhang W, Zhao W, de Andrade M, de Vries PS, van Zuydam NR, Anand SS, Bertram L, Beutner F, Dedoussis G, Frossard P, Gauguier D, Goodall AH, Gottesman O, Haber M, Han BG, Huang J, Jalilzadeh S, Kessler T, König IR, Lannfelt L, Lieb W, Lind L, Lindgren CM, Lokki ML, Magnusson PK, Mallick NH, Mehra N, Meitinger T, Memon FU, Morris AP, Nieminen MS, Pedersen NL, Peters A, Rallidis LS, Rasheed A, Samuel M, Shah SH, Sinisalo J, Stirrups KE, Trompet S, Wang L, Zaman KS, Ardissino D, Boerwinkle E, Borecki IB, Bottiger EP, Buring JE, Chambers JC, Collins R, Cupples LA, Danesh J, Demuth I, Elosua R, Epstein SE, Esko T, Feitosa MF, Franco OH, Franzosi MG, Granger CB, Gu D, Gudnason V, Hall AS, Hamsten A, Harris TB, Hazen SL, Hengstenberg C, Hofman A, Ingelsson E, Iribarren C, Jukema JW, Karhunen PJ, Kim BJ, Kooner JS, Kullo JJ, Lehtimäki T, Loos RJF, Melander O, Metspalu A, März W, Palmer CN, Perola M, Quertermous T, Rader DJ, Ridker PM, Ripatti S, Roberts R, Salomaa V, Sanghera DK, Schwartz SM, Seedorf U, Stewart AF, Stott DJ, Thiery J, Zalloua PA, O'Donnell CJ, Reilly MP, Assimes TL, Thompson JR, Erdmann J, Clarke R, Watkins H, Kathiresan S, McPherson R, Deloukas P, Schunkert H, Samani NJ, Farrall M. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet* 2015; **47**: 1121-1130 [PMID: 26343387 DOI: 10.1038/ng.3396]
- 9 **Willer CJ**, Schmidt EM, Sengupta S, Peloso GM, Gustafsson S, Kanoni S, Ganna A, Chen J, Buchkovich ML, Mora S, Beckmann JS, Bragg-Gresham JL, Chang HY, Demirkiran A, Den Hertog HM, Do R, Donnelly LA, Ehret GB, Esko T, Feitosa MF, Ferreira T, Fischer K, Fontanillas P, Fraser RM, Freitag DF, Gurdasani D, Heikkilä K, Hyppönen E, Isaacs A, Jackson AU, Johansson Å,

- Johnson T, Kaakinen M, Kettunen J, Kleber ME, Li X, Luan J, Lyytikäinen LP, Magnusson PKE, Mangino M, Mihailov E, Montasser ME, Müller-Nurasyid M, Nolte IM, O'Connell JR, Palmer CD, Perola M, Petersen AK, Sanna S, Saxena R, Service SK, Shah S, Shungin D, Sidore C, Song C, Strawbridge RJ, Surakka I, Tanaka T, Teslovich TM, Thorleifsson G, Van den Herik EG, Voight BF, Volcik KA, Waite LL, Wong A, Wu Y, Zhang W, Absher D, Asiki G, Barroso I, Been LF, Bolton JL, Bonnycastle LL, Brambilla P, Burnett MS, Cesana G, Dimitriou M, Doney ASF, Döring A, Elliott P, Epstein SE, Ingi Eyjolfsson G, Gigante B, Goodarzi MO, Grallert H, Gravito ML, Groves CJ, Hallmans G, Hartikainen AL, Hayward C, Hernandez D, Hicks AA, Holm H, Hung YJ, Illig T, Jones MR, Kaleebu P, Kastelein JJP, Khaw KT, Kim E, Klopp N, Komulainen P, Kumari M, Langenberg C, Lehtimäki T, Lin SY, Lindström J, Loos RJF, Mach F, McArdle WL, Meisinger C, Mitchell BD, Müller G, Nagaraja R, Narisu N, Nieminen TVM, Nsubuga RN, Olafsson I, Ong KK, Palotie A, Papamarkou T, Pomilla C, Pouta A, Rader DJ, Reilly MP, Ridker PM, Rivadeneira F, Rudan I, Ruokonen A, Samani N, Scharnagl H, Seeley J, Silander K, Stančáková A, Stirrups K, Swift AJ, Tiret L, Uitterlinden AG, van Pelt LJ, Vedantam S, Wainwright N, Wijmenga C, Wild SH, Willemse G, Wilsgaard T, Wilson JF, Young EH, Zhao JH, Adair LS, Arveiler D, Assimes TL, Bandinelli S, Bennett F, Bochud M, Boehm BO, Boomsma DI, Borecki IB, Bornstein SR, Bovet P, Burnier M, Campbell H, Chakravarti A, Chambers JC, Chen YI, Collins FS, Cooper RS, Danesh J, Dedoussis G, de Faire U, Feranil AB, Ferrières J, Ferrucci L, Freimer NB, Gieger C, Groop LC, Gudnason V, Gyllensten U, Hamsten A, Harris TB, Hingorani A, Hirschhorn JN, Hofman A, Hovingh GK, Hsiung CA, Humphries SE, Hunt SC, Hveem K, Iribarren C, Järvelin MR, Jula A, Kähönen M, Kaprio J, Kesäniemi A, Kivimaki M, Kooper JS, Koudstaal PJ, Krauss RM, Kuh D, Kuusisto J, Kyvik KO, Laakso M, Lakka TA, Lind L, Lindgren CM, Martin NG, März W, McCarthy MI, McKenzie CA, Meneton P, Metspalu A, Moilanen L, Morris AD, Munroe PB, Njølstad I, Pedersen NL, Power C, Pramstaller PP, Price JF, Psaty BM, Quertermous T, Rauramaa R, Saleheen D, Salomaa V, Sanghera DK, Saramies J, Schwarz PEH, Sheu WH, Shuldiner AR, Siegbahn A, Spector TD, Stefansson K, Strachan DP, Tayo BO, Tremoli E, Tuomilehto J, Uusitupa M, van Duijn CM, Vollenweider P, Wallentin L, Wareham NJ, Whitfield JB, Wolfenbuttel BHR, Ordovas JM, Boerwinkle E, Palmer CNA, Thorsteinsdóttir U, Chasman DI, Rotter JI, Franks PW, Ripatti S, Cupples LA, Sandhu MS, Rich SS, Boehnke M, Deloukas P, Kathiresan S, Mohlke KL, Ingelsson E, Abecasis GR; Global Lipids Genetics Consortium. Discovery and refinement of loci associated with lipid levels. *Nat Genet* 2013; **45**: 1274-1283 [PMID: 24097068 DOI: 10.1038/ng.2797]
- 10 **Locke AE**, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J, Croteau-Chonka DC, Esko T, Fall T, Ferreira T, Gustafsson S, Kutalik Z, Luan J, Mägi R, Randall JC, Winkler TW, Wood AR, Workalemahu T, Faul JD, Smith JA, Zhao JH, Zhao W, Chen J, Fehrman R, Hedman ÅK, Karjalainen J, Schmidt EM, Absher D, Amin N, Anderson D, Beckman M, Bolton JL, Bragg-Gresham JL, Buyske S, Demirkhan A, Deng G, Ehret GB, Feenstra B, Feitosa MF, Fischer K, Goel A, Gong J, Jackson AU, Kanoni S, Kleber ME, Kristiansson K, Lim U, Lotay V, Mangino M, Leach IM, Medina-Gomez C, Medland SE, Nalls MA, Palmer CD, Pasko D, Pechlivanis S, Peters MJ, Prokopenko I, Shungin D, Stančáková A, Strawbridge RJ, Sung YJ, Tanaka T, Teumer A, Trompet S, van der Laan SW, van Setten J, Van Vliet-Ostaptchouk JV, Wang Z, Yengo L, Zhang W, Isaacs A, Albrecht E, Ärnlöv J, Arscott GM, Attwood AP, Bandinelli S, Barrett A, Bas IN, Bellis C, Bennett AJ, Berne C, Blagieva R, Blüher M, Böhringer S, Bonnycastle LL, Böttcher Y, Boyd HA, Bruinenberg M, Caspersen IH, Chen YI, Clarke R, Daw EW, de Craen AJM, Delgado G, Dimitriou M, Doney ASF, Eklund N, Estrada K, Eury E, FolkerSEN L, Fraser RM, Garcia ME, Geller F, Giedraitis V, Gigante B, Go AS, Golay A, Goodall AH, Gordon SD, Gorski M, Grabe HJ, Grallert H, Grammer TB, Gräßler J, Grönberg H, Groves CJ, Gusto G, Haessler J, Hall P, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hengstenberg C, Holmen O, Hottenga JJ, James AL, Jeff JM, Johansson Å, Jolley J, Juliusdottir T, Kinnunen L, Koenig W, Koskenvuo M, Kratzer W, Laitinen J, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindström J, Lo KS, Lobbens S, Lorbeer R, Lu Y, Mach F, Magnusson PKE, Mahajan A, McArdle WL, McLachlan S, Menni C, Merger S, Mihailov E, Milani L, Moayyeri A, Monda KL, Morken MA, Mulas A, Müller G, Müller-Nurasyid M, Musk AW, Nagaraja R, Nöthen MM, Nolte IM, Pilz S, Rayner NW, Renstrom F, Rettig R, Ried JS, Ripke S, Robertson NR, Rose LM, Sanna S, Scharnagl H, Scholtens S, Schumacher FR, Scott WR, Seufferlein T, Shi J, Smith AV, Smolonska J, Stanton AV, Steinhorsdóttir V, Stirrups K, Stringham HM, Sundström J, Swertz MA, Swift AJ, Syvänen AC, Tan ST, Tayo BO, Thorand B, Thorleifsson G, Tyrer JP, Uh HW, Vandenput L, Verhulst FC, Vermeulen SH, Verweij N, Vonk JM, Waite LL, Warren HR, Waterworth D, Weedon MN, Wilkens LR, Willenborg C, Wilsgaard T, Wojczynski MK, Wong A, Wright AF, Zhang Q; LifeLines Cohort Study, Brennan EP, Choi M, Dastani Z, Drong AW, Eriksson P, Franco-Cereceda A, Gådin JR, Gharavi AG, Goddard ME, Handsaker RE, Huang J, Karpe F, Kathiresan S, Keildson S, Kiryluk K, Kubo M, Lee JY, Liang L, Lifton RP, Ma B, McCarroll SA, McKnight AJ, Min JL, Moffatt MF, Montgomery GW, Murabito JM, Nicholson G, Nyholt DR, Okada Y, Perry JRB, Dorajoo R, Reinmaa E, Salem RM, Sandholm N, Scott RA, Stolk L, Takahashi A, Tanaka T, van 't Hooft FM, Vinkhuyzen AAE, Westra HJ, Zheng W, Zondervan KT; ADIPOGen Consortium; AGEN-BMI Working Group; CARDIOGRAMplusC4D Consortium; CKDGen Consortium; GLGC; ICBP; MAGIC Investigators; MuTHER Consortium; MiGen Consortium; PAGE Consortium; ReproGen Consortium; GENIE Consortium; International Endogene Consortium, Heath AC, Arveiler D, Bakker SJL, Beilby J, Bergman RN, Blangero J, Bovet P, Campbell H, Caulfield MJ, Cesana G, Chakravarti A, Chasman DI, Chines PS, Collins FS, Crawford DC, Cupples LA, Cusi D, Danesh J, de Faire U, den Ruijter HM,

- Dominiczak AF, Erbel R, Erdmann J, Eriksson JG, Farrall M, Felix SB, Ferrannini E, Ferrières J, Ford I, Forouhi NG, Forrester T, Franco OH, Gansevoort RT, Gejman PV, Gieger C, Gottesman O, Gudnason V, Gyllensten U, Hall AS, Harris TB, Hattersley AT, Hicks AA, Hindorff LA, Hingorani AD, Hofman A, Homuth G, Hovingh GK, Humphries SE, Hunt SC, Hyppönen E, Illig T, Jacobs KB, Jarvelin MR, Jöckel KH, Johansen B, Jousilahti P, Jukema JW, Jula AM, Kaprio J, Kastelein JJP, Keinanen-Kiukaanniemi SM, Kiemeneij LA, Knekt P, Kooner JS, Kooperberg C, Kovacs P, Kraja AT, Kumari M, Kuusisto J, Lakka TA, Langenberg C, Marchand LL, Lehtimäki T, Lysenko V, Männistö S, Marette A, Matise TC, McKenzie CA, McKnight B, Moll FL, Morris AD, Morris AP, Murray JC, Nelis M, Ohlsson C, Oldehinkel AJ, Ong KK, Madden PAF, Pasterkamp G, Peden JF, Peters A, Postma DS, Pramstaller PP, Price JF, Qi L, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ridker PM, Rioux JD, Ritchie MD, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schunkert H, Schwarz PEH, Sever P, Shuldiner AR, Sinisalo J, Stolk RP, Strauch K, Tönjes A, Tréguoët DA, Tremblay A, Tremoli E, Virtamo J, Vohl MC, Völker U, Waeber G, Willemse G, Witteman JC, Zillikens MC, Adair LS, Amouyel P, Assimes TL, Bochud M, Boehm BO, Boerwinkle E, Bornstein SR, Bottiger EP, Bouchard C, Cauchi S, Chambers JC, Chanock SJ, Cooper RS, de Bakker PIW, Dedoussis G, Ferrucci L, Franks PW, Froguel P, Groop LC, Haiman CA, Hamsten A, Hui J, Hunter DJ, Hveem K, Kaplan RC, Kivimaki M, Kuh D, Laakso M, Liu Y, Martin NG, März W, Melbye M, Metspalu A, Moebus S, Munroe PB, Njølstad I, Oostra BA, Palmer CNA, Pedersen NL, Perola M, Pérusse L, Peters U, Power C, Quertermous T, Rauramaa R, Rivadeneira F, Saaristo TE, Saleheen D, Sattar N, Schadt EE, Schlessinger D, Slagboom PE, Snieder H, Spector TD, Thorsteinsdóttir U, Stumvoll M, Tuomilehto J, Uitterlinden AG, Uusitupa M, van der Harst P, Walker M, Wallachofski H, Wareham NJ, Watkins H, Weir DR, Wichmann HE, Wilson JF, Zanen P, Borecki IB, Deloukas P, Fox CS, Heid IM, O'Connell JR, Strachan DP, Stefansson K, van Duijn CM, Abecasis GR, Franke L, Frayling TM, McCarthy MI, Visscher PM, Scherag A, Willer CJ, Boehnke M, Mohlke KL, Lindgren CM, Beckmann JS, Barroso I, North KE, Ingelsson E, Hirschhorn JN, Loos RJF, Speliotes EK. Genetic studies of body mass index yield new insights for obesity biology. *Nature* 2015; **518**: 197-206 [PMID: 25673413 DOI: 10.1038/nature14177]
- 11 **Shungin D**, Winkler TW, Croteau-Chonka DC, Ferreira T, Locke AE, Mägi R, Strawbridge RJ, Pers TH, Fischer K, Justice AE, Workalemahu T, Wu JMW, Buchkovich ML, Heard-Costa NL, Roman TS, Drong AW, Song C, Gustafsson S, Day FR, Esko T, Fall T, Kutalik Z, Luan J, Randall JC, Scherag A, Vedantam S, Wood AR, Chen J, Fehrman R, Karjalainen J, Kahali B, Liu CT, Schmidt EM, Absher D, Amin N, Anderson D, Beekman M, Bragg-Gresham JL, Buyske S, Demirkan A, Ehret GB, Feitosa MF, Goel A, Jackson AU, Johnson T, Kleber ME, Kristiansson K, Mangino M, Leach IM, Medina-Gomez C, Palmer CD, Pasko D, Peclivanis S, Peters MJ, Prokopenko I, Stančáková A, Sung YJ, Tanaka T, Teumer A, Van Vliet-Ostaptchouk JV, Yengo L, Zhang W, Albrecht E, Ärnööv J, Arscott GM, Bandinelli S, Barrett A, Bellis C, Bennett AJ, Berne C, Blüher M, Böhringer S, Bonnet F, Böttcher Y, Bruinenberg M, Carba DB, Caspersen IH, Clarke R, Daw EW, Deelen J, Deelman E, Delgado G, Doney AS, Eklund N, Erdos MR, Estrada K, Eury E, Friedrich N, Garcia ME, Giedraitis V, Gigante B, Go AS, Golay A, Grallert H, Grammer TB, Gräßler J, Grewal J, Groves CJ, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heikkilä K, Herzig KH, Helmer Q, Hillege HL, Holmen O, Hunt SC, Isaacs A, Ittermann T, James AL, Johansson I, Juliusdottir T, Kalafati IP, Kinnunen L, Koenig W, Kooner IK, Kratzer W, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindström J, Lobbens S, Lorentzon M, Mach F, Magnusson PK, Mahajan A, McArdle WL, Menni C, Merger S, Mihailov E, Milani L, Mills R, Moayyeri A, Monda KL, Mooijaart SP, Mühlleisen TW, Mulas A, Müller G, Müller-Nurasyid M, Nagaraja R, Nalls MA, Narisu N, Glorioso N, Nolte IM, Olden M, Rayner NW, Renstrom F, Ried JS, Robertson NR, Rose LM, Sanna S, Scharnagl H, Scholtens S, Sennblad B, Seufferlein T, Siltani CM, Smith AV, Stirrups K, Stringham HM, Sundström J, Swertz MA, Swift AJ, Syvänen AC, Tayo BO, Thorand B, Thorleifsson G, Tomaschitz A, Troffa C, van Oort FV, Verweij N, Vonk JM, Waite LL, Wennauer R, Wilsgaard T, Wojczynski MK, Wong A, Zhang Q, Zhao JH, Brennan EP, Choi M, Eriksson P, Folkersen L, Franco-Cereceda A, Gharavi AG, Hedman ÅK, Hirvonen MF, Huang J, Kanoni S, Karpe F, Keildson S, Kiryluk K, Liang L, Lifton RP, Ma B, McKnight AJ, McPherson R, Metspalu A, Min JL, Moffatt MF, Montgomery GW, Murabito JM, Nicholson G, Nyholt DR, Olsson C, Perry JR, Reinmaa E, Salem RM, Sandholm N, Schadt EE, Scott RA, Stolk L, Vallejo EE, Westra HJ, Zondervan KT; ADIPOGen Consortium; CARDIOGRAMplusC4D Consortium; CKDGen Consortium; GEFOS Consortium; GENIE Consortium; GLGC; ICBP; International Endogene Consortium; LifeLines Cohort Study; MAGIC Investigators; MuTHER Consortium; PAGE Consortium; ReproGen Consortium, Amouyel P, Arveiler D, Bakker SJ, Beilby J, Bergman RN, Blangero J, Brown MJ, Burnier M, Campbell H, Chakravarti A, Chines PS, Claudi-Boehm S, Collins FS, Crawford DC, Danesh J, de Faire U, de Geus EJ, Dörr M, Erbel R, Eriksson JG, Farrall M, Ferrannini E, Ferrières J, Forouhi NG, Forrester T, Franco OH, Gansevoort RT, Gieger C, Gudnason V, Haiman CA, Harris TB, Hattersley AT, Heliövaara M, Hicks AA, Hingorani AD, Hoffmann W, Hofman A, Homuth G, Humphries SE, Hyppönen E, Illig T, Jarvelin MR, Johansen B, Jousilahti P, Jula AM, Kaprio J, Kee F, Keinanen-Kiukaanniemi SM, Kooner JS, Kooperberg C, Kovacs P, Kraja AT, Kumari M, Kuulasmaa K, Kuusisto J, Lakka TA, Langenberg C, Le Marchand L, Lehtimäki T, Lysenko V, Männistö S, Marette A, Matise TC, McKenzie CA, McKnight B, Musk AW, Möhlenkamp S, Morris AD, Nelis M, Ohlsson C, Oldehinkel AJ, Ong KK, Palmer LJ, Penninx BW, Peters A, Pramstaller PP, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ridker PM, Ritchie MD, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schwarz PE, Shuldiner AR, Staessen JA, Steinhorsdottir V, Stolk RP, Strauch K,

Tönjes A, Tremblay A, Tremoli E, Vohl MC, Völker U, Vollenweider P, Wilson JF, Witteman JC, Adair LS, Bochud M, Boehm BO, Bornstein SR, Bouchard C, Cauchi S, Caulfield MJ, Chambers JC, Chasman DI, Cooper RS, Dedoussis G, Ferrucci L, Froguel P, Grabe HJ, Hamsten A, Hui J, Hveem K, Jöckel KH, Kivimaki M, Kuh D, Laakso M, Liu Y, März W, Munroe PB, Njolstad I, Oostra BA, Palmer CN, Pedersen NL, Perola M, Péruisse L, Peters U, Power C, Quertermous T, Rauramaa R, Rivadeneira F, Saaristo TE, Saleheen D, Sinisalo J, Slagboom PE, Snieder H, Spector TD, Stefansson K, Stumvoll M, Tuomilehto J, Uitterlinden AG, Uusitupa M, van der Harst P, Veronesi G, Walker M, Wareham NJ, Watkins H, Wichmann HE, Abecasis GR, Assimes TL, Berndt SI, Boehnke M, Borecki IB, Deloukas P, Franke L, Frayling TM, Groop LC, Hunter DJ, Kaplan RC, O'Connell JR, Qi L, Schlessinger D, Strachan DP, Thorsteinsdottir U, van Duijn CM, Willer CJ, Visscher PM, Yang J, Hirschhorn JN, Zillikens MC, McCarthy MI, Speliotes EK, North KE, Fox CS, Barroso I, Franks PW, Ingelsson E, Heid IM, Loos RJ, Cupples LA, Morris AP, Lindgren CM, Mohlke KL. New genetic loci link adipose and insulin biology to body fat distribution. *Nature* 2015; **518**: 187-196 [PMID: 25673412 DOI: 10.1038/nature14132]

- 12 **Randall JC**, Winkler TW, Kutalik Z, Berndt SI, Jackson AU, Monda KL, Kilpeläinen TO, Esko T, Mägi R, Li S, Workalemahu T, Feitosa MF, Croteau-Chonka DC, Day FR, Fall T, Ferreira T, Gustafsson S, Locke AE, Mathieson I, Scherag A, Vedantam S, Wood AR, Liang L, Steinhorsdottir V, Thorleifsson G, Dermitzakis ET, Dimas AS, Karpe F, Min JL, Nicholson G, Clegg DJ, Person T, Krohn JP, Bauer S, Buechler C, Eisinger K, DIAGRAM Consortium, Bonnefond A, Froguel P; MAGIC Investigators, Hottenga JJ, Prokopenko I, Waite LL, Harris TB, Smith AV, Shuldiner AR, McArdle WL, Caulfield MJ, Munroe PB, Grönberg H, Chen YD, Li G, Beckmann JS, Johnson T, Thorsteinsdottir U, Teder-Laving M, Khaw KT, Wareham NJ, Zhao JH, Amin N, Oostra BA, Kraja AT, Province MA, Cupples LA, Heard-Costa NL, Kaprio J, Ripatti S, Surakka I, Collins FS, Saramies J, Tuomilehto J, Jula A, Salomaa V, Erdmann J, Hengstenberg C, Loley C, Schunkert H, Lamina C, Wichmann HE, Albrecht E, Gieger C, Hicks AA, Johansson A, Pramstaller PP, Kathiresan S, Speliotes EK, Penninx B, Hartikainen AL, Jarvelin MR, Gyllensten U, Boomsma DI, Campbell H, Wilson JF, Chanock SJ, Farrall M, Goel A, Medina-Gomez C, Rivadeneira F, Estrada K, Uitterlinden AG, Hofman A, Zillikens MC, den Heijer M, Kiemeneij LA, Maschio A, Hall P, Tyrer J, Teumer A, Völzke H, Kovacs P, Tönjes A, Mangino M, Spector TD, Hayward C, Rudan I, Hall AS, Samani NJ, Attwood AP, Sambrook JG, Hung J, Palmer LJ, Lokki ML, Sinisalo J, Boucher G, Huikuri H, Lorentzon M, Ohlsson C, Eklund N, Eriksson JG, Barlassina C, Rivolta C, Nolte IM, Snieder H, Van der Klauw MM, Van Vliet-Ostaptchouk JV, Gejman PV, Shi J, Jacobs KB, Wang Z, Bakker SJ, Mateo Leach I, Navis G, van der Harst P, Martin NG, Medland SE, Montgomery GW, Yang J, Chasman DI, Ridker PM, Rose LM, Lehtimäki T, Raitakari O, Absher D, Iribarren C, Basart H, Hovingh KG, Hyppönen E, Power C, Anderson D, Beilby JP, Hui J, Jolley J, Sager H, Bornstein SR, Schwarz PE, Kristiansson K, Perola M, Lindström J, Swift AJ, Uusitupa M, Atalay M, Lakka TA, Rauramaa R, Bolton JL, Fowkes G, Fraser RM, Price JF, Fischer K, Krjutå Kov K, Metspalu A, Mihailov E, Langenberg C, Luan J, Ong KK, Chines PS, Keinanen-Kiukaanniemi SM, Saaristo TE, Edkins S, Franks PW, Hallmans G, Shungin D, Morris AD, Palmer CN, Erbel R, Moebus S, Nöthen MM, Peclivanis S, Hveem K, Narisu N, Hamsten A, Humphries SE, Strawbridge RJ, Tremoli E, Grallert H, Thorand B, Illig T, Koenig W, Müller-Nurasyid M, Peters A, Boehm BO, Kleber ME, März W, Winkelmann BR, Kuusisto J, Laakso M, Arveiler D, Cesana G, Kuulasmaa K, Virtamo J, Yarnell JW, Kuh D, Wong A, Lind L, de Faire U, Gigante B, Magnusson PK, Pedersen NL, Dedoussis G, Dimitriou M, Kolovou G, Kanoni S, Stirrups K, Bonnycastle LL, Njølstad I, Wilsgaard T, Ganna A, Rehnberg E, Hingorani A, Kivimaki M, Kumari M, Assimes TL, Barroso I, Boehnke M, Borecki IB, Deloukas P, Fox CS, Frayling T, Groop LC, Haritunians T, Hunter D, Ingelsson E, Kaplan R, Mohlke KL, O'Connell JR, Schlessinger D, Strachan DP, Stefansson K, van Duijn CM, Abecasis GR, McCarthy MI, Hirschhorn JN, Qi L, Loos RJ, Lindgren CM, North KE, Heid IM. Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. *PLoS Genet* 2013; **9**: e1003500 [PMID: 23754948 DOI: 10.1371/journal.pgen.1003500]
- 13 **Dupuis J**, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, Wheeler E, Glazer NL, Bouatia-Naji N, Gloyn AL, Lindgren CM, Mägi R, Morris AP, Randall J, Johnson T, Elliott P, Rybin D, Thorleifsson G, Steinhorsdottir V, Henneman P, Grallert H, Dehghan A, Hottenga JJ, Franklin CS, Navarro P, Song K, Goel A, Perry JR, Egan JM, Lajunen T, Grarup N, Sparso T, Doney A, Voight BF, Stringham HM, Li M, Kanoni S, Shrader P, Cavalcanti-Proença C, Kumari M, Qi L, Timpson NJ, Gieger C, Zabena C, Rocheleau G, Ingelsson E, An P, O'Connell J, Luan J, Elliott A, McCarroll SA, Payne F, Roccasecca RM, Pattou F, Sethupathy P, Ardlie K, Ariyurek Y, Balkau B, Barter P, Beilby JP, Ben-Shlomo Y, Benediktsson R, Bennett AJ, Bergmann S, Bochud M, Boerwinkle E, Bonnefond A, Bonnycastle LL, Borch-Johnsen K, Böttcher Y, Brunner E, Bumpstead SJ, Charpentier G, Chen YD, Chines P, Clarke R, Coin LJ, Cooper MN, Cornelis M, Crawford G, Crisponi L, Day IN, de Geus EJ, Delplanque J, Dina C, Erdos MR, Fedson AC, Fischer-Rosinsky A, Forouhi NG, Fox CS, Frants R, Franzosi MG, Galan P, Goodarzi MO, Graessler J, Groves CJ, Grundy S, Gwilliam R, Gyllensten U, Hadjadj S, Hallmans G, Hammond N, Han X, Hartikainen AL, Hassanali N, Hayward C, Heath SC, Hercberg S, Herder C, Hicks AA, Hillman DR, Hingorani AD, Hofman A, Hui J, Hung J, Isomaa B, Johnson PR, Jørgensen T, Jula A, Kaakinen M, Kaprio J, Kesaniemi YA, Kivimaki M, Knight B, Koskeni S, Kovacs P, Kyvik KO, Lathrop GM, Lawlor DA, Le Bacquer O, Lecoeur C, Li Y, Lyssenko V, Mahley R, Mangino M, Manning AK, Martinez-Larrad MT, McAtee JB, McCulloch LJ, McPherson R, Meisinger C, Melzer D, Meyre D, Mitchell BD,

- Morken MA, Mukherjee S, Naitza S, Narisu N, Neville MJ, Oostra BA, Orrù M, Pakyz R, Palmer CN, Paolisso G, Pattaro C, Pearson D, Peden JF, Pedersen NL, Perola M, Pfeiffer AF, Pichler I, Polasek O, Posthuma D, Potter SC, Pouta A, Province MA, Psaty BM, Rathmann W, Rayner NW, Rice K, Ripatti S, Rivadeneira F, Roden M, Rolandsson O, Sandbaek A, Sandhu M, Sanna S, Sayer AA, Scheet P, Scott LJ, Seedorf U, Sharp SJ, Shields B, Sigurethsson G, Sijbrands EJ, Silveira A, Simpson L, Singleton A, Smith NL, Sovio U, Swift A, Syddall H, Syvänen AC, Tanaka T, Thorand B, Tchet J, Tönjes A, Tuomi T, Uitterlinden AG, van Dijk KW, van Hoek M, Varma D, Visvikis-Siest S, Vitart V, Vogelzangs N, Waeber G, Wagner PJ, Walley A, Walters GB, Ward KL, Watkins H, Weedon MN, Wild SH, Willemsen G, Witteman JC, Yarnell JW, Zeggini E, Zelenika D, Zethelius B, Zhai G, Zhao JH, Zillikens MC; DIAGRAM Consortium; GIANT Consortium; Global BPgen Consortium, Borecki IB, Loos RJ, Meneton P, Magnusson PK, Nathan DM, Williams GH, Hattersley AT, Silander K, Salomaa V, Smith GD, Bornstein SR, Schwarz P, Spranger J, Karpe F, Shuldiner AR, Cooper C, Dedoussis GV, Serrano-Ríos M, Morris AD, Lind L, Palmer LJ, Hu FB, Franks PW, Ebrahim S, Marmot M, Kao WH, Pankow JS, Sampson MJ, Kuusisto J, Laakso M, Hansen T, Pedersen O, Pramstaller PP, Wichmann HE, Illig T, Rudan I, Wright AF, Stumvoll M, Campbell H, Wilson JF; Anders Hamsten on behalf of Procardis Consortium; MAGIC investigators, Bergman RN, Buchanan TA, Collins FS, Mohlke KL, Tuomilehto J, Valle TT, Altshuler D, Rotter JI, Siscovick DS, Penninx BW, Boomsma DI, Deloukas P, Spector TD, Frayling TM, Ferrucci L, Kong A, Thorsteinsdóttir U, Stefansson K, van Duijn CM, Aulchenko YS, Cao A, Scuteri A, Schlessinger D, Uda M, Ruckonen A, Jarvelin MR, Waterworth DM, Vollenweider P, Peltonen L, Mooser V, Abecasis GR, Wareham NJ, Sladek R, Froguel P, Watanabe RM, Meigs JB, Groop L, Boehnke M, McCarthy MI, Florez JC, Barroso I. New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet* 2010; **42**: 105-116 [PMID: 20081858 DOI: 10.1038/ng.520]
- 14 Manning AK, Hivert MF, Scott RA, Grimsby JL, Bouatia-Naji N, Chen H, Rybin D, Liu CT, Bielak LF, Prokopenko I, Amin N, Barnes D, Cadby G, Hottenga JJ, Ingelsson E, Jackson AU, Johnson T, Kanoni S, Ladenbäck C, Lagou V, Lahti J, Lecoeur C, Liu Y, Martinez-Larrad MT, Montasser ME, Navarro P, Perry JR, Rasmussen-Torvik LJ, Salo P, Sattar N, Shungin D, Strawbridge RJ, Tanaka T, van Duijn CM, An P, de Andrade M, Andrews JS, Aspelund T, Atalay M, Aulchenko Y, Balkau B, Bandinelli S, Beckmann JS, Beilby JP, Bellis C, Bergman RN, Blangero J, Boban M, Boehnke M, Boerwinkle E, Bonnycastle LL, Boomsma DI, Borecki IB, Böttcher Y, Bouchard C, Brunner E, Budimir D, Campbell H, Carlson O, Chines PS, Clarke R, Collins FS, Corbatón-Anchuelo A, Couper D, de Faire U, Dedoussis GV, Deloukas P, Dimitriou M, Egan JM, Eiriksdóttir G, Erdos MR, Eriksson JG, Eury E, Ferrucci L, Ford I, Forouhi NG, Fox CS, Franzosi MG, Franks PW, Frayling TM, Froguel P, Galan P, de Geus E, Gigante B, Glazer NL, Goel A, Groop L, Gudnason V, Hallmans G, Hamsten A, Hansson O, Harris TB, Hayward C, Heath S, Hercberg S, Hicks AA, Hingorani A, Hofman A, Hui J, Hung J, Jarvelin MR, Jhun MA, Johnson PC, Jukema JW, Jula A, Kao WH, Kaprio J, Kardia SL, Keinanen-Kiukaanniemi S, Kivimaki M, Kolcic I, Kovacs P, Kumari M, Kuusisto J, Kyvik KO, Laakso M, Lakka T, Lannfelt L, Lathrop GM, Launer LJ, Leander K, Li G, Lind L, Lindstrom J, Lobbens S, Loos RJ, Luan J, Lyssenko V, Mägi R, Magnusson PK, Marmot M, Meneton P, Mohlke KL, Mooser V, Morken MA, Miljkovic I, Narisu N, O'Connell J, Ong KK, Oostra BA, Palmer LJ, Palotie A, Pankow JS, Peden JF, Pedersen NL, Pehlic M, Peltonen L, Penninx B, Pericic M, Perola M, Perusse L, Peyser PA, Polasek O, Pramstaller PP, Province MA, Räikkönen K, Rauramaa R, Rehnberg E, Rice K, Rotter JI, Rudan I, Ruokonen A, Saaristo T, Sabater-Lleal M, Salomaa V, Savage DB, Saxena R, Schwarz P, Seedorf U, Sennblad B, Serrano-Ríos M, Shuldiner AR, Sijbrands EJ, Siscovick DS, Smit JH, Small KS, Smith NL, Smith AV, Stančáková A, Stirrups K, Stumvoll M, Sun YY, Swift AJ, Tönjes A, Tuomilehto J, Trompet S, Uitterlinden AG, Uusitupa M, Vikström M, Vitart V, Vohl MC, Voight BF, Vollenweider P, Waeber G, Waterworth DM, Watkins H, Wheeler E, Widén E, Wild SH, Willemsen G, Wilson JF, Witteman JC, Wright AF, Yaghootkar H, Zelenika D, Zemunik T, Zgaga L; DIAbetes Genetics Replication And Meta-analysis (DIAGRAM) Consortium; Multiple Tissue Human Expression Resource (MUTHER) Consortium, Wareham NJ, McCarthy MI, Barroso I, Watanabe RM, Florez JC, Dupuis J, Meigs JB, Langenberg C. A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. *Nat Genet* 2012; **44**: 659-669 [PMID: 22581228 DOI: 10.1038/ng.2274]
- 15 Soranzo N, Sanna S, Wheeler E, Gieger C, Radke D, Dupuis J, Bouatia-Naji N, Langenberg C, Prokopenko I, Stolerman E, Sandhu MS, Heaney MM, Devaney JM, Reilly MP, Ricketts SL, Stewart AF, Voight BF, Willenborg C, Wright B, Altshuler D, Arking D, Balkau B, Barnes D, Boerwinkle E, Böhm B, Bonnefond A, Bonnycastle LL, Boomsma DI, Bornstein SR, Böttcher Y, Bumpstead S, Burnett-Miller MS, Campbell H, Cao A, Chambers J, Clark R, Collins FS, Coresh J, de Geus EJ, Dei M, Deloukas P, Döring A, Egan JM, Elosua R, Ferrucci L, Forouhi N, Fox CS, Franklin C, Franzosi MG, Gallina S, Goel A, Graessler J, Grallert H, Greinacher A, Hadley D, Hall A, Hamsten A, Hayward C, Heath S, Herder C, Homuth G, Hottenga JJ, Hunter-Merrill R, Illig T, Jackson AU, Jula A, Kleber M, Knouff CW, Kong A, Kooper J, Köttingen A, Kovacs P, Krohn K, Künnel B, Kuusisto J, Laakso M, Lathrop M, Lecoeur C, Li M, Li M, Loos RJ, Luan J, Lyssenko V, Mägi R, Magnusson PK, Mälarstig A, Mangino M, Martínez-Larrad MT, März W, McArdle WL, McPherson R, Meisinger C, Meitinger T, Melander O, Mohlke KL, Mooser VE, Morken MA, Narisu N, Nathan DM, Nauck M, O'Donnell C, Oexle K, Olla N, Pankow JS, Payne F, Peden JF, Pedersen NL, Peltonen L, Perola M, Polasek O, Porcu E, Rader DJ, Rathmann W, Ripatti S, Rocheleau G, Roden M, Rudan I, Salomaa V,

- Saxena R, Schlessinger D, Schunkert H, Schwarz P, Seedorf U, Selvin E, Serrano-Ríos M, Shrader P, Silveira A, Siscovich D, Song K, Spector TD, Stefansson K, Steinhorsdóttir V, Strachan DP, Strawbridge R, Stumvoll M, Surakka I, Swift AJ, Tanaka T, Teumer A, Thorleifsson G, Thorsteinsdóttir U, Tönjes A, Usala G, Vitart V, Völzke H, Wallaschofski H, Waterworth DM, Watkins H, Wichmann HE, Wild SH, Willemsen G, Williams GH, Wilson JF, Winkelmann J, Wright AF; WTCCC, Zabene C, Zhao JH, Epstein SE, Erdmann J, Hakonarson HH, Kathiresan S, Khaw KT, Roberts R, Samani NJ, Fleming MD, Sladek R, Abecasis G, Boehnke M, Froguel P, Groop L, McCarthy MI, Kao WH, Florez JC, Uda M, Wareham NJ, Barroso I, Meigs JB. Common variants at 10 genomic loci influence hemoglobin A₁(C) levels via glycemic and nonglycemic pathways. *Diabetes* 2010; **59**: 3229-3239 [PMID: 20858683 DOI: 10.2337/db10-0502]
- 16 **Hartwig FP**, Davies NM, Hemani G, Davey Smith G. Two-sample Mendelian randomization: avoiding the downsides of a powerful, widely applicable but potentially fallible technique. *Int J Epidemiol* 2016; **45**: 1717-1726 [PMID: 28338968 DOI: 10.1093/ije/dyx028]
- 17 **Hemani G**, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, Laurin C, Burgess S, Bowden J, Langdon R, Tan VY, Yarmolinsky J, Shihab HA, Timpton NJ, Evans DM, Relton C, Martin RM, Davey Smith G, Gaunt TR, Haycock PC. The MR-Base platform supports systematic causal inference across the human genome. *Elife* 2018; **7** [PMID: 29846171 DOI: 10.7554/elife.34408]
- 18 **Morgan RG**. Network Mendelian Randomization Study Design to Assess Factors Mediating the Causal Link Between Telomere Length and Heart Disease. *Circ Res* 2017; **121**: 200-202 [PMID: 28729448 DOI: 10.1161/CIRCRESAHA.117.311387]
- 19 **Weisman A**, Fazli GS, Johns A, Booth GL. Evolving Trends in the Epidemiology, Risk Factors, and Prevention of Type 2 Diabetes: A Review. *Can J Cardiol* 2018; **34**: 552-564 [PMID: 29731019 DOI: 10.1016/j.cjca.2018.03.002]
- 20 **Marušić S**, Meliš P, Lucijanić M, Grgurević I, Turčić P, Neto PRO, Bilić-Čurčić I. Impact of pharmacotherapeutic education on medication adherence and adverse outcomes in patients with type 2 diabetes mellitus: a prospective, randomized study. *Croat Med J* 2018; **59**: 290-297 [PMID: 30610771 DOI: 10.3325/cmj.2018.59.290]
- 21 **Tan JP**, Cheng KKF, Siah RC. A systematic review and meta-analysis on the effectiveness of education on medication adherence for patients with hypertension, hyperlipidaemia and diabetes. *J Adv Nurs* 2019; **75**: 2478-2494 [PMID: 30993749 DOI: 10.1111/jan.14025]
- 22 **Yorke E**, Atiase Y. Impact of structured education on glucose control and hypoglycaemia in Type-2 diabetes: a systematic review of randomized controlled trials. *Ghana Med J* 2018; **52**: 41-60 [PMID: 30013260 DOI: 10.4314/gmj.v52i1.8]
- 23 **Gage SH**, Bowden J, Davey Smith G, Munafò MR. Investigating causality in associations between education and smoking: a two-sample Mendelian randomization study. *Int J Epidemiol* 2018; **47**: 1131-1140 [PMID: 29961807 DOI: 10.1093/ije/dyy131]
- 24 **Akter S**, Goto A, Mizoue T. Smoking and the risk of type 2 diabetes in Japan: A systematic review and meta-analysis. *J Epidemiol* 2017; **27**: 553-561 [PMID: 28716381 DOI: 10.1016/j.je.2016.12.017]
- 25 **Xu H**, Wang Q, Sun Q, Qin Y, Han A, Cao Y, Yang Q, Yang P, Lu J, Liu Q, Xiang Q. In type 2 diabetes induced by cigarette smoking, activation of p38 MAPK is involved in pancreatic β-cell apoptosis. *Environ Sci Pollut Res Int* 2018; **25**: 9817-9827 [PMID: 29372523 DOI: 10.1007/s11356-018-1337-3]
- 26 **Tillmann T**, Vaucher J, Okbay A, Pikhart H, Peasey A, Kubinova R, Pajak A, Tamosiunas A, Malyutina S, Hartwig FP, Fischer K, Veronesi G, Palmer T, Bowden J, Davey Smith G, Bobak M, Holmes MV. Education and coronary heart disease: mendelian randomisation study. *BMJ* 2017; **358**: j3542 [PMID: 28855160 DOI: 10.1136/bmj.j3542]
- 27 **Böckerman P**, Viinikainen J, Pulkki-Råback L, Hakulinen C, Pitkänen N, Lehtimäki T, Pehkonen J, Raitakari OT. Does higher education protect against obesity? *Prev Med* 2017; **101**: 195-198 [PMID: 28645627 DOI: 10.1016/j.ypmed.2017.06.015]
- 28 **Velázquez-López L**, Muñoz-Torres AV, Medina-Bravo P, Vilchis-Gil J, Klönder-Klönder M, Escobedo-de la Peña J. Multimedia education program and nutrition therapy improves HbA1c, weight, and lipid profile of patients with type 2 diabetes: a randomized clinical trial. *Endocrine* 2017; **58**: 236-245 [PMID: 28921414 DOI: 10.1007/s12020-017-1416-0]
- 29 **Wadolowska L**, Hamulka J, Kowalkowska J, Ulewicz N, Hoffmann M, Gornicka M, Bronkowska M, Leszczynska T, Glibowski P, Korzeniowska-Ginter R. Changes in Sedentary and Active Lifestyle, Diet Quality and Body Composition Nine Months after an Education Program in Polish Students Aged 11-12 Years: Report from the ABC of Healthy Eating Study. *Nutrients* 2019; **11** [PMID: 30717465 DOI: 10.3390/nu11020331]
- 30 **Prospective Studies Collaboration**, Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, Qizilbash N, Collins R, Peto R. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* 2009; **373**: 1083-1096 [PMID: 19299006 DOI: 10.1016/S0140-6736(09)60318-4]
- 31 **Lyall DM**, Celis-Morales C, Ward J, Illiodromiti S, Anderson JJ, Gill JMR, Smith DJ, Ntuk UE, Mackay DF, Holmes MV, Sattar N, Pell JP. Association of Body Mass Index With Cardiometabolic Disease in the UK Biobank: A Mendelian Randomization Study. *JAMA Cardiol* 2017; **2**: 882-889 [PMID: 28678979 DOI: 10.1001/jamacardio.2016.5804]
- 32 **Zhu Z**, Zheng Z, Zhang F, Wu Y, Trzaskowski M, Maier R, Robinson MR, McGrath JJ, Visscher PM, Wray NR, Yang J. Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat Commun* 2018; **9**: 224 [PMID: 29335400 DOI:]

- 10.1038/s41467-017-02317-2]
- 33 **Cheng L**, Zhuang H, Ju H, Yang S, Han J, Tan R, Hu Y. Exposing the Causal Effect of Body Mass Index on the Risk of Type 2 Diabetes Mellitus: A Mendelian Randomization Study. *Front Genet* 2019; **10**: 94 [PMID: 30891058 DOI: 10.3389/fgene.2019.00094]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoftice@wjnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjnet.com>

