

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 58896

Manuscript Type: CASE REPORT

Early colon cancer with total gastrointestinal enteropathy-associated T-cell lymphoma: A case report

Zhang MY *et al.* Early colon cancer with T-cell lymphoma

Meng-Yuan Zhang, Cong-Cong Min, Wei-Wei Fu, Hua Liu, Xiao-Yan Yin, Cui-Ping Zhang, Zi-Bin Tian, Xiao-Yu Li

Match Overview

1	Internet 52 words crawled on 07-May-2020 www.wjgnet.com	2%
2	Crossref 22 words Sanghui Park, Young Hyeh Ko. "Peripheral T cell lymphoma i n Asia", International Journal of Hematology, 2014	1%
3	Internet 15 words crawled on 29-Jan-2020 journals.sagepub.com	1%
4	Crossref 14 words Sarah Ondrejka, Deepa Jagadeesh. "Enteropathy-Associat ... T-Cell Lymphoma", Current Hematologic Malignancy Reports	1%
5	Crossref 14 words Yifan Yang, Sukhjeet S. Batth, Mingyi Chen, Dariusz Borys, H o Phan. "Enteropathy-Associated T Cell Lymphoma Presen ...	1%
6	Internet 14 words crawled on 27-Apr-2020 f6publishing.blob.core.windows.net	1%



ALL

IMAGES

VIDEOS

4,480,000 Results

Any time ▾

[A case of colon perforation due to enteropathy-associated ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3607762>

Mar 21, 2013 · Enteropathy-associated T-cell lymphoma (EATL) is an extremely rare disease, which is often related to **gluten-sensitive enteropathy**. It is an uncommon **intestinal lymphoma** with very poor prognosis and high mortality rate. In the absence of specific symptoms or radiological findings, it is difficult to diagnose **early**.

Cited by: 9

Author: Jun Bong Kim, Seong Hwan Kim, Young ...

Publish Year: 2013

[Type I enteropathy-associated T-cell lymphoma in the colon ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4768892>

Feb 22, 2016 · Enteropathy-associated T-cell lymphoma (EATL) is a rare **gastrointestinal non-Hodgkin's lymphoma**, originating from **intraepithelial T-lymphocyte**, which is specifically associated with **celiac disease**. EATL most commonly presents in the sixth and seventh decades of life.

Cited by: 1

Author: Jiu-Cong Zhang, Yong Wang, Xiu-Feng W...

Publish Year: 2016

[Enteropathy-associated T-cell lymphoma presenting with ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4063612>

The current study presents a rare case of primary colon T-cell lymphoma in a 16-year-old male with poor prognosis, as well as a case of **gastrointestinal lymphoma** occurring in the duodenum and colon in a 62-year-old male with a 10-year history of NHL.

Cited by: 2

Author: Guohui Jiao, Zhongqing Zheng, Kui Jiang,...

Publish Year: 2014

[Enteropathy associated T-cell lymphoma - Cancer Therapy ...](#)

<https://www.cancertherapyadvisor.com/home/decision...> ▾

Enteropathy associated T-cell lymphoma What every physician needs to know: Enteropathy associated T-cell lymphoma (EATL) is an aggressive peripheral T-cell lymphoma with a ...

PEOPLE ALSO ASK

[What is intestinal T cell lymphoma?](#)

Early colon cancer with total gastrointestinal enteropathy-associated



ALL

IMAGES

VIDEOS

4,810,000 Results

Any time ▼

Enteropathy-associated T-cell lymphoma presenting with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4063612>

In contrast to **B-cell lymphoma**, **enteropathy-associated T-cell lymphoma (ETCL)** is a highly **aggressive T-cell lymphoma** with a poor prognosis and due to non-specific clinical and endoscopic observations, **early** diagnosis and appropriate treatment may be delayed .

Cited by: 2**Author:** Guohui Jiao, Zhongqing Zheng, Kui Jian...**Publish Year:** 2014

A case of colon perforation due to enteropathy-associated ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3607762>

Mar 21, 2013 · **Enteropathy-associated T-cell lymphoma (EATL)** is an extremely rare disease, which is often related to **gluten-sensitive enteropathy**. It is an uncommon **intestinal lymphoma** with very poor prognosis and high mortality rate. In the absence of specific symptoms or radiological findings, it is difficult to diagnose **early**.

Cited by: 9**Author:** Jun Bong Kim, Seong Hwan Kim, Young...**Publish Year:** 2013

Type I enteropathy-associated T-cell lymphoma in the colon ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4768892>

Enteropathy-associated T-cell lymphoma (EATL) is a rare **gastrointestinal non-Hodgkin's lymphoma**, originating from **intraepithelial T-lymphocyte**, which is specifically **associated** with **celiac disease**. EATL most commonly presents in the sixth and seventh decades of life.

Cited by: 1**Author:** Jiu-Cong Zhang, Yong Wang, Xiu-Feng ...**Publish Year:** 2016

PEOPLE ALSO ASK

What is intestinal T cell lymphoma?





Early colon cancer with total gastrointestinal enteropathy-associated



ALL IMAGES VIDEOS MAPS NEWS SHOPPING

4,670,000 Results Any time ▾

[Enteropathy-associated T-cell lymphoma presenting with ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4063612>

The **intestinal T-cell lymphoma** is endoscopically characterized by focally, multifocally or diffusely distributed polymorphic irregular ulcers, which most frequently involve the **ileocecum and colon**. Its location in the **intestine** is **associated with enteropathy** and develops from the **intraepithelial T-lymphocytes** of the intestine (5).

Cited by: 2 **Author:** Guohui Jiao, Zhongqing Zheng, Kui Jiang...

Publish Year: 2014

[Endoscopic features of gastro-intestinal lymphomas: From ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4177478>

Sep 28, 2014 - **Gastric MALT lymphoma**. MALT lymphomas arise in the context of the lymphoid infiltrates of the mucosa layer of hollow organs and from glandular tissues[22,28].The GI tract is the most frequently involved zone (almost 50% of cases)[], particularly the **stomach** (85% of cases).The gross intestine represents only a 0.4% of cases.

Cited by: 16 **Author:** Calogero Vetro, Alessandra Romano, Ire...

Publish Year: 2014

[A case of colon perforation due to enteropathy-associated ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3607762>

Mar 21, 2013 - **Enteropathy-associated T-cell lymphoma (EATL)** is an extremely rare disease, which is often related to **gluten-sensitive enteropathy**. It is an uncommon **intestinal lymphoma** with very poor prognosis and high mortality rate. In the absence of specific symptoms or radiological findings, it is difficult to diagnose **early**.

Cited by: 9 **Author:** Jun Bong Kim, Seong Hwan Kim, Young ...

Publish Year: 2013

[Primary Gastrointestinal Lymphoma](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5059001>

Oct 30, 2015 - INTRODUCTION. Primary **gastrointestinal lymphoma** (PGIL) is a tumor of **gastrointestinal** (GI) tract, as its primary lesion might invade the lymph nodes of the related drainage area, excluding tumors involving the liver, spleen, or lymphomas in patients who exhibit GI symptoms,

Enteropathy-associated T-cell lymphoma

Disease

Enteropathy-associated T-cell lymphoma, previously termed enteropathy-associated T-cell lymphoma, type I and at one time termed enteropathy-type T-cell lymphoma, is a complication of coeliac disease in which a malignant T-cell lymphoma develops in areas of the small intestine afflicted by the disease's intense inflammation. While a relatively rare disease, it is the most common type of primary gastrointestinal T-cell lymphoma.



People also search for

[Celiac Disease](#)

[Non-Hodgkin's Lymphoma](#)

[Post-transplant lymphoproliferative disorder](#)

[Mycosis fungoides](#)

[Waldenstrom Macroglobulinemia](#)

[See more ▾](#)

Data from: Wikipedia

[Suggest an edit](#)