

4
Name of Journal: *World Journal of Gastrointestinal Surgery*
Manuscript NO: 58901
Manuscript Type: ORIGINAL ARTICLE

Prospective Study
Perianal fistulodesis - A pilot study of a novel minimally invasive surgical and medical approach for closure of perianal fistulae

Roxanne Villiger, Daniela Cabalzar-Wondberg, Daniela Zeller, Pascal Frei, Luc Biedermann, Christian Schneider, Michael Scharl, Gerhard Rogler, Matthias Turina, Andreas Rickenbacher, Benjamin Misselwitz

Abstract

BACKGROUND

Perianal fistulae strongly impact on quality of life of affected patients.

Match Overview

1	Crossref 39 words Luana BERNARDI, Carlos Henrique Marques dos SANTOS, V erônica Assalin Zorgetto PINHEIRO, Rodrigo Juliano OLIVEIR	1%
2	Crossref 36 words Yuming Chong, Chang Han, Ji Li, Xiao Long. "Mapping global r esearch trends in stem cell therapy for inflammatory bowel c...	1%
3	Internet 13 words crawled on 27-Aug-2020 mafiadoc.com	<1%
4	Internet 13 words crawled on 02-Dec-2020 f6publishing.blob.core.windows.net	<1%
5	Internet 12 words crawled on 06-Dec-2020 kclpure.kcl.ac.uk	<1%
6	Internet 12 words crawled on 22-Jun-2020 www.hal.inserm.fr	<1%



Perianal fistulodesis - A pilot study of a novel minimally invasive surgical



ALL

IMAGES

VIDEOS

9,580 Results

Any time ▼

Novel Approaches to Ileocolic and Perianal Fistulising ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6280273>

With regard to **surgical** treatments, a **rectal** sleeve advancement flap is sometimes an option for the primary **closure of perianal** fistulas in the case of a single internal opening and in the absence of **rectal** inflammation. Additionally, therapy with antibiotics or azathioprine should be considered a second-line therapy in case of failure of the ...

Author: Gaetano Luglio, Gianluca Cassese, Al... **Publish Year:** 2018

E Management of patients with complex perianal fistulas in ...

<https://www.sciencedirect.com/science/article/pii/S1590865819309259>

May 01, 2020 · 6) Patients whose **perianal** fistulas were previously treated and have shown an inadequate response (absence of **closure** of part or all fistula tracks, or new fistula during induction treatment) or a loss of response (fistula relapse during maintenance treatment after initial fistula **closure**) while they were receiving either an immunosuppressive ...

Search Tools

[Turn off Hover Translation \(关闭取](#)

Feedback

Perianal fistulodesis - A pilot study of a novel minimally invasive sur



ALL

IMAGES

VIDEOS

16,400 Results

Any time ▼

[Novel Approaches to Ileocolic and Perianal Fistulising ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6280273>

With regard to **surgical** treatments, a **rectal** sleeve advancement flap is sometimes an option for the primary **closure of perianal** fistulas in the case of a single internal opening and in the absence of **rectal** inflammation. Additionally, therapy with antibiotics or azathioprine should be considered a second-line therapy in case of failure of the ...

Author: Gaetano Luglio, Gianluca Cassese, Al... **Publish Year:** 2018

[Minimally Invasive Anal Fistula Treatment-a True Minimal ...](#)

<https://europepmc.org/article/PMC/PMC4744224> ▼

Dec 15, 2015 · **Minimally Invasive** Anal Fistula Treatment-a True Minimal **Invasive Surgical** Procedure. ... The advent of minimal access surgery revolutionized the concept of **surgical** practice. The **approach** placed the patient at the center of all **surgical** decisions. It was for the first time that the comfort and needs of the patient were given paramount ...

Author: P. K. Chowbey **Publish Year:** 2015

[Symptom Amelioration in Crohn's Perianal Fistulas Using ...](#)

<https://academic.oup.com/ecco-jcc/article/12/9/1067/5003424> ▼

May 25, 2018 · VAAFT and **Fistula** tract Laser **Closure** [FiLaC™] are novel therapies for definitive sphincter-sparing surgical treatment of **anal fistulas**, which have largely been evaluated in cryptoglandular **fistula**. 18 The rationale behind these **minimally invasive** procedures, which probe the **fistula** tract via the external opening, is to ablate the tract either by electrocautery [VAAFT] or by laser ...

Cited by: 9 **Author:** Samuel O Adegbola, Samuel O Adegbola...

Publish Year: 2018

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

16,500 Results

Any time ▾

Novel Approaches to Ileocolic and Perianal Fistulising ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6280273>

With regard to **surgical** treatments, a **rectal** sleeve advancement flap is sometimes an option for the primary **closure of perianal** fistulas in the case of a single internal opening and in the absence of **rectal** inflammation. Additionally, therapy with antibiotics or azathioprine should be considered a second-line therapy in case of failure of the ...

Author: Gaetano Luglio, Gianluca Cassese, Alfo... **Publish Year:** 2018

Most recent papers with the keyword Perianal Fistulae ...

<https://read.qxmd.com/keyword/174238> ▾

AIM: Ligation of the intersphincteric fistula tract (LIFT) procedure is increasingly used for **surgical closure** of high **perianal** fistulas in Crohn's disease. Currently, data on postoperative MRI findings is scarce, while considered most important to assess healing and recurrences. Therefore, we evaluated fistula characteristics on MRI and the relation with clinical outcomes after LIFT.

Laparoscopy, Endoscopy, and Minimally Invasive Colorectal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5380456>

Transanal TME is a **novel minimally invasive approach** that has emerged in response to the challenges associated with traditional open and laparoscopic **surgical** approaches to **rectal** cancer. Although taTME is technically challenging, at least initially, its technical advantages as well as the potential oncologic benefit are exciting.

Cited by: 7 **Author:** Pasithorn A. Suwanabol, Justin A. Maykel

Publish Year: 2017

Endoscopic vacuum therapy with instillation (iEVT) – a ...

<https://europepmc.org/article/PMC/PMC7687668> ▾

Feb 24, 2020 · The mean number of iEVT sessions was 5.8 ± 2.3 . The mean time of iEVT management was 20.7 ± 8.8 days. Locally, a reduction in purulent discharge and defect's dimension with contraction