

Match Overview

1	Internet 96 words crawled on 29-Sep-2020 www.wjgnet.com	4%
2	Internet 30 words crawled on 24-Sep-2020 core.ac.uk	1%
3	Crossref 22 words "Clinical Investigation of Portal Hypertension", Springer Nature, 2019	1%
4	Crossref 19 words Umesha Boregowda, Chandraprakash Umapathy, Nasir Halim, Madhav Desai et al. "Update on the management of [...]"	1%
5	Crossref 13 words Shuntaro Yoshida, Hirotsugu Watabe, Masaaki Akahane, Yusuke Kyoden et al. "Usefulness of multi-detector helical CT"	1%
6	Internet 12 words crawled on 01-Dec-2020	1%

7
Name of Journal: *World Journal of Gastrointestinal Surgery*

Manuscript NO: 59214

Manuscript Type: CASE REPORT

Multidetector computed tomography three-dimensional and multiplanar reconstruction diagnosis of a rare cause of gastrointestinal bleeding: A case report

Yong Cai, Xiao-Bo Chen, Da-Hai Mai, Ping Wu, Yong-Hai Chen, Hua Chen



ALL

IMAGES

VIDEOS

6,850 Results

Any time ▾

Multidetector CT of Vascular Compression Syndromes in the ...

<https://pubs.rsna.org/doi/full/10.1148/rg.341125010>

May-Thurner syndrome: **case report** and **review** of the literature involving modern endovascular therapy. Vascular 2009;17(6): 330–335. Crossref, Medline, Google Scholar; 40. Oguzkurt L, Tercan F, Pourbagher MA, Kizilkilic O, Turkoz R, Boyvat F. **Computed tomography** findings in 10 cases of iliac **vein** compression (May-Thurner) syndrome.

Cited by: 72

Author: Ramit Lamba, Dawn T. Tanner, Simran Sekh...

Publish Year: 2014

(PDF) Role of MDCT angiography in assessment of vascular ...

<https://www.researchgate.net/publication/259129485...>

superior mesenteric artery had a common origin in three in- stances (2%), and the celiac axis was absent in two (1%), with the common hepatic, splenic and left gastric arising from the

(PDF) Multidetector CT Anatomy of Drainage Routes of ...

<https://www.researchgate.net/publication/234142902...>

Interportal communications with the left gastric **vein** and varices occurred from the left portal **vein** in 13, from the gastroepiploic **vein** in one, and from a **superior mesenteric vein** branch in one.

Search Tools

Turn on Hover Translation (开启取词)

激活 Windows

转到“设置”以激活 Windows。

Multidetector computed tomography three-dimensional and multip



ALL

IMAGES

VIDEOS

59,600 Results

Any time ▼

Chest computed tomography with multiplanar reformatted ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2556736>

Methods. **Three-dimensional reconstruction** of the airways using a workstation connected to a **multidetector chest computed tomography** (CT) scanner may change the diagnostic strategy in patients with blunt chest trauma with clinical signs evocative of bronchial rupture.

Cited by: 19

Author: Morgan Le Guen, Catherine Beigelman, ...

Publish Year: 2007

Diagnosis of incarcerated intramesosigmoid hernia aided by ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6195874>

We **report** an extremely **rare case** of incarcerated intramesosigmoid hernia that was diagnosed by **multidetector computed tomography** with **multiplanar reconstruction** images. The finding of narrow belt-shaped fluid retention contacting the tip of the incarcerated short intestinal loop is characteristic of intramesosigmoid hernia and will be useful ...

Cited by: 1

Author: Hideki Nagano, Takanori Goi, Seiichi Tag...

Publish Year: 2018

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

59,300 Results

Any time ▾

[Diagnosis of incarcerated intramesosigmoid hernia aided by ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6195874>

We **report** an extremely **rare case** of incarcerated intramesosigmoid hernia that was diagnosed by **multidetector computed tomography** with **multiplanar reconstruction** images. The finding of narrow belt-shaped fluid retention contacting the tip of the incarcerated short intestinal loop is characteristic of intramesosigmoid hernia and will be useful ...

Cited by: 1

Author: Hideki Nagano, Takanori Goi, Seiichi Taguc...

Publish Year: 2018

[Chest computed tomography with multiplanar reformatted ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2556736>

Methods. **Three-dimensional reconstruction** of the airways using a workstation connected to a **multidetector** chest **computed tomography** (CT) scanner may change the diagnostic strategy in patients with blunt chest trauma with clinical signs evocative of bronchial rupture.

Cited by: 19

Author: Morgan Le Guen, Catherine Beigelman, Bel...

Publish Year: 2007

[Role of three-dimensional multidetector computed ...](#)

<https://www.sciencedirect.com/science/article/pii/S0167527307004524>

Jun 23, 2008 · Superior mesenteric artery (SMA) aneurysms are the third most common visceral artery aneurysms. **Diagnosis** of SMA aneurysm can be achieved by abdominal ultrasonogram, postcontrast **computed tomography** (CT), **three-dimensional** (3-D) **multidetector** (MD) CT, or selective angiography of the SMA. 3-D MD CT may play an important role for the evaluation of characteristic morphology of ...

Cited by: 5

Author: Cheol Ung Choi, Seung-Woon Rha, Soon Yo...

Publish Year: 2008

[Diagnosis of incarcerated intramesosigmoid hernia aided by ...](#)

<https://link.springer.com/article/10.1186/s40792-018-0535-z> ▾