

Match Overview

1	Crossref 27 words Dina Bassiouny, Nadia Ismiil, Valerie Dubé, Guangming Han et al. "Comprehensive Clinicopathologic and Updated Immu..."	1%
2	Internet 20 words crawled on 30-Sep-2020 insights.ovid.com	1%
3	Crossref 17 words Katherine C. Kurnit, Abdulrahman K. Sinno, Bryan M. Fellman, Aaron Varghese et al. "Effects of Gastrointestinal-Type Cl..."	1%
4	Crossref Posted Content 17 words Yoshitaka Kiya, Yuichi Nagakawa, Chie Takishita, Hiroaki Osakabe et al. "Neuroendocrine carcinoma of the common bile (..."	1%
5	Internet 15 words crawled on 07-Oct-2020 pesquisa.bvsalud.org	1%
6	Crossref 13 words Chi-Hau Chen, Ming-Kwang Shyu, Shu-Wei Wang, Chih-Hsing Chou et al. "MUC20 promotes aggressive phenotypes of €..."	1%
7	Internet 12 words crawled on 17-Jan-2020 journals.sagepub.com	1%

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 59354

Manuscript Type: CASE REPORT

Gastrointestinal-type chemotherapy prolongs survival in an atypical primary ovarian mucinous carcinoma: A case report

Gastrointestinal-type chemotherapy in primary ovarian mucinous cancer patient

Abstract

BACKGROUND

Primary ovarian mucinous carcinoma is a rare histologic subtype of epithelial ovarian carcinoma and exhibits considerable morphologic overlap with secondary tumour. It is hard to differentiate primary from metastatic ovarian mucinous carcinoma by morphological and immunohistochemical features. Because of the histologic similarity between primary ovarian mucinous carcinoma and metastatic gastrointestinal carcinoma, it has been hypothesized that ovarian mucinous carcinomas might respond



ALL

IMAGES

VIDEOS

33,100 Results

Any time ▾

Amanda N. Fader's research works | Johns Hopkins ...

<https://www.researchgate.net/scientific...>

Objective: To estimate whether **gastrointestinal-type chemotherapy** was associated with improved **survival** compared with standard gynecologic regimens for women with **ovarian mucinous carcinoma**.

Borderline Ovarian Tumors - Atypical Proliferative Ovarian ...

https://www.glowm.com/section_view/heading... ▾

The pattern of gene expression of **mucinous carcinoma** can easily be distinguishable from serous, endometrioid and clear cell **carcinoma**. **Ovarian mucinous** tumors express the genes UC2, MUC3 and MUC17 that are typical of tumors of **mucinous** lineage, **primary** or metastatic. 190 Among other genes expressed by **mucinous** tumors is the CDx2, which is a very ...

Advanced Stage Mucinous Adenocarcinoma of the Ovary is ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC20862744>

Feb 01, 2011 · In conclusion, we found that advanced stage **primary mucinous carcinoma** of the ovary is very rare (0.5–1.5% of advanced **ovarian** epithelial invasive neoplasms), and that 61% (27/44) of the patients enrolled on this protocol as **primary mucinous** carcinomas were probably metastatic **carcinoma**

...

Cited by: 140

Author: Richard J. Zaino, Mark F. Brady, Subodh M. ...

Publish Year: 2011

PEOPLE ALSO ASK

What are the symptoms of mucinous cancer in the breast? ▾

What are the symptoms of ductal carcinoma? ▾

Gastrointestinal-type chemotherapy prolongs survival in an atypical



ALL

IMAGES

VIDEOS

31,300 Results

Any time ▾

Effects of Gastrointestinal-Type Chemotherapy in Women ...

<https://journals.lww.com/greenjournal/Fulltext/...>

Gastrointestinal-type chemotherapy with or without bevacizumab was associated with **improved survival** and should be considered in patients with **ovarian mucinous carcinoma** requiring adjuvant therapy.

Gastrointestinal-type chemotherapy is associated with **improved survival** in **ovarian mucinous carcinoma** when compared with standard **gynecologic regimens**.

Types of Ovarian Tumors - Ovarian Cancer | Johns Hopkins ...

<https://pathology.jhu.edu/ovarian-cancer/types-of-tumors> ▾

Mucinous carcinoma. A malignant epithelial tumor composed of gastrointestinal-type cells. **Mucinous carcinoma** accounts for 3-4% of all **primary ovarian** carcinomas. Most tumors are confined to the ovary at presentation. Advanced stage **primary mucinous ovarian carcinoma** is rare.

PEOPLE ALSO ASK

What are the symptoms of mucinous cancer in the breast? ▾

What is the incidence of serous ovarian tumors? ▾

What are the types of ovarian tumors? ▾

What are the risk factors for mucinous carcinoma? ▾

Feedback

A descriptive report of outcomes of primary mucinous ...

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

WORK

31,600 Results

Any time ▾

Open links in new tab



[Effects of Gastrointestinal-Type Chemotherapy in Women ...](#)

<https://journals.lww.com/greenjournal/Fulltext/...>

Gastrointestinal-type chemotherapy with or without **bevacizumab** was associated with improved **survival** and should be considered in patients with ovarian mucinous carcinoma requiring adjuvant therapy.

Gastrointestinal-type chemotherapy is associated with improved survival in ovarian **mucinous carcinoma** when compared with standard gynecologic regimens.

[Types of Ovarian Tumors - Ovarian Cancer | Johns Hopkins ...](#)

<https://pathology.jhu.edu/ovarian-cancer/types-of-tumors> ▾

Mucinous carcinoma. A malignant epithelial tumor composed of gastrointestinal-type cells. **Mucinous carcinoma** accounts for 3-4% of all **primary ovarian** carcinomas. Most tumors are confined to the ovary at presentation. Advanced stage **primary mucinous ovarian carcinoma** is rare.

[A descriptive report of outcomes of primary mucinous ...](#)

<https://ijgc.bmj.com/content/29/5/904> ▾

Jun 01, 2019 · Objective We described progression-free **survival** and overall **survival** in patients with **primary mucinous ovarian** cancer receiving adjuvant gynecologic versus gastrointestinal **chemotherapy** regimens. Methods We identified all **primary mucinous ovarian** cancer patients receiving adjuvant gynecologic or gastrointestinal **chemotherapy** regimens at a single institution from 1994 to 2016.

Cited by: 1 **Author:** Brooke A Schlappe, Qin C Zhou, Roisin O'C...

Publish Year: 2019

[Mucinous adenocarcinoma of the intestinal type arising ...](#)

<https://ovarianresearch.biomedcentral.com/articles/10.1186/1757-2215-5-41> ▾

Dec 05, 2012 · Mature cystic teratomas (MCTs) are the most common germ cell tumors of the ovary. Malignant transformation occurs in 1-2% of these neoplasms. Although most of the malignancies arising from MCTs are squamous cell carcinomas, adenocarcinoma of the gastrointestinal type is extremely rare. We herein present a **case** of adenocarcinoma of the intestinal type arising from a MCT.

Cited by: 11 **Author:** Masaaki Takai, Masanori Kanemura, Hiros...

Publish Year: 2012

[Primary ovarian mucinous carcinoid tumor: A case report ...](#)

<https://www.sciencedirect.com/science/article/pii/S1028455919301299>