

Oral and perioral herpes simplex virus type 1 (HSV-1 ...

<https://www.researchgate.net/publication/7080271...>

Herpes simplex virus type 1 (HSV-1) gives rise to a variety of clinical disorders and is a major cause of morbidity and mortality worldwide. HSV-1 infections are common in **oral and perioral** area.

Nongenital Herpes Simplex Virus - American Family Physician

<https://www.aafp.org/afp/2010/1101/p1075.html> ▾

**Diagnosis**

Epidemiology

Pathophysiology

Cause

Prognosis



Nongenital herpes simplex virus type 1 is a common infection usually transmitted during childhood via nonsexual contact. Most of these infections involve the oral mucosa or lips (herpes labialis). The diagnosis of an infection with herpes simplex virus type 1 is usually made by the appearance of the lesions (grouped vesicles or ulcers on an erythematous base) and patient history. However, if uncertain, the diagnosis of herpes labialis can be made by viral culture, polymerase chain reaction, serology, direc...

[See more on aafp.org](#)

Cited by: 58

Author: Richard P. Usatine, Rochelle Tinitigan

Publish Year: 2010

Neonatal Herpes Simplex Virus Type 1 Infection and Jewish ...

<https://academic.oup.com/jpids/article/4/2/126/1019841> ▾

Jul 23, 2014 · Jewish ritual circumcision rarely but occasionally includes a procedure involving direct **oral** suction of the wound, which can expose an **infant** to **infection** with **herpes simplex virus type 1** (HSV-1). This practice has provoked international controversy in recent years, but no systematic review of the clinical literature has previously been published.

Cited by: 12

Author: Brian F. Leas, Craig A. Umscheid

Publish Year: 2015

国内版 国际版

Oral and Perioral Herpes Simplex Virus Type I Infection in A five-moi



ALL IMAGES VIDEOS

21,800 Results

Any time ▼

Nongenital Herpes Simplex Virus - American Family Physician

<https://www.aafp.org/afp/2010/1101/p1075.html> ▼

< **Diagnosis** Epidemiology Pathophysiology Cause Prognosis >

Nongenital herpes simplex virus type 1 is a common infection usually transmitted during childhood via nonsexual contact. Most of these infections involve the oral mucosa or lips (herpes labialis). The diagnosis of an infection with herpes simplex virus type 1 is usually made by the appearance of the lesions (grouped vesicles or ulcers on an erythematous base) and patient history. However, if uncertain, the diagnosis of herpes labialis can be made by viral culture, polymerase chain reaction, serology, direc...

[See more on aafp.org](#)

Cited by: 57

Author: Richard P. Usatine, Rochelle Tinitigan

Publish Year: 2010

Oral and perioral herpes simplex virus type 1 (HSV-1 ...

<https://www.researchgate.net/publication/7080271...>

Herpes simplex virus type 1 (HSV-1) gives rise to a variety of clinical disorders and is a major cause of morbidity and mortality worldwide. HSV-1 infections are common in **oral and perioral** area.

Neonatal Herpes Simplex Virus Type 1 Infection and Jewish ...

<https://academic.oup.com/jpids/article/4/2/126/1019841> ▼

Jul 23, 2014 · Jewish ritual circumcision rarely but occasionally includes a procedure involving direct **oral** suction of the wound, which can expose an **infant** to **infection** with **herpes simplex virus type 1** (HSV-1). This practice has provoked international controversy in recent years, but no systematic review of the clinical literature has previously been published.

Match Overview

There are no matching sources for this report.

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 59468

Manuscript Type: CASE REPORT

Oral and Perioral Herpes Simplex Virus Infection Type I in A Five-Month-Old Infant: A case report and discussion.

Aloyouny *et al.* HSV in A Five-Month-old Infant

Abstract**BACKGROUND**

Herpes simplex virus (HSV) is a highly infectious pathogen that is easily transmitted via the bodily fluids of an infected individual. This virus usually affects individuals older than six months of age, and rarely causes lesions or symptoms in younger patients.

CASE SUMMARY

国内版

国际版

Oral and perioral herpes simplex virus infection type I in a five-mo



ALL

IMAGES

VIDEOS

35,000 Results

Any time ▾

Primary Herpes Simplex Virus Infection of the Nipple in a ...

<https://www.mdedge.com/dermatology/article/109374/infectious-diseases/primary-herpes...>

Herpes mastitis should be included in the differential diagnosis for breast pain during lactation. Children of breastfeeding women diagnosed with **herpes** mastitis require immediate evaluation for a possible source of the **infection**, as complications of **herpes** viral **infection** in infants can be severe and life threatening.

Primary Maternal Herpes Simplex Virus-1 Gingivostomatitis ...

<https://academic.oup.com/jpids/article-abstract/1/4/299/951050> ▾

Jul 10, 2012 · Neonatal **herpes simplex virus** (HSV) has long been recognized as a devastating consequence of maternal HSV **type** 1 (HSV-1) and HSV **type** 2 (HSV-2) genital **infections** [1, 2]. Although recurrent genital HSV **infection** is the most common manifestation of HSV during pregnancy, women who have a primary genital HSV **infection** at term are at the greatest risk of transmitting the **virus** ...

Cited by: 1

Author: Sara A. Healy, Kathleen M. Mohan, Ann ...

Publish Year: 2012

Case Report: Concurrent meningococcal and herpes simplex ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992534>

Apr 15, 2014 · The facial rash had settled but the swabs of the area returned positive for **herpes simplex virus** (HSV) **type** 1. At the time there were no clinical features to indicate that the **infection** had spread to the CSF, but the secondary fever that had evolved since stopping the topical aciclovir and the subsequent rise in C reactive protein (CRP) and WCC ...

Cited by: 2

Author: Jasmin Ali, Hannah Walsh, Swathi Sana...

Publish Year: 2014

PEOPLE ALSO ASK