

## Match Overview

1	<b>Internet</b> 17 words crawled on 10-Nov-2020 <a href="http://www.hindawi.com">www.hindawi.com</a>	1%
2	<b>Internet</b> 12 words crawled on 03-Jan-2021 <a href="http://pesquisa.bvsalud.org">pesquisa.bvsalud.org</a>	<1%
3	<b>Crossref</b> 12 words Park, Jun Seok, Gyu-Seog Choi, Seon Hahn Kim, Hyeong Ro k Kim, Nam Kyu Kim, Kang Young Lee, Sung Bum Kang, Ji Y	<1%
4	<b>Crossref</b> 12 words "Basic Research Poster Session", Journal of Endourology, 11/ 2008	<1%

**Name of Journal:** *World Journal of Gastrointestinal Surgery*

**Manuscript NO:** 59973

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*

**Retrospective research of neoadjuvant therapy on tumor-downstaging, post-operative complications and prognosis in locally advanced rectal cancer**

Research of Neoadjuvant therapy in LARC

**Abstract**

BACKGROUND

Neoadjuvant therapy is becoming increasingly important in locally advanced rectal cancer. Hence, such research has become a concerning problem.

AIM

To evaluate the downstaging effect of neoadjuvant therapy, its impact on postoperative

Retrospective Research of Neoadjuvant Therapy on tumor-downsta



ALL

IMAGES

VIDEOS

81,200 Results

Any time ▾

## A single-institution retrospective analysis of outcomes ...

<https://www.journalrcr.org/article.asp?issn=2588...>

Journal of Radiation and **Cancer Research**, India. Background: **Neoadjuvant** chemoradiotherapy (NACTRT) is an established standard of care for the management of **locally advanced rectal cancer** (LARC). A **retrospective** analysis was conducted to assess the results of treatment and predictors of overall survival (OS) for patients treated in a tertiary care hospital in India from 2013 to 2019.

**Author:** Kannan Venkatesan, Vivek Anand, Ranj... **Publish Year:** 2020

## Frontiers | Radiomics-Based Preoperative Prediction of ...

<https://www.frontiersin.org/articles/10.3389/fonc.2020.00604> ▾

May 11, 2020 · Introduction. **Neoadjuvant therapy** followed by total mesorectal excision (TME) is the standard treatment for patients with **locally advanced rectal cancer** (LARC) ().After **neoadjuvant therapy**, ~50–60% of patients are downstaged, and ~20% show pathologic complete response (1–3).Although TME is effective at providing local tumor control, it is also associated with significant genitourinary ...

**Cited by:** 3 **Author:** Xuezhi Zhou, Xuezhi Zhou, Yongju Yi, Yongj...

**Publish Year:** 2020

## Neoadjuvant therapy and subsequent treatment in rectal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6768820>

Standard **Therapy** and Indications. **Neoadjuvant** (chemo) radiotherapy followed by TME has revolutionized the oncological outcomes of patients with resectable **rectal cancer** in last decades, leading to a local recurrence rate as low as 5%-6% 2, 11).**Neoadjuvant** treatment also contributes to the **tumor downstaging** and downsizing, which facilitate surgical resection and sphincter preservation.

**Cited by:** 1 **Author:** Qiyuan Qin, Lei Wang

**Publish Year:** 2018



ALL

IMAGES

VIDEOS

83,600 Results

Any time ▾

## A single-institution retrospective analysis of outcomes ...

<https://www.journalrcr.org/article.asp?issn=2588...>

Journal of Radiation and **Cancer Research**, India. Background: **Neoadjuvant** chemoradiotherapy (NACTRT) is an established standard of care for the management of **locally advanced rectal cancer** (LARC). A **retrospective** analysis was conducted to assess the results of treatment and predictors of overall survival (OS) for patients treated in a tertiary care hospital in India from 2013 to 2019.

**Author:** Kannan Venkatesan, Vivek Anand, Ranj... **Publish Year:** 2020

## A Retrospective Analysis on Two-week Short-course Pre ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5122946>

The **neoadjuvant** radiotherapy regimen for **locally advanced rectal cancer** includes pre-operative short-course radiotherapy (represented by five fractions of 5Gy over 1 week) and pre-operative long-course chemo-radiotherapy (CRT) consisting of 5-fluorouracil (5-Fu) or capecitabine.

**Cited by:** 2 **Author:** Chen Shi, Hao Zhou, Xiaofan Li, Yong Cai

**Publish Year:** 2016

## Neoadjuvant-intensified treatment for rectal cancer: Time ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3662944>

May 28, 2013 · Core tip: Management of **rectal cancer** requires a multimodality treatment approach. The objective of this study was to determine whether **neoadjuvant-intensified** radiochemotherapy, using traditional radiation **therapy** in combination with oxaliplatin and 5-fluorouracil (5-FU), could improve the overall and disease-free survival rates in patients with **locally advanced rectal cancer**.

**Cited by:** 18 **Author:** Daniela Musio, Francesca De Felice, Nadia ...

**Publish Year:** 2013

[ALL](#)[IMAGES](#)[VIDEOS](#)[MAPS](#)[NEWS](#)[SHOPPING](#)

81,900 Results

Any time ▾

### [A Retrospective Analysis on Two-week Short-course Pre ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5122946>

The **neoadjuvant** radiotherapy regimen for **locally advanced rectal cancer** includes pre-operative short-course radiotherapy (represented by five fractions of 5Gy over 1 week) and pre-operative long-course...

**Cited by:** 2**Author:** Chen Shi, Hao Zhou, Xiaofan Li, Yong Cai**Publish Year:** 2016

### [Neoadjuvant therapy for locally advanced rectal cancer ...](#)

<https://www.sciencedirect.com/science/article/pii/S1533002821000013>

Jan 05, 2021 · Since **neoadjuvant therapy** became standard of care for **locally advanced rectal cancer**, the additional benefit of adjuvant chemotherapy has been reexamined. Several studies have attempted t...

### [A single-institution retrospective analysis of outcomes ...](#)

<https://www.journalrcr.org/article.asp?issn=2588...>

Journal of Radiation and **Cancer Research**, India. Background: **Neoadjuvant** chemoradiotherapy (NACTRT) is an established standard of care for the management of **locally advanced rectal cancer**...

**Author:** Kannan Venkatesan, Vivek Anand, Ranj... **Publish Year:** 2020

### [Neoadjuvant treatment \(FOLFOX4 plus hypofractionated ...](#)

<https://journals.sagepub.com/doi/full/10.1177/1758835920977139>

Dec 08, 2020 · **Neoadjuvant** concurrent fluoropyrimidine-based chemoradiotherapy (CRT) followed by total mesorectal excision (TME) is considered as the standard treatment for patients with **locally**...

**Author:** Alessandro Passardi, Ilario Giovanni Ra... **Publish Year:** 2020

### [Total Neoadjuvant Therapy: A Shifting Paradigm in Locally ...](#)

<https://www.sciencedirect.com/science/article/pii/S1533002817300543>

Mar 01, 2018 · Introduction. Colorectal carcinoma (CRC) is the second leading cause of **cancer**-related