

51,500 Results

Any time ▾

[Long-term control with chemoradiation of initially ...](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6431024>

Extended **long-term** control of a **poorly differentiated metastatic** (stage IV) mixed adenoneuroendocrine **carcinoma** is rare. The **multimodal** first- and second-line regimens of radiotherapy and chemotherapy...

Cited by: 1

Author: S. Semrau, A. Agaimy, M. Pavel, D. Lubgan, ...

Publish Year: 2019

[Long-term control with chemoradiation of initially ...](#)<https://www.ncbi.nlm.nih.gov/pubmed/30902067>

Mar 23, 2019 · The patient has been recurrence-free for more than 5 years. CONCLUSIONS: Extended **long-term** control of a **poorly differentiated metastatic** (stage IV) mixed adenoneuroendocrine **carcinoma**...

Cited by: 1

Author: S. Semrau, A. Agaimy, M. Pavel, D. Lubgan, ...

Publish Year: 2019

[\[PDF\] Long-term control with chemoradiation of initially ...](#)<https://jmedicalcasereports.biomedcentral.com/...>

metastasis and chemotherapy resulted in a clinical **complete response**. The patient has been recurrence-free for more than 5 years. Conclusions: Extended **long-term** control of a **poorly differentiated metastatic**...

[Long-term control with chemoradiation of initially ...](#)<https://jmedicalcasereports.biomedcentral.com/...> ▾

Mar 23, 2019 · The patient has been recurrence-free for more than 5 years. Extended **long-term** control of a **poorly differentiated metastatic** (stage IV) mixed adenoneuroendocrine **carcinoma** is rare. The...

Cited by: 1

Author: S. Semrau, A. Agaimy, M. Pavel, D. Lubgan, ...

Publish Year: 2019

[Poorly Differentiated Neuroendocrine Tumors | Oncohemakey](#)<https://oncohemakey.com/poorly-differentiated-neuroendocrine-tumors> ▾

Poorly differentiated neuroendocrine carcinomas of the gastrointestinal tract are rare and limited data are available to guide treatment. These tumors have been treated akin to small cell lung **cancer** given th...

国内版 国际版

Long term complete response in metastatic poorly-differentiated ne



ALL IMAGES VIDEOS

50,200 Results Any time ▾

[Long-term control with chemoradiation of initially ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/30902067>

Mar 23, 2019 · The patient has been recurrence-free for more than 5 years. CONCLUSIONS:

Extended **long-term** control of a poorly differentiated **metastatic** (stage IV) mixed...

Cited by: 1

Author: S. Semrau, A. Agaimy, M. Pavel, D. Lubg...

Publish Year: 2019

[Long term outcomes of neuroendocrine carcinomas \(high ...](#)

<https://www.sciencedirect.com/science/article/pii/S1878788613001689>

Feb 01, 2014 · Feb 01, 2014 · Introduction. **Neuroendocrine** carcinomas (NEC) of the colon and **rectum** are rare, extremely aggressive neoplasms with a high mortality rate .The **term**,...

Cited by: 23

Author: E. Aytac, Y. Ozdemir, G. Ozuner

Publish Year: 2014

[Poorly Differentiated Neuroendocrine Tumors | Oncohemakey](#)

<https://oncohemakey.com/poorly-differentiated-neuroendocrine-tumors> ▾

Poorly differentiated **neuroendocrine** carcinomas of the gastrointestinal tract are rare and limited data are available to guide treatment. These tumors have been treated akin to small cell lung...

[Long-term control with chemoradiation of initially ...](#)

<https://jmedicalcasereports.biomedcentral.com/...> ▾

Mar 23, 2019 · The patient has been recurrence-free for more than 5 years. Extended **long-term** control of a poorly differentiated **metastatic** (stage IV) mixed adenoneuroendocrine **carcinoma** i...

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 60168

Manuscript Type: CASE REPORT

Long term complete response in metastatic poorly-differentiated neuroendocrine rectal carcinoma with multimodal approach: A case report

della Torre S *et al.* Neuroendocrine tumor and multimodal therapeutic approach

Abstract

BACKGROUND

Neuroendocrine gastrointestinal tumors (NETs) are rare and they have different natural behavior. Surgery is the gold standard treatment in local disease while radiotherapy has been demonstrated inefficacy. Poorly differentiated neuroendocrine carcinomas (NECs) represent only 5–10% among digestive NETS. Due to aggressive growth and rapid metastatic diffusion, early diagnosis and multidisciplinary approach are mandatory.

The role of surgery and radiotherapy in this setting is still debated. Chemotherapy

Match Overview

1	Internet 58 words crawled on 01-Jan-2021 www.wjgnet.com	3%
2	Internet 54 words crawled on 02-Nov-2020 erc.bioscientifica.com	3%
3	Internet 21 words crawled on 21-Sep-2020 insights.ovid.com	1%

国内版 国际版

ig

Long term complete response in metastatic poorly-differentiated ne



ALL

IMAGES

VIDEOS

51,400 Results

Any time ▾

Long-term control with chemoradiation of initially ...

<https://www.ncbi.nlm.nih.gov/pubmed/30902067>

Mar 23, 2019 · The patient has been recurrence-free for more than 5 years. **CONCLUSIONS:** Extended **long-term control of a poorly differentiated metastatic (stage IV) mixed adenoneuroendocrine carcinoma** is rare. The **multimodal first- and second-line regimens of radiotherapy and chemotherapy** described in this **case report** represent a new therapeutic **approach**.

Cited by: 1

Author: S. Semrau, A. Agaimy, M. Pavel, D. Lubg...

Publish Year: 2019

[PDF] Long-term control with chemoradiation of initially ...

<https://jmedicalcasereports.biomedcentral.com/track/pdf/10.1186/s13256-019-1995-x>

Conclusions: Extended **long-term control of a poorly differentiated metastatic (stage IV) mixed adenoneuroendocrine carcinoma** is rare. The **multimodal first- and second-line regimens of radiotherapy and chemotherapy** described in this **case report** represent a new therapeutic **approach**. Encouraged by the results in this **case**, we compiled a review of the

Long-term control with chemoradiation of initially ...

<https://jmedicalcasereports.biomedcentral.com/...> ▾

Mar 23, 2019 · The patient has been recurrence-free for more than 5 years. Extended **long-term control of a poorly differentiated metastatic (stage IV) mixed adenoneuroendocrine carcinoma** is rare. The **multimodal first- and second-line regimens of radiotherapy and chemotherapy** described in this **case report** represent a new therapeutic **approach**.

Cited by: 1

Author: S. Semrau, A. Agaimy, M. Pavel, D. Lubg...

Publish Year: 2019

Long term outcomes of neuroendocrine carcinomas (high ...

<https://www.sciencedirect.com/science/article/pii/S1878788613001689>

Feb 01, 2014 · Introduction. **Neuroendocrine carcinomas (NEC)** of the colon and rectum are rare, extremely aggressive neoplasms with a high mortality rate. The term, **neuroendocrine carcinoma** or high-grade **neuroendocrine tumor** describes malignant epithelial tumors with evidence of **neuroendocrine** differentiation based on light microscopic, ultrastructural, or immunohistochemical ...

Cited by: 23

Author: E. Aytac, Y. Ozdemir, G. Ozuner

Publish Year: 2014