

国内版 国际版

ing

Transperineal core needle biopsy of a rectal subepithelial lesion guic



ALL IMAGES VIDEOS

14,200 Results Any time Open links in new tab

Transperineal In-Bore 3-T MR Imaging-guided Prostate ...

<https://pubs.rsna.org/doi/10.1148/radiol.14140221>

Inclusion criteria were at least one suspected lesion at typical 3-T multiparametric MR imaging and at least one of the following: (a) elevated prostate-specific antigen (PSA), (b) prior negative transrectal US-guided biopsy, or (c) inability to undergo transrectal biopsy owing to rectal surgery. Exclusion criterion was prior prostatectomy or ...

Cited by: 76 Author: Tobias Penzkofer, Kemal Tuncali, Andriy ...

Publish Year: 2015

Contrast enhanced ultrasound guided biopsy shows higher ...

<https://tcr.amegroups.com/article/view/7298/6781>

The purpose of this study was to compare the diagnostic accuracy of conventional ultrasound (US) guided vs. contrast-enhanced ultrasound (CEUS)-guided core needle biopsy for GISTs. In our series, all patients were scheduled for Imatinib treatment ...

Images of transperineal Core Needle Biopsy of A Rectal Sube...

<bing.com/images>



See all images >

Evaluation of MRI/Ultrasound Fusion-Guided Prostate Biopsy ...

<https://www.hindawi.com/journals/bmri/2017/2176471>

< Abstract Introduction Materials and Methods Results Discu >

ALL IMAGES VIDEOS

255 Results Any time ▾

Anorectal ultrasound for neoplastic and inflammatory lesions

https://www.researchgate.net/publication/7302499_A...
Endorectal ultrasound (ERUS) and endoanal ultrasound (ERAS) are also used in the diagnosis of benign mucosal lesions, fistula in ano, fecal incontinence, anorectal abscesses, and extrarectal masses.

Role of ultrasound in colorectal diseases. - Abstract ...

europepmc.org/abstract/MED/27920469
New ultrasound techniques such as elastography, contrast enhanced and Doppler ultrasound, mini-probes rectal and transperineal ultrasonography have broadened the indication. It gives an overview of the methodology of various ultrasound examinations, presents the morphology of normal bowel wall and the typical changes in different colonic diseases.

Transperineal ultrasound in the assessment of haemorrhoids ...

<https://www.researchgate.net/publication/43135323...>
Transperineal ultrasound is an inexpensive, safe and painless technique that dynamically and non-invasively evaluates the anorectal area. It has multiple indications, mainly in urology ...


Advances in endoscopic ultrasound imaging of colorectal ...

<https://www.researchgate.net/publication/291423578...>
Ultrasound is an undervalued non-invasive examination in the diagnosis of colonic diseases. It has been replaced by the considerably more expensive magnetic resonance imaging and computed

Search Tools

Turn off Hover Translation (关闭翻译)

激活 Windows
转到设置以激活 Windows。



Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 60432

Manuscript Type: CASE REPORT

Transperineal core-needlebiopsy of a rectal subepithelial lesion guided by endorectal ultrasound after contrast-enhanced ultrasound: A case report

Qiong Zhang *et al.* Transperineal CNB of rectal subepithelial lesion

Qiong Zhang, Jie-Ying Zhao, Hua Zhuang*, Chun-Yan Lu, Jin Yao, Yuan Luo, Yong-Yang Yu

Match Overview

1	Internet 20 words crawled on 24-Jul-2020 www.wjgnet.com	1%
2	Internet 19 words crawled on 12-Jun-2020 ebs.amegroups.com	1%
3	Crossref Posted Content 14 words Shunsuke Watanabe, Jun Miyoshi, Masao Toiki, Komei Kamayashi et al. "Effectiveness of introducing a 20-gauge cor..."	1%
4	Internet 13 words crawled on 20-Jun-2019 publishing.blab.core.windows.net	1%

激活 Windows
转到设置以激活 Windows。

Tests to Diagnose and Stage Prostate Cancer

<https://www.cancer.org/cancer/prostate-cancer...>

A **core needle biopsy** is the main method used to diagnose prostate cancer. It is usually done by a urologist. It is usually done by a urologist. During the **biopsy**, the doctor usually looks at the prostate with an imaging test such as transrectal **ultrasound (TRUS)** or MRI, or a ...

Screening Tests for Prostate Cancer · Stage · Can Prostate Cancer Be Found Early

Transperineal In-Bore 3-T MR Imaging-guided Prostate ...

<https://pubs.rsna.org/doi/10.1148/radiol.14140221>

Inclusion criteria were at least one suspected lesion at typical 3-T multiparametric MR imaging and at least one of the following: (a) elevated prostate-specific antigen (PSA), (b) prior negative transrectal **US-guided biopsy**, or (c) inability to undergo transrectal **biopsy** owing to **rectal surgery**. Exclusion criterion was prior prostatectomy or ...

Cited by: 79 Author: Tobias Penzkofer, Kemal Tuncali, Andriy ...
Publish Year: 2015

Images of Transperineal Core Needle Biopsy of A Rectal Sube...

<bing.com/images>



See all images >

PEOPLE ALSO ASK

Can prostate cancer be diagnosed with a biopsy?