

Risk perception and knowledge of COVID-19 in patients with celiac disease

Zhen J¹, Stefanolo JP², Temprano MP², Seiler CL¹, Caminero A¹, de-Madaria E³, Montoro Huguet M^{4,5}, Vivas S⁶, Niveloni S², Smecul E², Uscanga L⁷, Trucco E⁸, Lopez V⁸, Olano C⁸, Mansueto P⁹, Caroccio A⁹, Green PH¹⁰, Duerksen D¹¹, Day AS¹², Tye-Din JA¹³, Bai JC², Ciacchi C¹⁴, Verdu EF¹, Lebwohl B¹⁰, Pinto-Sanchez MI¹

Match Overview

1	Crossref 158 words J. Zhen, J.P. Stefanolo, M.P. Temprano, S. Tedesco et al. "THE RISK OF CONTRACTING COVID-19 IS NOT INC...	3%
2	Crossref 99 words Jamie Zhen, Juan Pablo Stefanolo, Maria de la Paz Temp rano, Sebastian Tedesco et al. "The Risk of Contracting ...	2%
3	Internet 18 words crawled on 11-Nov-2020 www.ifsqn.com	<1%
4	Internet 17 words crawled on 20-Aug-2020 eurjmedres.biomedcentral.com	<1%
5	Internet 17 words crawled on 29-Oct-2020 journals.plos.org	<1%
6	Internet 16 words crawled on 20-Jul-2020 helda.helsinki.fi	<1%
7	Crossref 14 words Sarah Dryhurst, Claudia R. Schneider, John Kerr, Alexa ra L. J. Freeman et al. "Risk perceptions of COVID-19 aro	<1%



ALL

IMAGES

VIDEOS

4,630,000 Results

Any time ▾

COVID-19 and celiac disease: Risks and precautions

<https://www.medicalnewstoday.com/articles/covid-19-and-celiac-disease> ▾

Apr 15, 2020 · As international headlines surround **the COVID-19 pandemic**, those with **celiac disease** may have concerns about what **the condition** means for ...

Author: Rachel Nall, MSN, CRNA

Knowledge and Perceptions of the COVID-19 Pandemic ...

<https://onlinelibrary.wiley.com/doi/abs/10.1002/mus.27130?af=R>

Introduction/Aims **Patients** with myasthenia gravis (MG) may be particularly vulnerable during the SARS-CoV-2 pandemic due to **risk** of worsening **disease** during infection, potential adverse impacts of ...

UMSOM researchers find greatest risk factors for death in ...

<https://www.news-medical.net/news/20201218/UMSOM...> ▾

Dec 19, 2020 · **Knowledge** is power in many ways, so I think understanding which hospitalized **COVID-19 patients** are at highest **risk** of mortality can help guide difficult treatment decisions."

Risk factors for COVID-19 vaccine acceptance

<https://www.news-medical.net/news/20201214/Risk...> ▾

Dec 14, 2020 · Overall, the important variables driving vaccine hesitancy in the UK were the individual's sex, age, **risk perception** about the virus, and the level of trust in WHO-backed **COVID-19** ...

Assessment of Knowledge, Attitudes, and Perception of ...

<https://link.springer.com/article/10.1007/s10900-020-00882-0> ▾

Jul 07, 2020 · The most common statements accepted by participants as causes of **perception of fear of COVID-19** infection were the following: **fear of** transmission of infection to their families (98.5%), the disease is highly transmissible (90.4%), COVID-19 new disease with no available vaccine (78.6%) or treatment (87%), the fatality of the disease (82.1%), **fear of** entering **COVID-19** isolation hospitals (86.5%), and stigma related to **COVID-19** ...

Cited by: 8

Author: Wafaa Yousif Abdel Wahed, Enas Mamd...

Publish Year: 2020

Covid-19's impact on cancer services and research

<https://www.news-medical.net/news/20201209/Covid...> ▾

Dec 10, 2020 · For cancer **patients**, this means understanding how **risk** factors such as cancer type, treatment type, immune system health and any other health issues influence their Covid-19 **risk**.

Risk perception and knowledge of COVID-19 in patients with celiac d



Sign in



ALL

IMAGES

VIDEOS

3,220,000 Results

Any time ▾

Open links in new tab



COVID-19 and celiac disease: Risks and precautions

<https://www.medicalnewstoday.com/articles/covid-19-and-celiac-disease> ▾

As international headlines surround the **COVID-19 pandemic**, those with **celiac disease** may have concerns about what the **condition** means for them and their **dietary treatments**. Currently, there is no...

Knowledge, perceptions and preventive practices towards ...

<https://pubmed.ncbi.nlm.nih.gov/32437432>

Results: Of the 247 respondents, 205 (83.0%) knew the main **clinical symptoms of COVID-19**. 72.0% knew that older people who have **chronic illnesses** are at high **risk** of developing a severe form of **COVID-19**. About 95.1% knew that the **COVID-19 virus** spreads via respiratory droplets of infected people, while 77 (31.2%) of the respondents knew about the possibility of asymptomatic transmission.

Ethical dilemmas, perceived risk, and motivation among ...

<https://journals.sagepub.com/doi/10.1177/...>

Background: Positioned at the frontlines of the battle against **COVID-19 disease**, nurses are at increased **risk** of contraction, yet as they feel obligated to provide care, they also experience ethical...

Assessment of Knowledge, Attitudes, and Perception of ...

<https://link.springer.com/article/10.1007/s10900-020-00882-0> ▾

The most common statements accepted by participants as causes of **perception of fear of COVID-19**

Search Tools

[Turn off Hover Translation \(关闭取词\)](#)

激活 Windows

转到“设置”以激活 Windows。

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

2,900,000 Results

Any time ▾

COVID-19 and celiac disease: Risks and precautions

<https://www.medicalnewstoday.com/articles/covid-19-and-celiac-disease> ▾

Apr 15, 2020 · As international headlines surround the **COVID-19** pandemic, those with **celiac disease** may have concerns about what the **condition** means for them and their ...

Author: Rachel Nall, MSN, CRNA

COVID-19 Severity Is Tripled in the Diabetes Community: A ...

<https://care.diabetesjournals.org/content/44/2/526> ▾

Feb 01, 2021 · First, after adjustment for age, race, and other **risk** factors, the odds of a **COVID-19**-related hospitalization and greater illness severity for **patients** with type 1 **diabetes** are three- to fourfold higher than for **patients** without **diabetes**. This increased **risk** is approximately the same for **patients** with type 2 **diabetes** (Fig. 1 A and C).

Cited by: 3

Author: Justin M Gregory, James C Slaughter, Sara ...

Publish Year: 2021

COVID-19 Prevalence Maps and Public Knowledge, Risk ...

<https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2774639> ▾

Jan 06, 2021 · Individual **risk perception** was assessed on a scale of 9 items about **perceived susceptibility and severity** of getting **COVID-19**. Scores ranged from 1 to 7, with higher scores indicating greater **perceived susceptibility and severity** of getting **COVID-19**.

Author: Alistair Thorpe, Aaron M. Scherer, Paul ...

Publish Year: 2021

Some common-cold antibodies may help fight COVID-19 ...

<https://news.yahoo.com/common-cold-antibodies-may-help-202915732.html> ▾

Feb 24, 2021 · **Celiac disease** does not boost **COVID-19** risks **Celiac disease** does not increase **patients'** **risk for infection** or severe illness from the new coronavirus, new data show. People with **celiac disease** ...

Knowledge, perceptions and preventive practices towards ...

<https://pubmed.ncbi.nlm.nih.gov/32437432>

Results: Of the 247 respondents, 205 (83.0%) knew the main clinical symptoms of **COVID-19**. 72.0% knew that older people who have chronic illnesses are at high **risk** of developing a severe form of **COVID-19**.