

## Match Overview

1	<b>Internet</b> 86 words crawled on 16-Apr-2020 <a href="http://journals.sagepub.com">journals.sagepub.com</a>	3%
2	<b>Internet</b> 41 words crawled on 14-Aug-2020 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	2%
3	<b>Internet</b> 28 words crawled on 20-Jan-2018 <a href="http://noblehospitalspune.com">noblehospitalspune.com</a>	1%
4	<b>Crossref</b> 20 words Alexis M.P. Schizas, Andrew B. Williams. "Anorectal investigation", Elsevier BV, 2014	1%
5	<b>Internet</b> 19 words crawled on 07-Oct-2018 <a href="http://gut.bmj.com">gut.bmj.com</a>	1%
6	<b>Crossref</b> 18 words Ann Yates. "Faecal incontinence: a healthcare taboo". Briti	1%

**Name of Journal:** *World Journal of Gastroenterology*

**Manuscript NO:** 61458

**Manuscript Type:** MINIREVIEWS

**Diagnostic approach to faecal incontinence: What test and when to perform?**

Wisam Sbeit, Tawfik Khoury, Amir Mari

Diagnostic Approach to Faecal Incontinence: What Test and When



Sign in



ALL

IMAGES

VIDEOS

251,000 Results

Any time ▼

## [Faecal incontinence in adults - Diagnosis Approach | BMJ ...](https://bestpractice.bmj.com/topics/en-gb/840/diagnosis-approach)

<https://bestpractice.bmj.com/topics/en-gb/840/diagnosis-approach> ▼

Proctography is reserved for patients with suspected prolapse. Stool cultures, **blood tests** (full blood count, C-reactive protein), and rectal biopsy are necessary only if the history suggests incontinence is due to diarrhoea; otherwise they are not routinely requested. Anal manometry gives an objective assessment of anal continence.

## [Diagnosis of Fecal Incontinence | NIDDK](https://www.niddk.nih.gov/.../diagnosis)

<https://www.niddk.nih.gov/.../diagnosis> ▼

**electromyography**—a **test** that checks how well the muscles and nerves of your anus and pelvic floor are working; Endoscopy. Your doctor may perform an endoscopy to look inside your anus, rectum, and colon for signs of inflammation and digestive tract problems that may be causing your fecal incontinence. Endoscopies for fecal incontinence include

### PEOPLE ALSO ASK

What tests are done to check for fecal incontinence? ▼

What is faecal incontinence? ▼

## Fecal Incontinence

Also known as: soiling

Inability to control the passage of gas or stools through the anus.

### Common causes

Fecal incontinence is not always related to an underlying condition. It may be caused by:

- Constipation
- Diarrhea
- Scarred or stiffened rectal walls
- Surgery complications
- Vaginal child birth
- Mental retardation

### Treatment

**Self-treatment:** Self- care steps that may be helpful in some less- serious cases:

Diagnostic approach to faecal incontinence: What test and when to



ALL

IMAGES

VIDEOS

327,000 Results

Any time ▼

Open links in new tab



## [Diagnosis of Fecal Incontinence | NIDDK](https://www.niddk.nih.gov/.../diagnosis)

<https://www.niddk.nih.gov/.../diagnosis> ▼

**electromyography**—a test that checks how well the muscles and nerves of your anus and pelvic floor are working; Endoscopy. Your doctor may perform an endoscopy to look inside your anus, rectum, and colon for signs of inflammation and digestive tract problems that may be causing your fecal incontinence. Endoscopies for fecal incontinence include

## [Faecal incontinence in adults - Diagnosis Approach | BMJ ...](https://bestpractice.bmj.com/topics/en-gb/840/diagnosis-approach)

<https://bestpractice.bmj.com/topics/en-gb/840/diagnosis-approach> ▼

Approach. Faecal incontinence is a symptom, and the cause is often multifactorial. A structured approach to management is required, beginning with diagnosis and correction of reversible contributing factors. A baseline assessment should include history, examination, and **anoscopy/sigmoidoscopy**. A cognitive assessment may also be appropriate if learning difficulties, dementia, or behavioural issues are ...

### PEOPLE ALSO ASK

What tests are used to diagnose fecal incontinence?





ALL

IMAGES

VIDEOS

218,000 Results

Any time ▾

Open links in new tab



## Diagnosis of Fecal Incontinence | NIDDK

<https://www.niddk.nih.gov/.../diagnosis> ▾

**electromyography**—a test that checks how well the muscles and nerves of your anus and pelvic floor are working; Endoscopy. Your doctor may perform an endoscopy to look inside your anus, rectum, and colon for signs of inflammation and digestive tract problems that may be causing your fecal incontinence. Endoscopies for fecal incontinence include

## Fecal incontinence - Diagnosis and treatment - Mayo Clinic

<https://www.mayoclinic.org/diseases-conditions/...> ▾

Diagnosis

Treatment

Clinical Trials

Lifestyle and Home Remedies



Your doctor will ask questions about your condition and perform a physical exam that usually includes a visual inspection of your anus. A probe may be used to examine this area for nerve damage. Normally, this touching causes your anal sphincter to contract and your anus to pucker.

[See more on mayoclinic.org](#)

### PEOPLE ALSO ASK

What tests are used to diagnose fecal incontinence?



How do doctors check for fecal incontinence?



What is faecal incontinence?



What is fecal leakage?



Feedback

## Fecal incontinence in adults - Diagnosis Approach | BMJ ...

<https://bestpractice.bmj.com/topics/en-us/840/diagnosis-approach> ▾

Proctography is reserved for patients with suspected prolapse. Stool cultures, **blood tests** (complete blood count, C-reactive protein), and **rectal biopsy** are necessary only if the history suggests incontinence is due to diarrhea; otherwise they are not routinely requested. Anal manometry gives an objective assessment of anal continence.

Faecal incontinence in adults

Nov 26, 2018

## Fecal Incontinence

Also known as: soiling

Inability to control the passage of gas or stools through the anus.

### Common causes

Fecal incontinence is not always related to an underlying condition. It may be caused by:

- Constipation
- Diarrhea
- Scarred or stiffened rectal walls
- Surgery complications
- Vaginal child birth
- Mental retardation

### Treatment

**Self-treatment:** Self-care steps that may be helpful in some less-serious cases:

- Regular exercise, consumption of high-fiber diet & plenty of fluids to prevent constipation
- Avoid straining while passing stools
- Adopt a sitting or squatting position to avoid straining

**See a doctor if you notice:**

- It after vaginal child birth
- It interferes with daily activities at home or workplace
- Social withdrawal or low self-esteem issues
- Difficulties in interpersonal relationships
- It in a patient having some other conditions like diabetes, infections, multiple sclerosis

**See a doctor immediately if you notice:**

- It occurs suddenly
- It following a head injury or accident
- It in children above the age of 3 years
- Pain while passing stools

### Related conditions

[See all \(4+\)](#)