

[A Small Bowel Ulcer due to Clopidogrel with ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6047549>

We report the first case of small bowel ulcers due to clopidogrel in a 74-year-old man. He presented with diarrhea and melena after having been taking low-dose aspirin (LDA) and clopidogrel. There was no evidence of bleeding in the stomach, duodenum, or colon. Capsule endoscopy showed multiple ulcers and erosions in the small intestine.

Author: Masahide Ebi, Satoshi Inoue, Tom... Publish Year: 2018

[Diagnosis and Management of Upper Gastrointestinal Bleeding](#)

<https://www.aafp.org/afp/2012/0301/p469.html>



🌱 Make a difference simply by se
MAYBE LAT

47,700 Results Any time

A Small Bowel Ulcer due to Clopidogrel with ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6047549
Jun 16, 2018 · We report the first case of small bowel ulcers due to clopidogrel in a 74-year-old man. He presented with diarrhea and melena after having been taking low-dose aspirin (LDA) and clopidogrel. There was no evidence of bleeding in the stomach, duodenum, or colon. Capsule endoscopy showed multiple ulcers and erosions in the small intestine.
Author: Masahide Ebi, Satoshi Inoue, Tomoya S... Publish Year: 2018

Clopidogrel Uses, Side Effects & Warnings - Drugs.com

https://www.drugs.com/mtm/clopidogrel.html
Jan 23, 2020 · You should not use clopidogrel if you are allergic to it, or if you have: any active bleeding, or a stomach ulcer or bleeding in the brain (such as from a head injury). Tell your doctor if you have ever had: an ulcer in your stomach or intestines; or a bleeding disorder or blood clotting disorder.

Before Taking This Medicine, How Should I Take Clopidogrel?, What Happens if I Miss A Dose?
You should not use clopidogrel if you are allergic to it, or if you have: 1. any active bleeding, or 2. a stomach ulcer or bleeding in the brain (s...
Follow all directions on your prescription label and read all medication guides or instruction sheets. Use these medicines exactly as directed. Clop...
Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

- PEOPLE ALSO ASK
Can clopidogrel cause stomach ulcers?
Does clopidogrel stop bleeding?
What is the initial assessment for bowel bleeding?
Can you take Plavix if you are allergic to clopidogrel?

clopidogrel and bleeding - MedHelp

https://healthquestions.medhelp.org/clopidogrel-and-bleeding
The prolonged use of clopidogrel (Plavix) greatly reduces the risk of late thrombosis in DES, but clopidogrel itself poses problems. Aside from the expense and the risk of bleeding posed by the long-term use of this drug, using clopidogrel (because of the pronounced bleeding tendency it causes) essentially precludes any invasive medical ...

Small bowel bleeding: a comprehensive review ...

https://academic.oup.com/gastro/article/2/4/262/574111
May 29, 2014 · The diagnostic yield of small bowel radiography has been reported to be 5–10% in patients with suspected small bowel bleeding [81, 82]. In a meta-analysis, the yields of small bowel barium radiography were 8% for any findings and 6% for clinically significant findings; in contrast, the yields of CE were 67% and 42%, respectively. Small bowel radiography is unlikely to be helpful in diagnosing vascular lesions, which are the most common cause of small bowel ...
Cited by: 64 Author: Deepak Ourjan, Vishal Sharma, Surinder S... Publish Year: 2014

Plavix Uses, Dosage & Side Effects - Drugs.com

https://www.drugs.com/plavix.html
Mar 02, 2020 · You should not use Plavix if you are allergic to clopidogrel, or if you have: any active bleeding, or a stomach ulcer or bleeding in the brain (such as from a head injury). Tell your doctor if you have ever had: an ulcer in your stomach or intestines, or a bleeding disorder or blood clotting disorder.

Before Taking This Medicine, How Should I Take Plavix?, Plavix Dosing Information
You should not use Plavix if you are allergic to clopidogrel, or if you have: 1. any active bleeding, or 2. a stomach ulcer or bleeding in the bra...
Take Plavix exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sh...
Usual Adult Dose for Acute Coronary Syndrome: Unstable Angina (UA)/Non-ST-Elevation Myocardial Infarction (NSTEMI) - Loading dose: 300 mg orally once...

Which lab tests should be performed in suspected peptic ...

https://www.medscape.com/answers/181753-13889
Mar 14, 2021 · Chan FK, Ching JY, Hung LC, et al. Clopidogrel versus aspirin and esomeprazole to prevent recurrent ulcer bleeding. N Engl J Med. 2005 Jan 20. 352(3):238-44. [Medline].

Plavix (Clopidogrel) - Side Effects, Interactions, Uses ...

https://www.everydayhealth.com/drugs/plavix
Jun 23, 2020 · You should not use clopidogrel if you are allergic to it, or if you have: any active bleeding, or a stomach ulcer or bleeding in the brain (such as from a head injury).

Management of bleeding and procedures in patients on ...

https://www.sciencedirect.com/science/article/pii/S0268960X1930133X
Jan 01, 2020 · They found no significant difference in bleeding rates between ticagrelor-treated patients in whom the drug was discontinued either >120 h pre-operatively or between 72 and 120 h, with increased bleeding at <72 h. Whereas the clopidogrel-treated patients had more bleeding episodes if clopidogrel was discontinued between 72 and 120 h compared with >120 h pre-operatively. Another study ...

Clopidogrel: Side Effects, Dosage, Uses, and More

https://www.healthline.com/health/clopidogrel-oral-tablet
Clopidogrel oral tablet is a prescription drug that's available as the brand-name drug Plavix. It's also available as a generic drug. Generic drugs usually cost less than the brand-name version.



Name of Journal: *World Journal of Clinical Cases*
Manuscript NO: 61612
Manuscript Type: CASE REPORT

Small bowel ulcer bleeding due to suspected clopidogrel use in a patient with clopidogrel resistance: A case report

Lee SH *et al.* Clopidogrel resistance presenting with hematochezia

Sang Hoon Lee, Dong Ryeol Ryu, Sung Joon Lee, Sung Chul Park, Byung Ryul Cho, Seung Koo Lee, Sang Ji Choi, Hyun Seok Cho

Abstract

BACKGROUND

Clopidogrel is a platelet aggregation inhibitor used for the management of cardiovascular disease. While antiplatelet therapy decreases cardiovascular events after successful coronary drug-eluting stent, it increases the risk of gastrointestinal (GI)

Match Overview

1	Internet 94 words crawled on 05-Mar-2021 www.wjnet.com	5%
2	Crossref 93 words Masahide Ebi, Satoshi Inoue, Tomoya Sugiyama, Kazuhiro Yamamoto et al. 'A Small Bowel Ulcer due to Clopidogr	5%
3	Crossref 71 words Tsai, Tsung-Jun, Kwok-Hung Lai, Ping-I Hsu, Chieh-Ku L in, Hoi-Hung Chan, Hsien-Chung Yu, Huey-Min Wang, Ku	4%
4	Crossref 40 words Sooyun Chang, Soo Jung Park, Sun Wook Kim, Moo-Nyu n Jin, Jung-Hee Lee, Hyun Ju Kim, Sung Pil Hong, Tae Il	2%
5	Internet 37 words crawled on 14-Mar-2020 www.science.gov	2%
6	Internet 33 words crawled on 05-Mar-2021 coak.info	2%
7	Internet 25 words crawled on 22-Feb-2011 clinicaltrials.gov	1%
8	Internet 21 words crawled on 28-Feb-2021 e-sc.org	1%

Small bowel ulcer bleeding due to suspected clopidogrel use in a p:



ALL IMAGES VIDEOS

49,300 Results Any time

A Small Bowel Ulcer due to Clopidogrel with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6047549>

We report the first case of small bowel ulcers due to clopidogrel in a 74-year-old man. He presented with diarrhea and melena after having been taking low-dose aspirin (LDA) and clopidogrel. There was no evidence of bleeding in the stomach, duodenum, or colon. Capsule endoscopy showed multiple ulcers and erosions in the small intestine.

Author: Masahide Ebi, Satoshi Inoue, Tomoya... Publish Year: 2018

Clopidogrel Uses, Side Effects & Warnings - Drugs.com

<https://www.drugs.com/mtm/clopidogrel.html>

Jan 23, 2020 · You should not use clopidogrel if you are allergic to it, or if you have: any active bleeding; or a stomach ulcer or bleeding in the brain (such as from a head injury). Tell your doctor if you have ever had: an ulcer in your stomach or intestines; or a bleeding ...

Before Taking This Medicine

You should not use clopidogrel if you are allergic to it, or if you have: 1. any active bleeding; or 2. a stomach ulcer or bleeding in the brain (s...

How Should I Take Clopidogrel?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use these medicines exactly as directed.Clop...

What Happens If I Miss A Dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

PEOPLE ALSO ASK

Can clopidogrel cause stomach ulcers?

Does clopidogrel stop bleeding?

What is the initial assessment for bowel bleeding?

Can you take Plavix if you are allergic to clopidogrel?

Feedback