

ALL IMAGES VIDEOS

Add the Give with Bing extension

1,450,000 Results Any time

Cirrhosis as new indication for statins | Gut

<https://gut.bmj.com/content/69/5/953>

May 01, 2020 · These have been summarised in a recent meta-analysis.⁹ Globally, these studies showed that patients with cirrhosis receiving statins had a lower rate of cirrhosis decompensation, lower incidence of liver cancer, lower rate of infections and increased survival.^{9 11 12 75 76} Of note, in the detailed analysis provided by Mchanly et al,⁷⁷ the effects of statins reducing mortality was ...

Cited by: 7 Author: Jaime Bosch, Jordi Gracia-Sancho, Juan ...
Publish Year: 2020

Statin therapy in patients with cirrhosis | Frontline ...

<https://fg.bmj.com/content/6/4/255>

Oct 01, 2015 · Improvement in portal hypertension, reduction in the incidence of hepatocellular carcinoma (HCC) and delays in hepatic decompensation have been associated with statin therapy among patients with cirrhosis.^{14–16} In this review, we will explore the burden of CVD among patients with cirrhosis, review available evidence on the safety and efficacy of statins in this population, and summarise recent

Cirrhosis of the Liver

Medical Condition

A degenerative disease of the liver resulting in scarring and liver failure.

Common (More than 200,000 cases per year in US)

Requires lab test or imaging

Treatments can help manage condition no known cure

Can last several years or be lifelong

Alcohol abuse and Hepatitis B & C virus infections are the most common causes of

ALL IMAGES VIDEOS

1,540,000 Results Any time ▾

The Use of Statins in Patients With Cirrhosis ...

<https://www.gastroenterologyandhepatology.net/...>

Addition of simvastatin to standard therapy for the prevention of variceal rebleeding does not reduce rebleeding but increases survival in patients with cirrhosis. Gastroenterology

EXPLORE FURTHER

Use of Statins in Patients with Chronic Liver Disease and ...

pubmed.ncbi.nlm.nih.gov

The Use of Statins in Patients With Chronic Liver Disease ...

pubmed.ncbi.nlm.nih.gov

Use of statins in patients with liver disease

pubmed.ncbi.nlm.nih.gov

Cirrhosis as new indication for statins | Gut

gut.bmj.com

Recommended to you based on what's popular • Feedback

Efficacy of Statin Association With Standard Treatment in ...

<https://clinicaltrials.gov/ct2/show/NCT01095185>

Mar 30, 2010 · Patients with liver cirrhosis and variceal bleeding with standard treatment will be stratified and after randomized to undergo either statins (20 mg for 15 days, and after 40 mg/day) or placebo. They will be followed up for 12 months.

PEOPLE ALSO ASK

Are Statins safe for cirrhosis? ▾

Are Statins safe for dialysis patients? ▾

Is statin harmful to liver? ▾

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 61954

Manuscript Type: MINIREVIEWS

Addition of statins to the standard treatment in patients with cirrhosis: Safety and efficacy

Muñoz AE *et al.* Statins in patients with cirrhosis

Abstract

This review summarizes the safety and efficacy of statins in patients with cirrhosis. Due to concerns about the safety of statins in patients with impaired liver function, they have recently been investigated as a potential treatment option in cirrhosis. The most clinically significant adverse event is statin-related myopathy, and this may be related to the high serum statin concentrations in the setting of severely impaired liver function. Rhabdomyolysis is the most serious and potentially life-threatening

Match Overview

Match	Source	Words	Match %
1	Crossref	159 words Juan G. Abráides, Candid Villanueva, Carles Araci, Juan Tu mes et al. "Addition of Simvastatin to Standard Therapy for t	2%
2	Crossref	156 words Elisa Pose, Jonel Trebicka, Rajeshwar P Mookerjee, Paol ... Angeli, Pere Ginés. "Statins: Old drugs as new therapy for li	2%
3	Crossref	141 words Alberto E. Muñoz, Florencia Pollarsky, Mónica Marino, María no Carlier et al. "Safety of Chronic Simvastatin Treatment in	2%
4	Internet	120 words crawled on 11-Aug-2016 www.vjgnet.com	2%
5	Crossref	117 words Jonathan C Price, Phyllis C Tien. "Editorial: Statins and Liver Disease: Is it Time to Recommend Statins to Prevent Liv...	2%
6	Internet	117 words crawled on 20-Apr-2013 ddrc.ac.ir	2%
7	Internet	109 words crawled on 26-Apr-2021 iibcbarcelona.eu	2%
8	Internet	74 words crawled on 30-Sep-2019	1%

国内版 国际版

Addition of statins to the standard treatment in patients with cirrho:



ALL IMAGES VIDEOS

529,000 Results Any time ▾

Conclusions: The addition of simvastatin to the standard therapy in patients with decompensated cirrhosis and cardiovascular risk factors was efficient as it decreased the patient's mortality. Furthermore, the simvastatin was safe as patients showed good tolerance, considering that they did not develop adverse effects or serious **adverse** effects.

Addition of simvastatin to the standard therapy increases ...

www.oatext.com/addition-of-simvastatin-to-the-standard-therapy-increases-surv...

Was this helpful?

PEOPLE ALSO ASK

How are statins used to treat cirrhosis of the liver? ▾

What is the concern related to statin-induced hepatotoxicity? ▾

Is it safe to take statins with NAFLD? ▾

Is there a link between statins and liver disease? ▾

Feedback

Efficacy of Statin Association With Standard Treatment in ...

<https://clinicaltrials.gov/ct2/show/NCT01095185> ▾

Mar 30, 2010 · Patients with liver cirrhosis and variceal bleeding with standard treatment will be stratified and after randomized to undergo either statins (20 mg for 15 days, and after 40 mg/day) or placebo. They will be followed up for 12 months.

Actual Enrollment : 150 participants

Allocation: Randomized

Intervention Model: Parallel Assignment

Study Type : Interventional (Clinical Trial)

Addition of simvastatin to the standard therapy increases ...

<https://www.oatext.com/addition-of-simvastatin-to...> ▾