

ALL IMAGES VIDEOS

90,800 Results Any time

Angiosarcoma - National Cancer Institute

https://www.cancer.gov/.../rare-tumors/rare-vascular-tumors/angiosarcoma

Angiosarcoma is a rare cancer that develops in the inner lining of blood vessels and lymph vessels. This cancer can occur anywhere in the body but most often is in the skin, breast, liver and spleen.

Primary hepatic malignant vascular tumors: a follow-up ...

https://pubmed.ncbi.nlm.nih.gov/32795351

Background: Owing to its low incidence, there is insufficient clinical awareness and diagnostic experience with primary hepatic malignant vascular tumors (PHMVTs). The aim of our study was to investigate the imaging and clinicopathological features of patients with ...

PEOPLE ALSO ASK

What are the symptoms of primary liver angiosarcoma?

Epithelioid hemangioendothelioma Rare Tumor

Epithelioid hemangioendothelioma is a rare tumor, first characterized by Sharon Weiss and Franz Enzinger that both clinically and histologically is intermediate between angiosarcoma and hemangioma. However, a distinct, disease-defining genetic alteration recently described for EHE indicates that it is an entirely separate entity from both angiosarcoma and hemangioma.

Wikipedia

### Malignant vascular tumors of liver in neonates - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S0022346805008365>

Jan 01, 2006 · Vascular tumors of the liver reported in infants and young children are infantile **hemangioendothelioma**, **cavernous hemangioma**, malignant epithelioid **hemangioendothelioma**, and angiosarcoma. The most common is infantile **hemangioendothelioma**, a lesion most often congenital, usually multifocal, frequently symptomatic in infancy and may be complicated by hemodynamic and ...

**Cited by:** 22      **Author:** Zafar Nazir, Shahid Pervez  
**Publish Year:** 2006

### Childhood Vascular Tumors Treatment (PDQ®)–Health ...

<https://www.cancer.gov/types/soft-tissue-sarcoma/...>

In one child diagnosed with angiosarcoma secondary to malignant transformation from infantile hemangioma, response to treatment with **bevacizumab**, a monoclonal antibody against vascular endothelial growth factor, combined with systemic chemotherapy, has been reported.[17,27] A report of eight cases of liver angiosarcoma in children highlighted the misuse of the term ...

### Images of Primary Malignant Vascular Tumors of the Liver in ...

<bing.com/images>



See all images >

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 62289

Manuscript Type: MINIREVIEWS

**Primary Malignant Vascular Tumors of the Liver in Children: Angiosarcoma and Epithelioid Hemangioendothelioma**

Malignant vascular tumors of the liver in children

Sami Bannoura, Juan Putra

Match Overview

Match	Source	Words	Percentage
1	Crossref Ely Coumoyer, Alyaa Al-Brateem, Elissa Engel, Guiraz Chaouy, Stacie Stapleton, Denise M. Adams. "Clinical o...	26	1%
2	Crossref Hu-Zhi Zhang, Lei Dong, Su-Ying Wang, Xiao-Qian Yang. "TFE3 rearranged epithelioid hemangioendothelioma of bon...	21	1%
3	Crossref Simone Hellmer, Geoffrey Andrieux, Jochen Hochrein, Philipp Kurz et al. "Epithelioid hemangioendotheliomas of the...	21	1%
4	Crossref Govind B. Chhabra, Iram Siddiqui, Kaitira M. Ingley, Ashish A. Gupta. "Rare malignant liver tumors in children". <i>Pediatr...</i>	20	1%
5	Crossref Anna McGuire, Israel Fernandez-Pineda, Steven J Fahrenholtz, Belinda Hsi Dickie. "Pediatric Hepatic Vascular Tumors".	18	1%
6	Internet created on 23-Apr-2020 <a href="http://www.archivesofpathology.org">www.archivesofpathology.org</a>	16	1%
7	Internet created on 18-Jun-2017 <a href="http://en.wikipedia.org">en.wikipedia.org</a>	15	1%
8	Crossref Leigh A. Compton, Leona A. Doyle. "Advances in the Genetic Characterization of Cutaneous Mesenchymal Neoplasms"	14	1%
9	Crossref <i>Surgical Pathology of Liver Tumors</i> , 2015.	14	1%

国内版 国际版

Primary malignant vascular tumors of the liver in children: Angiosar



ALL IMAGES VIDEOS

63,600 Results Any time

### Childhood Vascular Tumors Treatment (PDQ®)—Health ...

[https://www.cancer.gov/types/soft-tissue-sarcoma/...](https://www.cancer.gov/types/soft-tissue-sarcoma/)



In one child diagnosed with angiosarcoma secondary to malignant transformation from infantile hemangioma, response to treatment with bevacizumab, a monoclonal antibody against vascular endothelial growth factor, combined with systemic chemotherapy, has ...

### Malignant vascular tumors of liver in neonates - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S0022346805008365>

Jan 01, 2006 - Vascular tumors of the liver reported in infants and young children are infantile hemangioendothelioma, cavernous hemangioma, malignant epithelioid hemangioendothelioma, and angiosarcoma. The most common is infantile hemangioendothelioma, a lesion most often congenital, usually multifocal, frequently symptomatic in infancy and may be complicated by hemodynamic and ...


Cited by: 23

Author: Zafar Nazir, Shahid Pervez

Publish Year: 2006

### Images of Primary Malignant Vascular tumors of the Liver in ...

<bing.com/images>



See all images >

PEOPLE ALSO ASK