

Endoscopic Suturing for Weight Gain After Bariatric Surgery

<https://www.uclahealth.org/comet/endoscopic...> ▾

Endoscopic Suturing for Weight Gain After Bariatric Surgery. Studies show that more than 90 percent of bariatric patients lose 50 percent or more of their excessive body weight and are able to maintain a long-term weight loss of 50 percent excess body weight or more. (1) Of even greater significance, many bariatric surgery patients experience improvements in co-occurring disorders, ...

Endoscopic Treatment of Weight Regain Following Roux-en ...

<https://pubmed.ncbi.nlm.nih.gov/29082456>

Roux-en-Y gastric bypass (RYGB) is the most commonly performed bariatric procedure. Despite its high efficacy, some patients regain part of their lost weight. Several endoscopic therapies have been introduced as alternatives to treat weight regain, but most ...

Cited by: 37 Author: Vitor Ottoboni Brunaldi, Pichamol Jirapiny...
Publish Year: 2018

PEOPLE ALSO ASK

Can weight regain after bariatric surgery? ▾

What is the weight loss percentage for RYGB? ▾

What is the best weight loss medication? ▾

Feedback

Endoscopic Therapy for Weight Regain After Bariatric ...

<https://www.researchgate.net/publication/287280455...>

Given the proliferation of endoluminal therapies with early evidence suggesting safety and efficacy in the treatment of weight regain, it is likely that endoscopic revision will be an increasingly...

Endoscopic Therapy for Weight Regain After Bariatric ...

<https://abdominalkey.com/endoscopic-therapy-for...> ▾

Endoscopic Therapy for Weight Regain After Bariatric Surgery Fig. 18.1 The gastrojejunal

ALL IMAGES VIDEOS

143,000 Results Any time

Endoscopic treatment can help when the stomach stretches out, and weight is regained after surgery. Using a non-surgical, minimally invasive procedure, it is possible to reduce the size of the stomach pouch again. This can restore the more quick feeling of fullness, and encourage patients to eat smaller meals again.

Weight Regain After Bariatric Surgery - Dr. Reem Sharaiha drsharaiha.com/weight-regain-after-bariatric-surgery/

Was this helpful?

PEOPLE ALSO ASK

- What to do when you regain weight after bariatric surgery?
Why is bariatric surgery the best weight loss method?
What happens to your body when you gain weight after surgery?
How to reduce the need for revision surgery?

Feedback

Endoscopic Therapy for Weight Regain After Bariatric ...
https://link.springer.com/chapter/10.1007/978-1-4419-1710-2_18
Dec 27, 2012 Given the proliferation of endoluminal therapies with early evidence suggesting safety and efficacy in the treatment of weight regain, it is likely that endoscopic revision will be an increasingly prominent element in bariatric care.

Endoscopic Treatment of Weight Regain Following Roux-en-Y ...
https://pubmed.ncbi.nlm.nih.gov/29082456
Roux-en-Y gastric bypass (RYGB) is the most commonly performed bariatric procedure. Despite its high efficacy, some patients regain part of their lost weight. Several endoscopic therapies have been introduced as alternatives to treat weight regain, but most ...

Weight Regain After Bariatric Surgery: Prevalence ...
https://pubmed.ncbi.nlm.nih.gov/30168043
Recent findings: Given the high percentage of patients with weight regain after surgery, significant effort has been placed on developing treatment options in the last few years. Tremendous work has taken place in the realm of cognitive behavior therapy, appetite suppressants, and endoscopic procedures with the hope of reducing the need for revision surgery which can be associated with significant ...

Endoscopic Therapy for Weight Regain After Bariatric ...
https://www.researchgate.net/publication/287280455
Given the proliferation of endoluminal therapies with early evidence suggesting safety and efficacy in the treatment of weight regain, it is likely that endoscopic revision will be an increasingly ...

Endoscopic Therapy for Weight Regain After Bariatric ...
https://abdominalkey.com/endoscopic-therapy-for...
May 30, 2017 Endoscopic post-bypass revisions. Tech Gastrointest Endosc 2010;12 (3):160-6; with permission) The first published report in patients with dilated GJA and weight regain reported weight loss in 15 of 20 patients at 2 months [4]. A subsequent study of 28 patients with dilated GJA in 2007 resulted in loss of >75% of regained weight in 64% of patients in a mean 2.3 sessions.

Related searches for Endoscopic therapy of weight regain after ba...

- endoscopic weight loss surgery
endoscopic bariatric surgery
endoscopic bariatric therapy
endoscopic gastric surgery
endoscopic weight loss procedures
new weight loss surgery procedures
endoscopic gastric bypass procedure

Bariatric Surgery

Medical Specialty
A surgery that helps reduce food consumption and help weight loss by removing a part of stomach thereby reducing its size.
Type of procedure: Minimally invasive
Recovery time: Can take several weeks
Duration: Few hours
Hospital stay: Typically a few days

May treat

Obesity
Very common
Obesity is a medical condition in which excess body fat has ...
Read more

What to expect before procedure

- Inform the doctor about your medical history and go through all the required physical and pathological tests.
Counseling with a psychiatrist and a nutritionist is usually done before the surgery.
Do not smoke for at least 4 to 5 weeks before the surgery.
Do not consume any foods or beverages for at least 10 to 12 hours before the surgery.
Bowel clearing laxatives and enema will be given before the surgery.
Come prepared and pack your belongings for at least 3 days, accompanied with an adult family member.

What to expect during procedure

- Anesthesia is given to numb the site of surgery.
Minimal incisions are made and laparoscope is inserted through one.
Depending upon the type of surgery, structural changes are made.
Incisions are sealed with stitches and bandaged.

What to expect after procedure

- During the hospital stay after the surgery, monitoring for pulse, blood pressure and other details, are done.
After discharge, medications should be taken regularly.
For few days, only bland liquids can be taken, which will gradually increase in consistency. Regular food intake is permitted after several weeks of surgery.
Do not lift heavy weights.
Regularly follow up with the doctor and nutritionist.
Due to sudden weight loss, fatigue, body ache, hairloss, skin changes and irritation may be experienced in first few months after the surgery.

Risks and Complications

- Infection and bleeding at the surgical site
Internal bleeding
Lung or breathing problems
Perforation and leak from the bowel
Malabsorption of other nutrients
Loose, sagging skin
Blood clots
Slipping of gastric band leading to blocked stomach
Hypoglycemia, vomiting, ulcers are long term risks

People also search for

- Sleeve Gastrectomy
Bariatrics
Biliopancreatic Diversion with Duodenal Switch
Choline
Gastroplasty
See more

Data from Focus Medica. Reviewed by a panel of doctors. Learn more
Consult a medical professional for advice

Suggest an edit

Search Tools

Turn off Hover Translation (关闭翻译)

Name of Journal: *World Journal of Gastrointestinal Surgery*

Manuscript NO: 63049

Manuscript Type: MINIREVIEWS

Endoscopic therapy of weight regain after bariatric surgery

Bulajic M *et al.* Endoscopic therapy weight regain bariatric surgery

Milutin Bulajic, Salvatore Francesco Vadalà di Prampero, Ivo Boškoski, Guido Costamagna

Abstract

Weight regain following primary bariatric surgery occurs in a significant proportion of patients and is attributed to epidemiological, anatomical and metabolic factors. Surgical revision of these patients has significant risks and limited benefits. Endoscopic revisions that reduce gastric pouch size and diameter of the gastrojejunal anastomosis may offer

Match Overview

| | | |
|---|--|----|
| 1 | Crossref 121 words Daniel B. Mizgall, Aayad R. Alqanem, Barham K. Abu Dayeh, Mohamed Elahmedi et al. "Revisonal endoscopic st | 2% |
| 2 | Internet 97 words crawled on 15-Nov-2020 www.thieme-connect.com | 2% |
| 3 | Crossref 75 words Alex Marie Florent Heylen, Anja Jacobs, Monika Lybeer, Ruediger L. Probst. "The OTSC®-Clip in Revisional Enc... | 1% |
| 4 | Internet 72 words www.ncsl.nm.nih.gov | 1% |
| 5 | Crossref 67 words Pichamol Jirapinyo, Diogo T.H. de Moura, William Y. Dong, Galileu Farias, Christopher C. Thompson. "Dose respo | 1% |
| 6 | Crossref 66 words Diogo Tunes Hounneaux De Moura, Christopher C. Thompso. "Endoscopic management of weight regain follow ... | 1% |
| 7 | Crossref 66 words Rabindra R. Walten. "The Use of the Overstitch for Bar... ric Weight Loss". <i>Gastrointestinal Endoscopy Clinics of N</i> | 1% |
| 8 | Crossref 66 words Sharaha, Reem Z., Prashant Kadia, Nihel Kumta, Louis J. Aronne, and Michel Kahalish. "Endoscopic sleeve plic... | 1% |

国内版 国际版

Endoscopic therapy of weight regain after bariatric surgery



ALL IMAGES VIDEOS

145,000 Results Any time ▾

Roux-en-Y gastric bypass (RYGB) is the most commonly performed bariatric procedure. Despite its high efficacy, some patients regain part of their lost weight. Several endoscopic therapies have been introduced as alternatives to treat weight regain, but most of the **articles** are relatively small with unclear long-term data.

Cited by: 51

Publish Year: 2018

[Endoscopic Treatment of Weight Regain Following Roux-e...](#)

pubmed.ncbi.nlm.nih.gov/29082456/

Was this helpful?

PEOPLE ALSO ASK

What to do when you regain weight after bariatric surgery? ▾

Why is bariatric surgery the best weight loss method? ▾

What happens to your body when you gain weight after surgery? ▾

How to reduce the need for revision surgery? ▾

Feedback

[Endoscopic Therapy for Weight Regain After Bariatric ...](#)

https://link.springer.com/chapter/10.1007/978-1-4419-1710-2_18 ▾

Dec 27, 2012 · Given the proliferation of endoluminal therapies with early evidence suggesting safety and efficacy in the treatment of weight regain, it is likely that endoscopic revision will be an increasingly prominent element in bariatric care.

Cited by: 1

Author: Nitin Kumar, Christopher C. Thompson

Publish Year: 2013

[Endoscopic Treatment of Weight Regain Following Roux-en ...](#)