

World Journal of *Gastroenterology*

World J Gastroenterol 2021 August 28; 27(32): 5297-5459



OPINION REVIEW

- 5297** Management of Flood syndrome: What can we do better?
Strainiene S, Peculyte M, Strainys T, Stundiene I, Savlan I, Liakina V, Valantinas J

REVIEW

- 5306** Radiomics and machine learning applications in rectal cancer: Current update and future perspectives
Stanzione A, Verde F, Romeo V, Boccadifuoco F, Mainenti PP, Maurea S
- 5322** Could the burden of pancreatic cancer originate in childhood?
Diaconescu S, Gilcă-Blanariu GE, Poamaneagra S, Marginean O, Paduraru G, Stefanescu G

MINIREVIEWS

- 5341** Application of artificial intelligence in preoperative imaging of hepatocellular carcinoma: Current status and future perspectives
Feng B, Ma XH, Wang S, Cai W, Liu XB, Zhao XM
- 5351** Artificial intelligence application in diagnostic gastrointestinal endoscopy - Deus ex machina?
Correia FP, Lourenço LC
- 5362** Faecal microbiota transplantation enhances efficacy of immune checkpoint inhibitors therapy against cancer
Kang YB, Cai Y
- 5376** Immune checkpoint inhibitor-related hepatotoxicity: A review
Remash D, Prince DS, McKenzie C, Strasser SI, Kao S, Liu K

ORIGINAL ARTICLE

Basic Study

- 5392** Therapeutic effect of *Cistanche deserticola* on defecation in senile constipation rat model through stem cell factor/C-kit signaling pathway
Zhang X, Zheng FJ, Zhang Z
- 5404** Recombinant angiopoietin-like protein 4 attenuates intestinal barrier structure and function injury after ischemia/reperfusion
Wang ZY, Lin JY, Feng YR, Liu DS, Zhao XZ, Li T, Li SY, Sun JC, Li SF, Jia WY, Jing HR

Retrospective Study

- 5424** Prolonged survival in patients with hand-foot skin reaction secondary to cooperative sorafenib treatment
Ochi M, Kamoshida T, Araki M, Ikegami T

- 5438** Contrast-enhanced ultrasound imaging for intestinal lymphoma

Cui NY, Gong XT, Tian YT, Wang Y, Zhang R, Liu MJ, Han J, Wang B, Yang D

Observational Study

- 5448** Intestinal ischemic manifestations of SARS-CoV-2: Results from the ABDOCOVID multicentre study

Norsa L, Bonaffini PA, Caldato M, Bonifacio C, Sonzogni A, Indriolo A, Valle C, Furfaro F, Bonanomi A, Franco PN, Gori M, Smania V, Scaramella L, Forzenigo L, Vecchi M, Solbiati M, Costantino G, Danese S, D'Antiga L, Sironi S, Elli L

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Luca Mastracci, MD, Associate Professor, Anatomic Pathology, Department of Surgical Sciences and Integrated Diagnostics (DISC), University of Genova, Ospedale Policlinico San Martino, Largo Rosanna Benzi 10, Genova 16132, Italy. luca.mastracci@unige.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2021 edition of Journal Citation Report® cites the 2020 impact factor (IF) for WJG as 5.742; Journal Citation Indicator: 0.79; IF without journal self cites: 5.590; 5-year IF: 5.044; Ranking: 28 among 92 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2020 is 6.9 and Scopus CiteScore rank 2020: Gastroenterology is 19/136.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Jia-Hui Li; Production Department Director: Yu-Jie Ma; Editorial Office Director: Ze-Mao Gong.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, Subrata Ghosh

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

August 28, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Could the burden of pancreatic cancer originate in childhood?

Smaranda Diaconescu, Georgiana Emmanuela Gîlcă-Blanariu, Silvia Poamaneagra, Otilia Marginean, Gabriela Paduraru, Gabriela Stefanescu

ORCID number: Smaranda Diaconescu 0000-0001-8018-7191; Georgiana Emmanuela Gîlcă-Blanariu 0000-0002-5590-6462; Silvia Poamaneagra 0000-0001-7312-5937; Otilia Marginean 0000-0003-2313-7643; Gabriela Paduraru 0000-0002-6765-6018; Gabriela Stefanescu 0000-0002-1394-8648.

Author contributions: All authors contributed equally to the conception, design, preparation and writing of this review.

Conflict-of-interest statement: The authors have no conflicts of interest to declare for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Gastroenterology

Smaranda Diaconescu, Gabriela Paduraru, Department of Pediatrics, “Grigore T. Popa” University of Medicine and Pharmacy, Iasi 700115, Romania

Smaranda Diaconescu, Silvia Poamaneagra, Gabriela Paduraru, Department of Pediatric Gastroenterology, St Mary Emergency Children's Hospital, Iasi 700309, Romania

Georgiana Emmanuela Gîlcă-Blanariu, Gabriela Stefanescu, Department of Gastroenterology and Hepatology, “Grigore T. Popa” University of Medicine and Pharmacy, Iasi 700115, Romania

Georgiana Emmanuela Gîlcă-Blanariu, Gabriela Stefanescu, Department of Gastroenterology and Hepatology, “St. Spiridon” Emergency Hospital, Iasi 700111, Romania

Silvia Poamaneagra, Doctoral School, George Emil Palade University of Medicine, Pharmacy, Science and Technology, Targu Mures 540142, Romania

Otilia Marginean, Department of Pediatrics, Research Center of Disturbance of Growth and Development on Children-Belive, University of Medicine and Pharmacy “Victor Babes” Timisoara, Timisoara 300041, Romania

Otilia Marginean, First Clinic of Pediatrics, “Louis Turcanu” Emergency Children's Hospital, Timisoara 300011, Romania

Corresponding author: Smaranda Diaconescu, MD, PhD, Professor, Department of Pediatrics, “Grigore T. Popa” University of Medicine and Pharmacy, 16 Universității str, Iasi 700115, Romania. turti23@yahoo.com

Abstract

The presence of pancreatic cancer during childhood is extremely rare, and physicians may be tempted to overlook this diagnosis based on age criteria. However, there are primary malignant pancreatic tumors encountered in pediatric patients, such as pancreatoblastoma, and tumors considered benign in general but may present a malignant potential, such as the solid pseudo-papillary tumor, insulinoma, gastrinoma, and vasoactive intestinal peptide secreting tumor. Their early diagnosis and management are of paramount importance since the survival rates tend to differ for various types of these conditions. Many pediatric cancers may present pancreatic metastases, such as renal cell carcinoma, which may evolve with pancreatic metastatic disease even after two or more decades. Several childhood diseases may create a predisposition for the development of pancreatic cancer during adulthood; hence, there is a need for extensive screening strategies and complex programs to facilitate the transition from pediatric to adult

and hepatology

Country/Territory of origin:

Romania

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): C, C, C
Grade D (Fair): 0
Grade E (Poor): 0

Received: January 28, 2021

Peer-review started: January 28, 2021

First decision: May 13, 2021

Revised: June 8, 2021

Accepted: July 30, 2021

Article in press: July 30, 2021

Published online: August 28, 2021

P-Reviewer: Huang CY, Liu B, Morelli L, Zhao CF

S-Editor: Zhang H

L-Editor: Filipodia

P-Editor: Liu JH



healthcare. Nevertheless, genetic studies highlight the fact the specific gene mutations and family aggregations may be correlated with a special predisposition towards pancreatic cancer. This review aims to report the main pancreatic cancers diagnosed during childhood, the most important childhood diseases predisposing to the development of pancreatic malignancies, and the gene mutations associates with pancreatic malignant tumors.

Key Words: Pancreatic cancer; Pancreatic metastasis; Childhood; Adult; Screening; Transition

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Pancreatic malignant tumors are rarely found during childhood, their prognosis being linked to the type of tumor and its capacity to evolve towards metastasis. Also, there are types of cancer diagnosed in pediatric patients which may present with pancreatic metastasis later, during adulthood. Various conditions, when diagnosed during childhood, may be associated with the later onset of pancreatic cancer. Also, several genetic mutations have been linked to the development of pancreatic malignancies. We discuss here the above-mentioned topics in the context of a comprehensive literature review.

Citation: Diaconescu S, Gilcă-Blanariu GE, Poamaneagra S, Marginean O, Paduraru G, Stefanescu G. Could the burden of pancreatic cancer originate in childhood? *World J Gastroenterol* 2021; 27(32): 5322-5340

URL: <https://www.wjgnet.com/1007-9327/full/v27/i32/5322.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v27.i32.5322>

INTRODUCTION

In spite of technological, economic and healthcare progress, pancreatic cancer remains correlated with high mortality rates, its diagnosis at an early stage being of paramount importance for a favorable outcome, but this objective has proven exceedingly difficult to achieve. Researchers in the field have attempted to correlate the appearance of malignant pancreatic tumors with various local or systemic diseases and/or genetic mutations. The preventive management in cases of high risk for pancreatic cancer begins during childhood, by assessing and, whenever possible, treating the predisposing factors[1].

Pancreatic tumors are rarely found among the pediatric population; therefore, their study is limited to isolated case reports or case series. From a histological point of view, pancreatic tumors differ during childhood and adulthood, with a tendency for a better prognosis and clinical outcome at younger ages[2].

This review aims at reporting the primary pancreatic malignant tumors encountered during childhood, the pediatric cancers with the capacity to evolve towards pancreatic metastatic disease, the conditions diagnosable at a pediatric age that may predispose to pancreatic cancer during adulthood, and the most important gene mutations correlated with a predisposition towards pancreatic cancer (Table 1).

PANCREATIC MALIGNANT TUMORS IN CHILDREN

In general, pancreatic tumors are divided into epithelial (exocrine, of acinar, ductal, or undetermined cell origin; or endocrine) and non-epithelial types. The most frequent pancreatic tumor among children is of acinar cell origin, pancreatoblastoma. Solid pseudo-papillary tumors present a low malignant potential and are mostly encountered in young female patients. Rare types of pancreatic cancer may be found in children, such as acinar cell carcinoma and the extremely rare ductal adenocarcinomas. In rare cases, neonatal hypoglycemia can be caused by a focal or diffuse neuroendocrine adenoma of the gland[3].

Table 1 Pediatric pancreatic cancers

Pediatric pancreatic malignant tumors
Pancreatoblastoma
Solid pseudo-papillary tumors
Insulinoma
Gastrinoma
VIPoma
Acinar cell carcinoma
Ductal adenocarcinoma
Mucinous cystic neoplasm
Lymphoma: Hodgkin lymphoma, large cell lymphoma, Burkitt's lymphoma
Germ cell tumors
Primitive neuroectoderm tumors: extraosseous Ewing's sarcoma
Mesenchymal tumors: rhabdomyosarcoma, leiomyosarcoma, schwannoma
Pediatric cancers evolving with pancreatic metastatic disease
Renal cell carcinoma
Sarcomas
Colorectal carcinoma
Neuroblastoma
Diseases (diagnosable) in pediatric patients predisposing to pancreatic cancer at adult age
Obesity
Diabetes mellitus, glucose metabolism disturbances and insulin resistance
Chronic and hereditary pancreatitis
Cystic fibrosis
McCune Albright syndrome
Multiple endocrine neoplasia type I
Von Hippel Lindau disease
Li Fraumeni syndrome

The pancreas may be also involved into neoplastic processes arising from the adjacent structures, such as lymphomas, teratomas, rhabdomyosarcomas and primitive neuroectodermal tumors. Secondary involvement may be seen during metastatic disease originating from neuroblastomas[4].

The study based on the data provided by the United States' National Cancer Institute for the Surveillance, Epidemiology and End Result (SEER) Database confirmed the scarcity of malignant pancreatic tumors in the pediatric population; data collected from a 30-year period of time identified only 58 patients aged under 20 years suffering from malignant pancreatic tumors[4].

Pancreatoblastoma

Pancreatoblastoma was described for the first time in a 15-mo-old male patient in 1957, and was originally termed "infantile adenocarcinoma of the pancreas"[5]. Later, histological similarities were found between the newly discovered type of tumor and the undifferentiated or incompletely differentiated fetal acini at 7-8 wk of gestation in the embryo, and the term "pancreatoblastoma" was imposed[6].

Most cases of pancreatoblastomas are sporadic but there have been cases of congenital nature associated with Beckwith-Wiedemann syndrome (macrosomia, macroglossia, visceromegaly, omphalocele and an increased risk for embryonal tumors, such as nephroblastoma, hepatoblastoma, rhabdomyosarcoma and mostly cystic pancreatoblastoma)[7].

Mostly encountered in pediatric patients aged 0 to 9-years-old, pancreatoblastomas are exceptionally rare in adults.

Pancreatoblastoma usually appears as a large well-defined solitary mass, with lobulated margins ranging between 1.5 cm and 20 cm, and with half of the cases arising from the cephalo-pancreatic region. It may present as a cystic mass accompanied by necrosis, especially when associated with Beckwith-Wiedemann syndrome[3].

Although it has been mostly cited in the male population, Rasalkar *et al*[8] reported two cases of pancreatoblastoma in 2 girls, aged 9-years-old and 11-years-old. The symptomatology included pain and discomfort in the epigastric region, accompanied by a local lump and food intolerance. In both cases, the mass had disordered the local anatomy by displacing the splenic vessels, right kidney, the renal veins and the inferior vena cava. The tumor was located on the head of the pancreas in the first case and on the distal body and pancreatic tail in the second case.

Other symptoms associated with pancreatoblastomas include weight loss, anorexia, fatigue, lethargy, and jaundice in rare cases. Local invasion has been cited, mostly involving the adjacent pancreas, biliary tree, and the local vessels, including the portal and the mesenteric veins. The most common sites of metastases are the liver and abdominal lymph nodes, but there have been case reports of rarely located metastasis, such as to the lungs, brain, omentum, colon, spleen and adrenal glands[9].

Solid pseudo-papillary tumors

Pseudo-papillary tumors of the pancreas are rarely encountered tumors of exocrine origin, having a strong female predilection and being diagnosed during childhood in half of the cases[3]. The literature suggests that they present no preference regarding location on the pancreas, a low-grade malignant potential and a favorable long-term prognosis after surgical treatment[10].

The histological origin of these tumors is not yet clear but it has been linked to ductal cells, endocrine or multipotent stem cells. One important issue is the possibility of recurrence and, hence, the mandatory imaging follow-up. Authors have reported cases of metastasis in the liver, regional lymph nodes, omentum, mesentery and peritoneum[11].

In a 15-year retrospective study, the authors found 11 patients with pseudo-papillary tumors presenting with abdominal pain or palpable abdominal mass, digestive intolerance, and 1 case of traumatic tumor rupture leading to hemoperitoneum; two cases were diagnosed incidentally during medical examinations for other diseases. In this lot, preoperative serum tumor markers were assessed as follows: α -fetoprotein, carcinoembryonic antigen, carbohydrate antigen (CA) 125 were within normal ranges, whereas elevated levels of CA 19-9 were found in 3 of 10 patients[10].

Insulinoma

Insulinomas are particularly rare in children, either the sporadic form or when associated with multiple endocrine neoplasia type 1 (MEN1). Although more than 90% of insulinomas are benign, there have been cases of malignant insulinomas, characterized by the World Health Organization (WHO) according to their metastatic potential and risk of recurrence and mortality. A tumor size larger than 2 cm as well as the presence of cytokeratin 19 and tumor staging and grading with Ki 67 > 2% are reported predictors for the metastatic disease. The most common sites of metastasis are the regional lymph nodes, the liver and bone[12].

The clinical manifestations include symptoms associated with neuroglycopenia, such as loss of consciousness, confusion, dizziness and lethargy, or with adrenergic stimulation, such as hunger, weight gain, tachycardia, palpitations and arterial hypertension; hypoglycemia may be difficult to recognize in neonates and small children up to 2 years of age because of the poor adrenergic symptoms. In these cases, the symptomatology associates with seizures, apnea and lethargy[13].

In a 20-year retrospective study, the fact that metastatic liver disease may appear years after the initial diagnosis and presumed curative surgical treatment of insulinomas was highlighted. In these cases, the metastatic liver disease manifested as recurrent, severe hypoglycemic episodes leading to coma[14].

Gastrinoma

Pancreatic gastrinomas are, in frequency, the second mostly encountered pancreatic neuroendocrine tumors (pNETs) after insulinomas. In children, this is an uncommon tumor, its diagnosis being based on complex biological and imagistic investigations. Because of its rarity, many cases of pediatric gastrinomas remain undiagnosed for

years; evidence of digestive ulcers associated with elevated levels of fasting gastrin is usually diagnostic for a gastrinoma associated with Zollinger-Ellison syndrome, but these findings have to be correlated with advanced imaging investigations in order to assess tumor localization and invasion. In the pediatric population, the malignancy rate is close to 30%, poor prognosis being associated with tumor size (> 3 cm), metastatic disease (especially hepatic, lymph nodes and bone metastasis), female gender, inadequate gastric hypersecretion control as well as the histopathological features[15].

According to the WHO classification, gastrinomas present with different potential of malignization corresponding to the degree of differentiation, as follows: Well differentiated gastrinomas with benign/uncertain behavior; well-differentiated gastrinomas with low-grade malignant behavior; and, poorly differentiated gastrinomas with high-grade malignant potential[16].

Massaro *et al*[17] reported a case of a gastrinoma with hepatic metastasis in a child investigated for Zollinger-Ellison syndrome who eventually underwent orthoptic liver transplant.

The symptomatology of sporadic pancreatic gastrinomas includes epigastric pain due to severe peptic ulcer disease, heartburn, chronic diarrhea, gastrointestinal (GI) bleeding, nausea and vomiting. There are pancreatic gastrinomas associated with *MEN1* mutation, which may manifest clinically at younger ages, before hyperparathyroidism; hence, complex management including assessment of ionized calcium and serum parathormone levels is important[16].

Vasoactive intestinal peptide secreting tumors

The vasoactive intestinal peptide secreting tumors (VIPomas) gather together a group of neuroendocrine tumors characterized by the secretion of the vasoactive intestinal peptide, resulting in the “WDHA” syndrome (featuring achlorhydria, hypokalemia and watery diarrhea). The literature suggests that 60%-80% of VIPomas have malignant potential and present with metastasis at first diagnosis. During childhood, VIPomas are usually diagnosed between 2-4 years of age, but there have been cases at younger age (Reindl *et al*[18] start their series with a 2-mo-old patient).

Metastasis are mostly located in the liver, but rare case of lymph node, lung and kidney metastasis have been described[19].

Acinar cell carcinoma

Carcinoma originating from the acinar cells represents a rare type of pancreatic cancer, although it is more frequently encountered during childhood compared to ductal cell adenocarcinoma. Even though the acinar cells represent the most commonly found cells in the pancreas, their malignant transformation accounts for only 1% of pancreatic exocrine tumors[20].

The histological differentiation between acinar cell carcinoma and pancreatoblastoma may be difficult in children, as their macroscopic and microscopic features tend to be similar. Other distinct characteristics include the presence of calcifications in one-third of the cases and the tendency for intratumoral hemorrhage[21].

Illyés *et al*[22] report the particular case of a 10-year-old male who presented with clinical and biological manifestations associated with Cushing's syndrome and was later diagnosed with pancreatic acinar cell carcinoma invading the retroperitoneum which evolved with multiple metastasis.

Ductal adenocarcinoma

Ductal adenocarcinomas are exceptionally rare entities in childhood. In spite of some case reports before 1993 and considering the current advancements in medicine and immunohistochemistry, these cases are no longer considered ductal adenocarcinomas [23].

The literature suggests that ductal adenocarcinomas appearing at a young age may present with a tendency for poor differentiation and a high degree of metastasis. The 5-year survival rate is similar to that in adults (2%-4%), although there have been authors who supported a lower survival rate due to delayed diagnosis in children and young adults[3].

Mucinous cystic neoplasms

Pancreatic cystic lesions usually present with a good prognosis in children, but rare cases of malignant (high grade dysplasia) or premalignant (low or intermediate grade dysplasia) course have been described. Recent studies suggest that mucinous cystic neoplasm is asymptomatic and discovered incidentally, and the malignant potential has to be considered, especially in tumors larger than 4 cm[24].

Lymphoma

Lymphomas may originate from the pancreatic tissue itself or may involve the pancreas with the true origin lying in the peripancreatic lymph nodes. During childhood, the most common type of lymphoma involving the pancreas is non-Hodgkin lymphoma; pancreatic and peripancreatic invasion can be seen in cases of large cell lymphoma and Burkitt's lymphoma. The pancreas involvement may be suggested by the presence of the mass itself or by a diffuse infiltration of the gland accompanied by local edema due to a process of acute pancreatitis or tumor lysis syndrome. There have been case reports of primary pancreatic lymphomas at a pediatric age manifested with obstructive jaundice, misleading the diagnosis towards a false acute cholestatic hepatitis[25,26].

Intrinsic pancreatic lymphomas usually affect female patients and manifest with nonspecific symptoms associated with the mass effect (abdominal pain, digestive intolerance). Imaging studies do not offer specific information and in order to make a definitive differential diagnosis from other pancreatic masses, a histological exam remains mandatory[27].

The literature reports less than 10 cases of pancreatic lymphoma manifesting as acute pancreatitis at debut. Athmani *et al*[28] report the case of a 6-year-old female with a rapidly progressive jaundice; in that case, abdominal imaging showed an important increase in the pancreatic size, homogeneous hepatomegaly and intra- and extrahepatic biliary ducts' dilatation. Ultimately, the histopathological examination set the diagnosis of Burkitt's lymphoma.

Standard diagnostic criteria for primary pancreatic lymphoma include absence of superficial lymphadenopathy, no enlargement of mediastinal lymph nodes visible on chest radiography, normal leukocyte count, the main mass located in the pancreas with peripancreatic lymph nodes involvement, and the absence of hepatic or splenic involvement[29].

Germ cell tumors

Extragenital germ cell tumors represent 1%-3% of all childhood tumors. Even though their occurrence, originating from an upper abdominal organ, is exceedingly uncommon, they have been described on the head of the pancreas and causing biliary dilatation and mass-effect related symptoms. Based on the morphological characteristics and on the degree of differentiation, the WHO classifies teratomas as mature and immature[30].

Even though the malignant potential of pancreatic teratomas is yet to be established, 7%-10% of retroperitoneal teratomas with other localization are malignant[31]. For example, in other teratomas, the presence of yolk sac tumor microfoci may promote a malignant relapse after an incomplete surgical resection[32].

Primitive neuroectodermal tumors

Primitive neuroectodermal tumors usually originate in the bone and soft tissue, and there have been cases where they originated from solid organs containing neuroendocrine cells, like the pancreas, which accounts for 0.3% of primary pancreatic neoplasms at all ages[33].

In a literature review of extraosseous Ewing's sarcoma of the pancreas, Bose *et al*[34] pointed out the fact that the symptomatology usually associates with abdominal pain and jaundice; there were 2 particular cases in females of pediatric age where the tumor manifested with precocious puberty and 2 cases of GI bleeding and secondary iron deficiency anemia.

Primitive neuroectodermal tumors present an aggressive behavior, mostly due to the metastatic power; the most usual sites for metastasis are lungs, liver, bone and bone marrow.

Mesenchymal tumors

In children, the most common malignant mesenchymal tumor involving the pancreas is rhabdomyosarcoma. A percentage (specifically 15%) of rhabdomyosarcomas diagnosed during childhood originate from a site other than the head, neck, genitourinary tract and extremities. When situated in the abdomen, rhabdomyosarcomas can have originated from any part of the GI tract, including the extrahepatic biliary tract, pancreas and omentum; malignant ascites is an uncommon feature. An appreciable amount (specifically 14%) of these children present with metastatic disease at the initial diagnosis; the most common sites for metastasis are the lungs (36%), bone marrow (22%) and bones (7%)[35].

In rare cases, leiomyosarcomas may arise from the pancreatic duct and blood vessel walls within the pancreas; although rarely encountered, there have been cases of pancreatic leiomyosarcomas in children aged 14 and older[36].

Pancreatic schwannomas are rarely described in the literature, and more than 90% of them are benign tumors. However, there have been cases, mostly associated with von Recklinghausen disease, where the authors have described large schwannomas with malignant characteristics. Usually found on the pancreatic head and body, they usually manifest as nonspecific abdominal pain, weight loss, jaundice or GI bleeding [37].

Complete surgical resection is the treatment of choice in cases of pancreatic schwannomas, especially due to the fact that the character of malignancy cannot be established pre/during surgery and the fact that they do not respond to radiotherapy or chemotherapy[38].

Sheng *et al*[39] reported the first case of a low-grade malignancy pancreatic solitary fibrous tumor in a pediatric patient, a 14-mo-old male toddler.

PEDIATRIC CANCERS EVOLVING WITH PANCREATIC METASTATIC DISEASE

Metastatic pancreatic disease accounts for less than 2% of the entire pancreatic malignancies[40].

Pancreatic metastasis are rare entities that usually appear years after the diagnosis of the primary tumor. Following a topical literature review, authors reported that 62.6% of pancreatic metastasis appeared secondary to renal cell carcinoma, 7.2% secondary to sarcomas and 6.2% secondary to colorectal carcinoma[41]. Renal cell carcinoma is extremely rare in children, representing 1.8% to 6.3% of all types of renal cancers. Metastases originating from renal cell carcinomas usually present themselves as solitary or multiple masses inside the pancreas, which may occur even 20 years after the primary tumor manifestation. It is mandatory that the differential diagnosis of a pancreatic mass in patients with a history of renal cell carcinoma include secondary pancreatic metastasis[42].

Kim *et al*[43] reported the case of a 4-year-old male with stage 4 neuroblastoma, who presented with pancreatic metastasis manifesting as acute pancreatitis and rapidly progressive severe cholestasis. Farah *et al*[44] reported the case of a 15-year-old female who developed pancreatic metastasis due to an alveolar rhabdomyosarcoma located into the paranasal sinuses.

Pediatric pancreatic malignant tumors present with different degrees of aggressiveness and different potential for metastatic disease or local recurrences. The overall survival rates of pancreatic cancer diagnosed during childhood is not well established, but the 5-year survival rate for particular tumors ranges from more than 95% for some to as low as 2%-4% for ductal adenocarcinomas. For example, Bien *et al*[45] highlighted the fact that the 5-year survival rate of patients with pancreatoblastomas ranges between 30%-50%, despite surgical treatment and chemotherapy.

The 5-year survival rate for solid pseudo-papillary tumors has been reported between 95%-98% after complete surgical resection[46]. Primitive neuroectodermal tumors are highly aggressive and present with very poor prognosis; the 5-year survival rate is reported to be 50%[47].

Brecht *et al*[4] analyzed 5-year overall survival in a group of 228 patients under the age of 30 diagnosed with malignant pancreatic tumors (100 carcinomas, 85 endocrine tumors, 8 solid pseudopapillary neoplasms, 11 pancreatoblastomas). The data were extracted from the United States National Cancer Institute's SEER Public-use Database from 1973 to 2004. According to this study, 5-year overall survival (OAS) varied widely as follows: According to stage, it was 87%, 68%, 21 % for local ($n = 54$), regional ($n = 42$), distant metastatic disease ($n = 108$), respectively. According to histological types, OAS of patients with carcinoma was 33%, endocrine tumors 58%, solid pseudo-papillary tumor of the pancreas 88%, pancreatoblastomas 66%. Following multivariate analysis, tumor stage, histology and age group are important and independent predictors for outcome.

Similar results were recently reported by Picado *et al*[48] analyzing the Data from the National Cancer Database (2004-2014) which included 109 children with pancreatic tumors. The 5-year overall survival by tumor histology was 95% for pseudopapillary tumors, 75% for neuroblastomas, 70% for pancreatoblastomas, 51% for endocrine tumors, 43% for sarcomas, and 34% for adenocarcinomas. On multivariable analysis, the strongest predictor of survival was the surgical resection[48].

Another recently published study, performed on 65 patients under the age of 21 diagnosed with pancreatic neoplasms, who followed the outcome after pancreaticoduodenectomy, reported that the 5-year overall survival by tumor histology was 95% for pseudopapillary neoplasm, 75% for neuroblastomas, 70% for pancreatoblastomas, 51% for endocrine tumors, 43% for sarcomas, and 34% for adenocarcinomas. In this study, the outcome was primarily associated with histology[49].

The burden of a diagnosis such as pancreatic cancer is often associated with adulthood, and pediatricians are tempted to overlook this possibility, especially in a toddler or young child. Diagnostic delays/mistakes are not uncommon at this age and they lead to worse outcome due to the aggressiveness of the local mass and the metastatic character.

DISEASES (DIAGNOSABLE) IN PEDIATRIC PATIENTS PREDISPOSING TO PANCREATIC CANCER AT ADULT AGE

Obesity

Various studies have highlighted a connection between increased body mass index (BMI) and the risk for pancreatic cancer[50-52].

It has been observed that a BMI of at least 30 kg/m² was associated with a significantly increased risk of pancreatic cancer compared with a BMI of less than 23 kg/m² [relative risk (RR): 1.72, 95%CI: 1.19-2.48], while an inverse relationship was also highlighted for moderate physical activity when comparing the highest to the lowest categories (RR: 0.45, 95%CI: 0.29-0.70), particularly among those with a BMI of at least 25 kg/m²[53].

Moreover, it has been suggested that obese individuals develop pancreatic cancer earlier in life compared to patients with a normal weight, while also having lower survival rates following the diagnosis of pancreatic cancer[51,54].

Recent studies have also shown that overweight status and obesity in late adolescence and early adulthood are associated with an increased risk of this type of cancer[55,56].

Taking these aspects into consideration, researchers also questioned whether increased body weight during childhood would also be associated with adult pancreatic cancer. A prospective Danish cohort study including over 293000 individuals for whom data on height and BMI between the ages of 7 and 13 years were available, linked this information with data from Danish Cancer Registry. This study highlighted that increased BMI at every age from 7-13 years was significantly and positively associated with pancreatic cancer in adulthood, diagnosed up until age 70 [57].

Furthermore, the American National Institutes of Health-AARP (formerly known as the American Association of Retired Persons) Diet and Health Study Cohort and several pooled analyses of cohort studies also revealed that adolescent or early adulthood body size would be positively associated with pancreatic cancer[51,58].

BMI in children aged 7-13 was identified as being associated with the development of pancreatic cancer in adulthood[57].

These positive associations were not dependent of the evolution of BMI later in life, indicating that the risk was established to a certain extent in earlier life. Consequently, in the context of increased prevalence of childhood obesity, improved strategies for prevention and early management of childhood obesity should be considered to diminish the risk for developing pancreatic cancer in adulthood, but also for other various diseases including cardiovascular diseases and other types of neoplasia[59-61].

Diabetes mellitus, glucose metabolism, and insulin resistance

The relationship between diabetes and pancreatic cancer has been investigated extensively, since numerous epidemiologic studies have described an association between these two entities; a meta-analysis including 88 cohort and case-control studies identified an around 2-fold risk (RR: 2.08, 95%CI: 1.87-2.32) of pancreatic malignancy in diabetic patients compared to patients without diabetes[62-64].

However, the exact direction of this association is under debate, with some data suggesting that diabetes may be a consequence rather than a cause of pancreatic cancer, while others support the bidirectional relationship between the two entities, taking into account some pathophysiological background, such as the fact that insulin resistance, as well as obesity may be mediated by reduced levels of plasma adiponectin, a fat-derived hormone that has insulin-sensitizing and anti-inflammatory properties[65-67].

All these data have come from studies including adult populations and mainly focused on type 2 diabetes mellitus, with less data on type 1 diabetes mellitus. A systematic review of studies that assessed risk of pancreatic cancer in people with diabetes, specifically mentioning whether it was type 1 or type 2 diabetes, or a surrogate for type 1 diabetes, such as young age at onset, suggested that the risk of pancreatic cancer is comparably increased both for type 1 and type 2 diabetes. The results of this systematic review also underlined some causality within this relationship. With regard to type 2 diabetes, a bidirectional relationship can be called upon, meaning that insulin resistance and diabetes may be induced by undiagnosed cancer or precancerous conditions of the pancreas, but considering the increased risk of pancreatic cancer reported as 1.5- to 2-fold higher in type 2 diabetes even when impaired glucose tolerance is detected more than 10 years before onset of cancer, suggesting that reverse causality does not individually and exclusively explain the association[68,69].

With regard to type 1 diabetes, it is likely, considering the early age at its onset, that it precedes pancreatic cancer rather than the other way round; therefore, type 1 diabetes should be considered a risk factor for pancreatic cancer[70].

Large cohort studies indicated that cancer risk in type 1 diabetes is rather low, but other authors reported that this risk is comparable between type 1 and type 2 diabetes. As older studies report, type 1 diabetes associates an overall modestly increased cancer risk[71].

Both epidemiological and genetic studies have tried to prove the association between type 2 diabetes and pancreatic cancer. Pierce *et al*[72] analyzed the 37 risk alleles attributed to type 2 diabetes and found a significant positive association between 2 of these alleles and pancreatic cancer.

However, the currently available data is limited and further studies including large type 1 and 2 diabetes and nondiabetic cohorts are needed.

Chronic pancreatitis

Chronic pancreatitis is a risk factor for pancreatic adenocarcinoma, independent of etiology[73,74].

Although the exact pathophysiology is not yet clear, several signaling pathways have been identified as activated during pancreatic inflammation, in the context of repeated DNA damage, error-prone repair mechanisms, and the progressive accumulation of genetic mutations, ultimately stimulating the development of pancreatic cancer[75]. Pancreas precancerous histologic changes are associated with a sequential accumulation of genetic defects, namely pancreatic intraepithelial neoplasms, which are present in sporadic pancreatic adenocarcinomas and also in patients with a history of chronic pancreatitis. Mutations found in the early stages involve the *kRas* gene, followed by *p16/CDKN2A*, *TP53* and *SMAD4/DPC4* genes[76]. Mutations in all four genes have been recognized as driver mutations that trigger neoplastic transformation and tumor progression[77].

Non-hereditary chronic pancreatitis is also an important risk factor for pancreatic cancer[73,74]. Reports from the International Pancreatitis Study Group identified a cumulative risk of 1.8% at 10 years and 4% at 20 years for pancreatic cancer, independent of the etiology of chronic pancreatitis[78]. More recent data included in a meta-analysis has suggested that the risk of pancreatic cancer was elevated 16-fold in patients with more than 2 years since their diagnosis of chronic pancreatitis, and even though this risk declines over time, it persists even after long-term follow-up. After at least 9 years of follow-up from the time of diagnosis of chronic pancreatitis, this patient category still had an over 3-fold increased risk of pancreatic cancer compared to patients without chronic pancreatitis[79].

Although these data were obtained from an adult population, they raise concern regarding the pediatric population, especially when considering patients with hereditary pancreatitis. Pancreatitis is rare in the pediatric population and frequently reported as idiopathic pancreatitis, showing gaps in the identification of the etiology. Recently, it was shown that genetic factors play a far more important role in the development of chronic pancreatitis in infancy and childhood than previously thought [80].

Hereditary pancreatitis is defined as two first-degree relatives or at least three second-degree relatives in two or more generations with chronic pancreatitis for which there is no other etiology. Several mutations have been associated with hereditary pancreatitis, from the initial description of the mutation in the cationic trypsinogen gene (*PRSS1*) by Whitcomb *et al*[81] to mutations in the *SPINK1*, *CPA1* and *CFTR* genes[82-85].

Due to a lack of specific signs and even to a sometimes quiet presentation, hereditary pancreatitis is frequently diagnosed at an advanced stage[86].

In some sporadic cases, mutations compatible with hereditary pancreatitis can be found without a corresponding family history, which could be explained by inheritance from unaffected carrier parents or even spontaneous *de novo* mutations[75, 87].

Although progress has been made in diagnosing hereditary pancreatitis, the therapeutic pathway for this entity in children is still under debate[80]. Furthermore, counseling in patients with hereditary pancreatitis and their families should cover several aspects, from highly penetrant childhood-onset disease to a predisposition for pancreatic cancer in adulthood[88].

Studies are suggesting that *PRSS1* and *CFTR* gene mutations may not be directly linked with the development of pancreatic cancer. The onset of hereditary pancreatitis during childhood or early adulthood favors the perpetual exposure to chronic inflammation. The link between inflammation and cancer may not be limited to a subset of tumors but may be present within different types of cancer including lung, bladder, digestive, skin, and vulvar cancer. Hence, even though mutations in the *PRSS1* and *CFTR* genes are not directly associated with pancreatic cancer, chronic inflammation may favor tumor formation[89].

Beyond hereditary pancreatitis, other genetically determined conditions, including exocrine pancreatic insufficiency could evolve towards chronic pancreatitis and therefore potentially associate an increased risk for pancreatic cancer. Among these, Shwachman-Diamond syndrome is included. This a rare multi-organ recessive disease, mainly characterized by bone marrow failure leading to increased risk of transformation to myelodysplastic syndrome, skeletal defects, short stature, and pancreatic insufficiency contributing to the failure to thrive[90].

Cystic fibrosis

The cystic fibrosis phenotype results from mutations in the gene encoding the cystic fibrosis transmembrane conductance regulator protein encoded by the *CFTR* gene and is among the most common autosomal recessive disease in White people having an increased risk of pancreatic cancer[91].

A meta-analysis investigating the risk for various cancers in cystic fibrosis patients identified a standardized incidence ratio of 6.18 (95%CI: 1.31-29.27), which is 2-fold to 5-fold higher in patients who had undergone lung transplantation in this setting[92]. Since there is an excess risk of early-onset pancreatic cancer in patients with cystic fibrosis, interventions for avoiding this setting should be identified. These could range from tackling environmental factors to gene therapy targeting mucin genes. Considering that nutritional deficiencies represent another possible risk factor for cancer, appropriate nutritional interventions to avoid or correct deficits should also be considered[93].

Mccune-Albright syndrome

McCune-Albright syndrome (MAS) represents a rare disorder, characterized by variable phenotypic expressions, including fibrous dysplasia of bone, hyperfunctioning endocrinopathies, and café-au-lait macules, in the context of a somatic gain-of-function mutation in the *GNAS* gene, which encodes the cAMP-regulating protein G_{α_s} , taking place early in embryonic development[94]. These types of *GNAS* mutations were previously identified in various types of neoplasias (pituitary, thyroid, appendiceal, and gonadal tumors, and breast cancer)[95-99].

Since there is such a wide range of pathology associated with MAS, it is essential to define them all and to establish an optimal screening strategy and clinical management. The activating *GNAS* mutations have been identified as somatic driver mutations in sporadic intraductal papillary mucinous neoplasms (IPMNs) and various GI lesions, also suggesting a potential role in pancreatic and GI tumorigenesis[100].

In a cohort of patients with MAS, about 46% developed IPMNs, which have risk for malignant degeneration[100,101]. This can lead to IPMN-associated adenocarcinoma but also to developing concurrent or distinct ductal adenocarcinoma, a pattern which has been described in a 2%-9% of patients who are being followed for IPMN[102,103].

Consequently, GI manifestations in MAS are common and include precursory pancreatic lesions for pancreatic cancer; therefore, patients with MAS may benefit from evaluation for GI pathology. Even though the optimal care for pancreatic lesions in MAS is not clearly shaped, a thorough GI history-taking at every visit and consideration of imaging with abdominal magnetic resonance imaging/magnetic resonance cholangiopancreatography would be useful[100].

A small observational study describes the link between MAS and pancreatic cancer, liver adenoma and choledochal cysts *via* the cAMP pathway. Taking into consideration this association, the authors suggest that all patients suffering from MAS should benefit from routine screening by magnetic resonance imaging (MRI) and if no lesion is observed, surveillance MRI may be performed every 5 years[104].

MEN1

MEN1 mutations cause a rare hereditary tumor syndrome, with an autosomal dominant transmission and high degree of penetrance. *MEN1* tumors are caused by inactivating mutations of the tumor suppressor gene *MEN1*, and is characterized by a predisposition to a multitude of endocrine and nonendocrine tumors[105]. The classical phenotype includes hyperplasia and/or tumors of parathyroid, enteropancreatic, and/or anterior pituitary origin, with 30%–70% of patients developing enteropancreatic tumors, either functional or non-functional pancreatic endocrine pancreatic tumors upon reaching middle age[106]. Germline *MEN1* mutations have also been noted in families with only a parathyroid disorder, namely familial isolated primary hyperparathyroidism, but this type of mutation has also been reported in some cases of “sporadic” pNETs[107–109].

Considering the pattern of transmission and variety of manifestations, the Endocrine Society’s clinical practice guidelines for *MEN1* recommend a complex surveillance scheme commencing at early pediatric age, aiming to detect early and manage optimally *MEN1*-associated manifestations and tumors[110]. Recommended screening for pancreatic tumors in this setting begins early, at age 5 for insulinoma, with recommendation for yearly evaluation of fasting glucose and insulin level; for other pNETs, screening is recommended beginning at age 10, with evaluation of chromogranin A, pancreatic polypeptide, glucagon, vasoactive intestinal peptide levels and annual imaging, comprising magnetic resonance imaging, computed tomography or endoscopic ultrasound, considering that large pancreatic tumors may develop between 10 and 20 years of age[105].

Von Hippel-Lindau disease

Von Hippel–Lindau (vHL) disease is an autosomal dominant syndrome which occurs secondary to germline mutations in the *VHL* tumor suppressor gene, with patients being at risk of developing visceral cysts and tumors throughout the body[111]. Most commonly, the pancreas and kidneys are involved, with development of cystic lesions, but there is also an increased risk of developing neoplasms, often with clear cell features, including cerebellar and retinal hemangioblastomas, adrenal pheochromocytomas, clear cell renal cell carcinomas, pancreatic neuroendocrine tumors and pancreatic serous cystadenomas.

Although the most frequent pancreatic lesions include pancreatic cysts, which are found in 75% of vHL patients, one-fifth of the vHL patients will present pNETs, arising earlier than in non-syndromic patients, mostly in the fourth decade of life[112, 113].

Considering the increased potential for metastatic disease of pNETs in vHL patients and also the risk for metastasis generally associated with pNETs, regardless of the histologic grade, surgical resection is typically recommended for all pNETs greater than 2 cm[114]. However, taking into account the possibility of recurrent pNETs in vHL patients, in the context of the particular genetic background, the therapeutic decision must be well considered[115]. Although these lesions most commonly present during adulthood, screening and surveillance should begin in the pediatric years for patients with vHL disease.

Li Fraumeni syndrome

Li Fraumeni syndrome (LFS) is inherited in an autosomal dominant manner, which implies an increased risk for multiple primary cancers in affected individuals[116]. The most frequently implied cancers include bone and soft tissue sarcomas, leukemia, breast cancer, and brain tumors; subsequent evidence highlighted a broader spectrum of neoplasia, including cancers of the lung, digestive tract, prostate, ovary and pancreas, as well as lymphoma and melanoma[117,118].

Germline mutations in the tumor suppressor gene *TP53* are mainly incriminated for this syndrome; however, the absence of detectable germline *TP53* mutations in some LFS families suggests other genes are potentially involved in this syndrome[119].

There are various diagnostic criteria for the syndrome, such as the one included in the classical definition, which was afterwards extended for Li Fraumeni-like syndrome (an entity which shares some features of LFS but does not meet the strict LFS diagnostic criteria), in order to establish the criteria for *TP53* genetic testing[120]. Since

there is such a diversity of types of neoplasia which could arise in the context of LFS, we currently lack appropriate clinical surveillance strategies in this setting. Moreover, in the absence of a thorough knowledge on potential clinical benefits, psychosocial and economic impact of a comprehensive clinical surveillance protocol on early cancer detection, including pancreatic cancer screening, in asymptomatic *TP53* mutation carriers remains unknown.

OTHER GENE MUTATIONS ASSOCIATED WITH A PREDISPOSITION FOR PANCREATIC CANCER

Genetic risk factors play a major role in the development of pancreatic cancer, even if we are considering hereditary pancreatic cancer, familial pancreatic cancer or cases of pancreatic cancer associated with a familial cancer syndrome. The genetic predisposition for developing a particular malignancy has been of interest over the last decades, and several gene variants have been linked to an increased risk for pancreatic cancer, such as *BRCA1* and *BRCA2*, *STK11*, *PRSS1*, *PALB2*, *ATM*, *CDKN2A*, *APC*, *MLH1*, *MSH6*, *MSH2*, and *PMS2*[121].

However, this area of genetic research is still providing new information with the expectation to develop tests in order to assess the individual risk factors for pancreatic cancer, especially inside families with pancreatic cancer patients.

The definition of familial pancreatic cancer implies the existence of at least one pair of first-degree relatives suffering from pancreatic cancer, in the absence of a particular syndrome inside the family[122].

BRCA1 and *BRCA2* mutations are highly associated with inherited breast and ovarian cancer, but in carriers other types of cancer occur at higher rates than in the general population, including of pancreatic cancer. Brose *et al*[123] emphasize the fact that the general population risk of 1.3% transforms into a 2.8 RR for pancreatic cancer in people who carry the *BRCA1* mutation.

Familial atypical multiple mole melanoma syndrome is disease related to a cell cycle regulator gene for the p16 protein product, *CDKN2A*. Screening should begin early in childhood, in order to capture the early debut of both malignant melanoma and pancreatic cancer[124].

APC gene mutations are associated with the onset of familial adenomatous polyposis, a condition which, classically, increases the risk for colorectal cancer. Nowadays, evidence strongly suggests a correlation between inherited mutation of this gene and duodenal, thyroid, hepatic and pancreatic cancers[125].

The hereditary non-polyposis colorectal cancer known as Lynch syndrome also accounts for an increased risk for pancreatic cancer (1.3%-4%)[126]. Geary *et al*[127] pointed out the 30-fold increased risk for pancreatic cancer in patients with particular germline mutations in genes associated with the appearance of Lynch syndrome, such as *MLH1*, *MSH2*, *MSH6* and *PMS2*.

Complete family medical history is of paramount importance in assessing the particular risk for pancreatic cancer. In risk stratification, early molecular studies and germline mutation testing are mandatory.

CONCLUSION

Pancreatic cancer is generally considered an adulthood condition. In spite of this fact, its origins may lay in diseases or predisposing conditions which may be diagnosed at an early age, hence the need for rigorous screening strategies in patients with hereditary diseases connected with pancreatic cancer or familial aggregation of such conditions. Clinicians should be aware of rare conditions diagnosed during childhood, which may associate an increased risk for adult-onset pancreatic cancer. Also, the rare diagnosis of pancreatic cancer should be taken into consideration even at a pediatric age, especially when facing previously considered idiopathic acute pancreatitis or cholestatic hepatitis. The clinical features associated with pancreatic or peripancreatic masses may mimic other diseases and their management should include extensive imaging and histopathological examinations.

Malignant pancreatic tumors remain extremely rare in children and young adults and limited data on incidence and outcomes are available. Entities change over the age groups towards a predominantly increased number of carcinomas with worse outcome in older patients. The overall survival of pediatric patients with pancreatic

tumors is grim, and the survival rates tend to vary between different tumors. Given the rarity and aggressivity of many histological types of pancreatic tumors in children, in many instances it is difficult to assess whether there is any link between pancreatic cancer in childhood and adulthood.

International collaboration is needed to learn more about pediatric pancreatic tumors and to study the link between pancreatic tumors in childhood and adulthood.

Prophylaxis strategies should include weight control during childhood, screening for diabetes mellitus at younger ages, early routine imagistic investigations in cases with a predisposition towards pancreatic cancer or pancreatic metastasis. MRI and endoscopic ultrasonography are the recommended screening methods for pancreatic lesions. Also, screening strategies should include parents, siblings and children of patients with pancreatic cancer, especially in cases where there are more than 2 pancreatic cancer patients within the same biological family.

The implementation of transition programs from pediatric to adult healthcare services remains mandatory for these patients, namely childhood pancreatic cancer survivors or children with predisposing conditions for pancreatic malignancies.

REFERENCES

- 1 **Yu DC**, Kozakewich HP, Perez-Atayde AR, Shamberger RC, Weldon CB. Childhood pancreatic tumors: a single institution experience. *J Pediatr Surg* 2009; **44**: 2267-2272 [PMID: [20006007](#) DOI: [10.1016/j.jpedsurg.2009.07.078](#)]
- 2 **van den Akker M**, Angelini P, Taylor G, Chami R, Gerstle JT, Gupta A. Malignant pancreatic tumors in children: a single-institution series. *J Pediatr Surg* 2012; **47**: 681-687 [PMID: [22498381](#) DOI: [10.1016/j.jpedsurg.2011.11.046](#)]
- 3 **Chung EM**, Travis MD, Conran RM. Pancreatic tumors in children: radiologic-pathologic correlation. *Radiographics* 2006; **26**: 1211-1238 [PMID: [16844942](#) DOI: [10.1148/rg.264065012](#)]
- 4 **Brecht IB**, Schneider DT, Klöppel G, von Schweinitz D, Barthlen W, Hamre MR. Malignant pancreatic tumors in children and young adults: evaluation of 228 patients identified through the Surveillance, Epidemiology, and End Result (SEER) database. *Klin Padiatr* 2011; **223**: 341-345 [PMID: [22012608](#) DOI: [10.1055/s-0031-1287836](#)]
- 5 **BECKER WF**. Pancreatoduodenectomy for carcinoma of the pancreas in an infant; report of a case. *Ann Surg* 1957; **145**: 864-70; discussions, 870 [PMID: [13425296](#) DOI: [10.1097/0000658-195706000-00008](#)]
- 6 **Horie A**, Yano Y, Kotoo Y, Miwa A. Morphogenesis of pancreatoblastoma, infantile carcinoma of the pancreas: report of two cases. *Cancer* 1977; **39**: 247-254 [PMID: [188539](#) DOI: [10.1002/1097-0142\(197701\)39:1<247::aid-cnrcr2820390138>3.0.co;2-f](#)]
- 7 **Cheng H**, Yang S, Ren Q, Yang W, Han W, Chang X, Zhu Z, Qin H, Wang H. Pancreatectomies for pediatric pancreatic tumors: A single institute experience from 2007 to 2018. *J Pediatr Surg* 2020; **55**: 1722-1726 [PMID: [31575410](#) DOI: [10.1016/j.jpedsurg.2019.08.051](#)]
- 8 **Rasalkar DD**, Paunipagar BK, Chu WC. Two Children with Pancreatoblastomas. *J Hong Kong Col Radiol* 2010; **13**: 133-136
- 9 **Montemarano H**, Lonergan GJ, Bulas DI, Selby DM. Pancreatoblastoma: imaging findings in 10 patients and review of the literature. *Radiology* 2000; **214**: 476-482 [PMID: [10671596](#) DOI: [10.1148/radiology.214.2.r00fe36476](#)]
- 10 **Park JY**, Kim SG, Park J. Solid pseudopapillary tumor of the pancreas in children: 15-year experience at a single institution with assays using an immunohistochemical panel. *Ann Surg Treat Res* 2014; **86**: 130-135 [PMID: [24761421](#) DOI: [10.4174/astr.2014.86.3.130](#)]
- 11 **Guo N**, Zhou QB, Chen RF, Zou SQ, Li ZH, Lin Q, Wang J, Chen JS. Diagnosis and surgical treatment of solid pseudopapillary neoplasm of the pancreas: analysis of 24 cases. *Can J Surg* 2011; **54**: 368-374 [PMID: [21939604](#) DOI: [10.1503/cjs.011810](#)]
- 12 **Nagtegaal ID**, Odze RD, Klimstra D, Paradis V, Rugge M, Schirmacher P, Washington KM, Carneiro F, Cree IA; WHO Classification of Tumours Editorial Board. The 2019 WHO classification of tumours of the digestive system. *Histopathology* 2020; **76**: 182-188 [PMID: [31433515](#) DOI: [10.1111/his.13975](#)]
- 13 **Miron I**, Diaconescu S, Aprodu G, Ioniuc I, Diaconescu MR, Miron L. Diagnostic Difficulties in a Pediatric Insulinoma: A Case Report. *Medicine (Baltimore)* 2016; **95**: e3045 [PMID: [26986124](#) DOI: [10.1097/MD.0000000000003045](#)]
- 14 **Hirshberg B**, Cochran C, Skarulis MC, Libutti SK, Alexander HR, Wood BJ, Chang R, Kleiner DE, Gordon P. Malignant insulinoma: spectrum of unusual clinical features. *Cancer* 2005; **104**: 264-272 [PMID: [15937909](#) DOI: [10.1002/cncr.21179](#)]
- 15 **Nath AL**, Saxena NA, Kulkarni BK, Borwankar SS, Lahoti HN, Oak SN. Zollinger-Ellison Syndrome in a 12-year-old Child. *J Indian Assoc Pediatr Surg* 2017; **22**: 168-169 [PMID: [28694576](#) DOI: [10.4103/0971-9261.207623](#)]
- 16 **Jensen RT**, Niederle B, Mitry E, Ramage JK, Steinmuller T, Lewington V, Scarpa A, Sundin A, Perren A, Gross D, O'Connor JM, Pauwels S, Kloppel G; Frascati Consensus Conference;

- European Neuroendocrine Tumor Society. Gastrinoma (duodenal and pancreatic). *Neuroendocrinology* 2006; **84**: 173-182 [PMID: [17312377](#) DOI: [10.1159/000098009](#)]
- 17 **Massaro SA**, Emre SH. Metastatic gastrinoma in a pediatric patient with Zollinger-Ellison syndrome. *J Pediatr Hematol Oncol* 2014; **36**: e13-e15 [PMID: [23426004](#) DOI: [10.1097/MPH.0b013e318282dadf](#)]
 - 18 **Reindl T**, Degenhardt P, Luck W, Riebel T, Sarioglu N, Henze G, Driever PH. [The VIP-secreting tumor as a differential diagnosis of protracted diarrhea in pediatrics]. *Klin Padiatr* 2004; **216**: 264-269 [PMID: [15455292](#) DOI: [10.1055/s-2004-44901](#)]
 - 19 **Yeh PJ**, Chen SH, Lai JY, Lai MW, Chiu CH, Chao HC, Wu RC, Wang CJ, Chen CC. Rare Cases of Pediatric Vasoactive Intestinal Peptide Secreting Tumor With Literature Review: A Challenging Etiology of Chronic Diarrhea. *Front Pediatr* 2020; **8**: 430 [PMID: [32850544](#) DOI: [10.3389/fped.2020.00430](#)]
 - 20 **Al-Hader A**, Al-Rohil RN, Han H, Von Hoff D. Pancreatic acinar cell carcinoma: A review on molecular profiling of patient tumors. *World J Gastroenterol* 2017; **23**: 7945-7951 [PMID: [29259370](#) DOI: [10.3748/wjg.v23.i45.7945](#)]
 - 21 **Park M**, Koh KN, Kim BE, Im HJ, Kim DY, Seo JJ. Pancreatic neoplasms in childhood and adolescence. *J Pediatr Hematol Oncol* 2011; **33**: 295-300 [PMID: [21464765](#) DOI: [10.1097/MPH.0b013e318206990a](#)]
 - 22 **Illyés G**, Luczay A, Benyó G, Kálmán A, Borka K, Köves K, Rácz K, Tulassay T, Schaff Z. Cushing's syndrome in a child with pancreatic acinar cell carcinoma. *Endocr Pathol* 2007; **18**: 95-102 [PMID: [17917000](#) DOI: [10.1007/s12022-007-0018-4](#)]
 - 23 **Lüttges J**, Stigge C, Pacena M, Klöppel G. Rare ductal adenocarcinoma of the pancreas in patients younger than age 40 years. *Cancer* 2004; **100**: 173-182 [PMID: [14692038](#) DOI: [10.1002/cncr.11860](#)]
 - 24 **Luo JJ**, Baksh FK, Pfeifer JD, Eastman JT, Beyer FC 3rd, Dehner LP. Abdominal mucinous cystic neoplasm in a male child. *Pediatr Dev Pathol* 2008; **11**: 46-49 [PMID: [18237233](#) DOI: [10.2350/07-01-0220.1](#)]
 - 25 **Sood V**, Agrawal N, Alam S, Rawat D, Khanna R, Bansal K, Bihari C. Primary Pancreatic Lymphoma Simulating Acute Cholestatic Hepatitis in a 7-Year-Old Child. *ACG Case Rep J* 2015; **2**: 190-192 [PMID: [26157960](#) DOI: [10.14309/crj.2015.51](#)]
 - 26 **Shet NS**, Cole BL, Iyer RS. Imaging of pediatric pancreatic neoplasms with radiologic-histopathologic correlation. *AJR Am J Roentgenol* 2014; **202**: 1337-1348 [PMID: [24848833](#) DOI: [10.2214/AJR.13.11513](#)]
 - 27 **Arenas García BR**. [Primary pancreatic lymphoma in pediatric patients]. *Radiologia* 2007; **49**: 125-127 [PMID: [17403343](#) DOI: [10.1016/s0033-8338\(07\)73733-5](#)]
 - 28 **Athmani I**, Le Rouzic MA, Valduga J, Mansuy L, Contet A, Galloy MA, Fouyssac F, Chastagner P. Pediatric pancreatic Burkitt lymphoma with obstructive jaundice: case report of a six-year-old child. *Arch Clin Cases* 2017; **4** [DOI: [10.22551/2017.17.0404.10114](#)]
 - 29 **Amodio J**, Brodsky JE. Pediatric Burkitt lymphoma presenting as acute pancreatitis: MRI characteristics. *Pediatr Radiol* 2010; **40**: 770-772 [PMID: [20135116](#) DOI: [10.1007/s00247-009-1475-3](#)]
 - 30 **Wang J**, Yin Y, Cai Z, Shen C, Yin X, Chen X, Zhao Z, Zhang B. Pediatric pancreatic teratoma: A case report and literature review. *Medicine (Baltimore)* 2019; **98**: e18001 [PMID: [31725669](#) DOI: [10.1097/MD.00000000000018001](#)]
 - 31 **Zhou XH**, Ma JK, Valluru B, Sharma K, Liu L, Hu JB. Diagnosis and differentiation of mature cystic teratoma of pancreas from its mimics: A case report. *Medicine (Baltimore)* 2020; **99**: e23267 [PMID: [33217853](#) DOI: [10.1097/MD.00000000000023267](#)]
 - 32 **Harms D**, Zahn S, Göbel U, Schneider DT. Pathology and molecular biology of teratomas in childhood and adolescence. *Klin Padiatr* 2006; **218**: 296-302 [PMID: [17080330](#) DOI: [10.1055/s-2006-942271](#)]
 - 33 **Teixeira U**, Goldoni M, Unterleider M, Diedrich J, Balbinot D, Rodrigues P, Monteiro R, Gomes D, Sampaio J, Fontes P, Waechter F. Primitive Neuroectodermal Tumor of the Pancreas: A Case Report and Review of the Literature. *Case Rep Surg* 2015; **2015**: 276869 [PMID: [26101685](#) DOI: [10.1155/2015/276869](#)]
 - 34 **Bose P**, Murugan P, Gillies E, Holter JL. Extrasosseous Ewing's sarcoma of the pancreas. *Int J Clin Oncol* 2012; **17**: 399-406 [PMID: [21892669](#) DOI: [10.1007/s10147-011-0311-6](#)]
 - 35 **McCarville MB**, Spunt SL, Pappo AS. Rhabdomyosarcoma in pediatric patients: the good, the bad, and the unusual. *AJR Am J Roentgenol* 2001; **176**: 1563-1569 [PMID: [11373233](#) DOI: [10.2214/ajr.176.6.1761563](#)]
 - 36 **Riddle ND**, Quigley BC, Browarsky I, Bui MM. Leiomyosarcoma arising in the pancreatic duct: a case report and review of the current literature. *Case Rep Med* 2010; **2010**: 252364 [PMID: [20589089](#) DOI: [10.1155/2010/252364](#)]
 - 37 **Aggarwal G**, Satsangi B, Shukla S, Lahoti BK, Mathur RK, Maheshwari A. Rare asymptomatic presentations of schwannomas in early adolescence: three cases with review of literature. *Int J Surg* 2010; **8**: 203-206 [PMID: [20167297](#) DOI: [10.1016/j.ijsu.2010.01.012](#)]
 - 38 **Cayirli H**, Tanriverdi HI, Ozguven AA, Gunsar C, Ersoy B, Kandiloglu AR. Schwannoma Localized Retroperitoneally in a 14-Year-Old Boy. *Case Rep Pediatr* 2016; **2016**: 1210874 [PMID: [28078159](#) DOI: [10.1155/2016/1210874](#)]
 - 39 **Sheng Q**, Xu W, Liu J, Shen B, Deng X, Wu Y, Wu W, Yu S, Wang X, Lv Z. Pancreatic solitary

- fibrous tumor in a toddler managed by pancreaticoduodenectomy: a case report and review of the literature. *Onco Targets Ther* 2017; **10**: 1853-1858 [PMID: 28392706 DOI: 10.2147/OTT.S133650]
- 40 **Lee M**, Song JS, Hong SM, Jang SJ, Kim J, Song KB, Lee JH, Cho KJ. Sarcoma metastasis to the pancreas: experience at a single institution. *J Pathol Transl Med* 2020; **54**: 220-227 [PMID: 32311873 DOI: 10.4132/jptm.2020.03.04]
 - 41 **Adler H**, Redmond CE, Heneghan HM, Swan N, Maguire D, Traynor O, Hoti E, Geoghegan JG, Conlon KC. Pancreatectomy for metastatic disease: a systematic review. *Eur J Surg Oncol* 2014; **40**: 379-386 [PMID: 24462547 DOI: 10.1016/j.ejso.2013.12.022]
 - 42 **Abdellah A**, Selma K, Elamin M, Asmae T, Lamia R, Abderrahmane M, Sanaa el M, Hanan E, Tayeb K, Nouredine B. Renal cell carcinoma in children: case report and literature review. *Pan Afr Med J* 2015; **20**: 84 [PMID: 26090042 DOI: 10.11604/pamj.2015.20.84.5791]
 - 43 **Kim EY**, Yoo SY, Kim JH, Sung KW. Pancreatic metastasis in a child suffering with treated stage 4 neuroblastoma. *Korean J Radiol* 2008; **9**: 84-86 [PMID: 18253081 DOI: 10.3348/kjr.2008.9.1.84]
 - 44 **Farah RA**, Kamen BA. Parameningeal alveolar rhabdomyosarcoma with an isolated pancreatic metastasis. *Pediatr Hematol Oncol* 1999; **16**: 463-467 [PMID: 10505324 DOI: 10.1080/088800199277038]
 - 45 **Bien E**, Godzinski J, Dall'igna P, Defachelles AS, Stachowicz-Stencel T, Orbach D, Bisogno G, Cecchetto G, Warmann S, Ellerkamp V, Brennan B, Balcerska A, Rapala M, Brecht I, Schneider D, Ferrari A. Pancreatoblastoma: a report from the European cooperative study group for paediatric rare tumours (EXPeRT). *Eur J Cancer* 2011; **47**: 2347-2352 [PMID: 21696948 DOI: 10.1016/j.ejca.2011.05.022]
 - 46 **You L**, Yang F, Fu DL. Prediction of malignancy and adverse outcome of solid pseudopapillary tumor of the pancreas. *World J Gastrointest Oncol* 2018; **10**: 184-193 [PMID: 30079144 DOI: 10.4251/wjgo.v10.i7.184]
 - 47 **Hsieh L**, Burjonrappa S. Pediatric pancreatic tumors: a review of current concepts. *J Pancreas* 2016; **17**: 257-262
 - 48 **Picado O**, Ferrantella A, Zabalo C, Rao K, Thorson CM, Sola JE, Perez EA. Treatment patterns and outcomes for pancreatic tumors in children: an analysis of the National Cancer Database. *Pediatr Surg Int* 2020; **36**: 357-363 [PMID: 31989243 DOI: 10.1007/s00383-020-04617-z]
 - 49 **Vasudevan SA**, Ha TN, Zhu H, Heaton TE, LaQuaglia MP, Murphy JT, Barry WE, Goodhue C, Kim ES, Aldrink JH, Polites SF, Leraas HJ, Rice HE, Tracy ET, Lautz TB, Superina RA, Davidoff AM, Langham MR Jr, Murphy AJ, Bütter A, Davidson J, Glick RD, Grijalva J, Gow KW, Ehrlich PF, Newman EA, Lal DR, Malek MM, Le-Nguyen A, Piché N, Rothstein DH, Short SS, Meyers R, Dasgupta R. Pancreaticoduodenectomy for the treatment of pancreatic neoplasms in children: A Pediatric Surgical Oncology Research Collaborative study. *Pediatr Blood Cancer* 2020; **67**: e28425 [PMID: 32658372 DOI: 10.1002/pbc.28425]
 - 50 **Carreras-Torres R**, Johansson M, Gaborieau V, Haycock PC, Wade KH, Relton CL, Martin RM, Davey Smith G, Brennan P. The Role of Obesity, Type 2 Diabetes, and Metabolic Factors in Pancreatic Cancer: A Mendelian Randomization Study. *J Natl Cancer Inst* 2017; **109** [PMID: 28954281 DOI: 10.1093/jnci/djx012]
 - 51 **Li D**, Morris JS, Liu J, Hassan MM, Day RS, Bondy ML, Abbruzzese JL. Body mass index and risk, age of onset, and survival in patients with pancreatic cancer. *JAMA* 2009; **301**: 2553-2562 [PMID: 19549972 DOI: 10.1001/jama.2009.886]
 - 52 **Aune D**, Greenwood DC, Chan DS, Vieira R, Vieira AR, Navarro Rosenblatt DA, Cade JE, Burley VJ, Norat T. Body mass index, abdominal fatness and pancreatic cancer risk: a systematic review and non-linear dose-response meta-analysis of prospective studies. *Ann Oncol* 2012; **23**: 843-852 [PMID: 21890910 DOI: 10.1093/annonc/mdr398]
 - 53 **Michaud DS**, Giovannucci E, Willett WC, Colditz GA, Stampfer MJ, Fuchs CS. Physical activity, obesity, height, and the risk of pancreatic cancer. *JAMA* 2001; **286**: 921-929 [PMID: 11509056 DOI: 10.1001/jama.286.8.921]
 - 54 **Yuan C**, Bao Y, Wu C, Kraft P, Ogino S, Ng K, Qian ZR, Robinson DA, Stampfer MJ, Giovannucci EL, Wolpin BM. Prediagnostic body mass index and pancreatic cancer survival. *J Clin Oncol* 2013; **31**: 4229-4234 [PMID: 24145341 DOI: 10.1200/JCO.2013.51.7532]
 - 55 **Genkinger JM**, Kitahara CM, Bernstein L, Berrington de Gonzalez A, Brotzman M, Elena JW, Giles GG, Hartge P, Singh PN, Stolzenberg-Solomon RZ, Weiderpass E, Adami HO, Anderson KE, Beane-Freeman LE, Buring JE, Fraser GE, Fuchs CS, Gapstur SM, Gaziano JM, Helzlsouer KJ, Lacey JV Jr, Linet MS, Liu JJ, Park Y, Peters U, Purdue MP, Robien K, Schairer C, Sesso HD, Viswanathan K, White E, Wolk A, Wolpin BM, Zeleniuch-Jacquotte A, Jacobs EJ. Central adiposity, obesity during early adulthood, and pancreatic cancer mortality in a pooled analysis of cohort studies. *Ann Oncol* 2015; **26**: 2257-2266 [PMID: 26347100 DOI: 10.1093/annonc/mdv355]
 - 56 **Stolzenberg-Solomon RZ**, Schairer C, Moore S, Hollenbeck A, Silverman DT. Lifetime adiposity and risk of pancreatic cancer in the NIH-AARP Diet and Health Study cohort. *Am J Clin Nutr* 2013; **98**: 1057-1065 [PMID: 23985810 DOI: 10.3945/ajcn.113.058123]
 - 57 **Nogueira L**, Stolzenberg-Solomon R, Gamborg M, Sørensen TIA, Baker JL. Childhood body mass index and risk of adult pancreatic cancer. *Curr Dev Nutr* 2017; **1** [PMID: 29388617 DOI: 10.3945/cdn.117.001362]
 - 58 **Genkinger JM**, Spiegelman D, Anderson KE, Bernstein L, van den Brandt PA, Calle EE, English DR, Folsom AR, Freudenheim JL, Fuchs CS, Giles GG, Giovannucci E, Horn-Ross PL, Larsson SC, Leitzmann M, Männistö S, Marshall JR, Miller AB, Patel AV, Rohan TE, Stolzenberg-Solomon RZ,

- Verhage BA, Virtamo J, Willcox BJ, Wolk A, Ziegler RG, Smith-Warner SA. A pooled analysis of 14 cohort studies of anthropometric factors and pancreatic cancer risk. *Int J Cancer* 2011; **129**: 1708-1717 [PMID: [21105029](#) DOI: [10.1002/ijc.25794](#)]
- 59 **Wang Y**, Lim H. The global childhood obesity epidemic and the association between socio-economic status and childhood obesity. *Int Rev Psychiatry* 2012; **24**: 176-188 [PMID: [22724639](#) DOI: [10.3109/09540261.2012.688195](#)]
- 60 **Park MH**, Falconer C, Viner RM, Kinra S. The impact of childhood obesity on morbidity and mortality in adulthood: a systematic review. *Obes Rev* 2012; **13**: 985-1000 [PMID: [22731928](#) DOI: [10.1111/j.1467-789X.2012.01015.x](#)]
- 61 **Kitahara CM**, Gamborg M, Berrington de González A, Sørensen TI, Baker JL. Childhood height and body mass index were associated with risk of adult thyroid cancer in a large cohort study. *Cancer Res* 2014; **74**: 235-242 [PMID: [24247722](#) DOI: [10.1158/0008-5472.CAN-13-2228](#)]
- 62 **Aggarwal G**, Kamada P, Chari ST. Prevalence of diabetes mellitus in pancreatic cancer compared to common cancers. *Pancreas* 2013; **42**: 198-201 [PMID: [23000893](#) DOI: [10.1097/MPA.0b013e3182592c96](#)]
- 63 **Bosetti C**, Rosato V, Li D, Silverman D, Petersen GM, Bracci PM, Neale RE, Muscat J, Anderson K, Gallinger S, Olson SH, Miller AB, Bas Bueno-de-Mesquita H, Scelo G, Janout V, Holcatova I, Lagiou P, Serraino D, Lucenteforte E, Fabianova E, Baghurst PA, Zatonski W, Foretova L, Fontham E, Bamlet WR, Holly EA, Negri E, Hassan M, Prizment A, Cotterchio M, Cleary S, Kurtz RC, Maisonneuve P, Trichopoulos D, Polesel J, Duell EJ, Boffetta P, La Vecchia C, Ghadirian P. Diabetes, antidiabetic medications, and pancreatic cancer risk: an analysis from the International Pancreatic Cancer Case-Control Consortium. *Ann Oncol* 2014; **25**: 2065-2072 [PMID: [25057164](#) DOI: [10.1093/annonc/mdu276](#)]
- 64 **Batabyal P**, Vander Hoorn S, Christophi C, Nikfarjam M. Association of diabetes mellitus and pancreatic adenocarcinoma: a meta-analysis of 88 studies. *Ann Surg Oncol* 2014; **21**: 2453-2462 [PMID: [24609291](#) DOI: [10.1245/s10434-014-3625-6](#)]
- 65 **Chari ST**, Leibson CL, Rabe KG, Timmons LJ, Ransom J, de Andrade M, Petersen GM. Pancreatic cancer-associated diabetes mellitus: prevalence and temporal association with diagnosis of cancer. *Gastroenterology* 2008; **134**: 95-101 [PMID: [18061176](#) DOI: [10.1053/j.gastro.2007.10.040](#)]
- 66 **Salvatore T**, Marfella R, Rizzo MR, Sasso FC. Pancreatic cancer and diabetes: A two-way relationship in the perspective of diabetologist. *Int J Surg* 2015; **21** Suppl 1: S72-S77 [PMID: [26123386](#) DOI: [10.1016/j.ijso.2015.06.063](#)]
- 67 **Bao Y**, Giovannucci EL, Kraft P, Stampfer MJ, Ogino S, Ma J, Buring JE, Sesso HD, Lee IM, Gaziano JM, Rifai N, Pollak MN, Cochrane BB, Kaklamani V, Lin JH, Manson JE, Fuchs CS, Wolpin BM. A prospective study of plasma adiponectin and pancreatic cancer risk in five US cohorts. *J Natl Cancer Inst* 2013; **105**: 95-103 [PMID: [23243202](#) DOI: [10.1093/jnci/djs474](#)]
- 68 **Gullo L**, Pezzilli R, Morselli-Labate AM; Italian Pancreatic Cancer Study Group. Diabetes and the risk of pancreatic cancer. *N Engl J Med* 1994; **331**: 81-84 [PMID: [8208269](#) DOI: [10.1056/NEJM199407143310203](#)]
- 69 **Huxley R**, Ansary-Moghaddam A, Berrington de González A, Barzi F, Woodward M. Type-II diabetes and pancreatic cancer: a meta-analysis of 36 studies. *Br J Cancer* 2005; **92**: 2076-2083 [PMID: [15886696](#) DOI: [10.1038/sj.bjc.6602619](#)]
- 70 **Stevens RJ**, Roddam AW, Beral V. Pancreatic cancer in type 1 and young-onset diabetes: systematic review and meta-analysis. *Br J Cancer* 2007; **96**: 507-509 [PMID: [17224924](#) DOI: [10.1038/sj.bjc.6603571](#)]
- 71 **Zendejdel K**, Nyrén O, Ostenson CG, Adami HO, Ekblom A, Ye W. Cancer incidence in patients with type 1 diabetes mellitus: a population-based cohort study in Sweden. *J Natl Cancer Inst* 2003; **95**: 1797-1800 [PMID: [14652242](#) DOI: [10.1093/jnci/djg105](#)]
- 72 **Pierce BL**, Austin MA, Ahsan H. Association study of type 2 diabetes genetic susceptibility variants and risk of pancreatic cancer: an analysis of PanScan-I data. *Cancer Causes Control* 2011; **22**: 877-883 [PMID: [21445555](#) DOI: [10.1007/s10552-011-9760-5](#)]
- 73 **Bang UC**, Benfield T, Hyldstrup L, Bendtsen F, Beck Jensen JE. Mortality, cancer, and comorbidities associated with chronic pancreatitis: a Danish nationwide matched-cohort study. *Gastroenterology* 2014; **146**: 989-994 [PMID: [24389306](#) DOI: [10.1053/j.gastro.2013.12.033](#)]
- 74 **Duell EJ**, Lucenteforte E, Olson SH, Bracci PM, Li D, Risch HA, Silverman DT, Ji BT, Gallinger S, Holly EA, Fontham EH, Maisonneuve P, Bueno-de-Mesquita HB, Ghadirian P, Kurtz RC, Ludwig E, Yu H, Lowenfels AB, Seminara D, Petersen GM, La Vecchia C, Boffetta P. Pancreatitis and pancreatic cancer risk: a pooled analysis in the International Pancreatic Cancer Case-Control Consortium (PanC4). *Ann Oncol* 2012; **23**: 2964-2970 [PMID: [22767586](#) DOI: [10.1093/annonc/mds140](#)]
- 75 **Weiss FU**. Pancreatic cancer risk in hereditary pancreatitis. *Front Physiol* 2014; **5**: 70 [PMID: [24600409](#) DOI: [10.3389/fphys.2014.00070](#)]
- 76 **Hruban RH**, Takaori K, Klimstra DS, Adsay NV, Albores-Saavedra J, Biankin AV, Biankin SA, Compton C, Fukushima N, Furukawa T, Goggins M, Kato Y, Klöppel G, Longnecker DS, Lütges J, Maitra A, Offerhaus GJ, Shimizu M, Yonezawa S. An illustrated consensus on the classification of pancreatic intraepithelial neoplasia and intraductal papillary mucinous neoplasms. *Am J Surg Pathol* 2004; **28**: 977-987 [PMID: [15252303](#) DOI: [10.1097/01.pas.0000126675.59108.80](#)]
- 77 **Korc M**. Driver mutations: a roadmap for getting close and personal in pancreatic cancer. *Cancer Biol Ther* 2010; **10**: 588-591 [PMID: [20716952](#) DOI: [10.4161/cbt.10.6.13128](#)]

- 78 **Lowenfels AB**, Maisonneuve P, Cavallini G, Ammann RW, Lankisch PG, Andersen JR, Dimagno EP, Andrén-Sandberg A, Domellöf L. Pancreatitis and the risk of pancreatic cancer. International Pancreatitis Study Group. *N Engl J Med* 1993; **328**: 1433-1437 [PMID: [8479461](#) DOI: [10.1056/NEJM199305203282001](#)]
- 79 **Kirkegård J**, Mortensen FV, Cronin-Fenton D. Chronic Pancreatitis and Pancreatic Cancer Risk: A Systematic Review and Meta-analysis. *Am J Gastroenterol* 2017; **112**: 1366-1372 [PMID: [28762376](#) DOI: [10.1038/ajg.2017.218](#)]
- 80 **Kargl S**, Kienbauer M, Duba HC, Schöfl R, Pumberger W. Therapeutic step-up strategy for management of hereditary pancreatitis in children. *J Pediatr Surg* 2015; **50**: 511-514 [PMID: [25840052](#) DOI: [10.1016/j.jpedsurg.2014.05.016](#)]
- 81 **Whitcomb DC**, Gorry MC, Preston RA, Furey W, Sossenheimer MJ, Ulrich CD, Martin SP, Gates LK Jr, Amann ST, Toskes PP, Liddle R, McGrath K, Uomo G, Post JC, Ehrlich GD. Hereditary pancreatitis is caused by a mutation in the cationic trypsinogen gene. *Nat Genet* 1996; **14**: 141-145 [PMID: [8841182](#) DOI: [10.1038/ng1096-141](#)]
- 82 **Witt H**, Luck W, Hennies HC, Classen M, Kage A, Lass U, Landt O, Becker M. Mutations in the gene encoding the serine protease inhibitor, Kazal type 1 are associated with chronic pancreatitis. *Nat Genet* 2000; **25**: 213-216 [PMID: [10835640](#) DOI: [10.1038/76088](#)]
- 83 **Witt H**, Beer S, Rosendahl J, Chen JM, Chandak GR, Masamune A, Bence M, Szmola R, Oracz G, Macek M Jr, Bhatia E, Steigenberger S, Lasher D, Bühler F, Delaporte C, Tebbing J, Ludwig M, Pilsak C, Saum K, Bugert P, Masson E, Paliwal S, Bhaskar S, Sobczynska-Tomaszewska A, Bak D, Balascak I, Choudhuri G, Nageshwar Reddy D, Rao GV, Thomas V, Kume K, Nakano E, Kakuta Y, Shimosegawa T, Durko L, Szabó A, Schnür A, Hegyi P, Rakonczay Z Jr, Pfützer R, Schneider A, Groneberg DA, Braun M, Schmidt H, Witt U, Friess H, Algül H, Landt O, Schuelke M, Krüger R, Wiedenmann B, Schmidt F, Zimmer KP, Kovacs P, Stumvoll M, Blüher M, Müller T, Janecke A, Teich N, Grützmann R, Schulz HU, Mössner J, Keim V, Löhr M, Férec C, Sahin-Tóth M. Variants in CPA1 are strongly associated with early onset chronic pancreatitis. *Nat Genet* 2013; **45**: 1216-1220 [PMID: [23955596](#) DOI: [10.1038/ng.2730](#)]
- 84 **Sharer N**, Schwarz M, Malone G, Howarth A, Painter J, Super M, Braganza J. Mutations of the cystic fibrosis gene in patients with chronic pancreatitis. *N Engl J Med* 1998; **339**: 645-652 [PMID: [9725921](#) DOI: [10.1056/NEJM199809033391001](#)]
- 85 **Sobczyńska-Tomaszewska A**, Bak D, Oralewska B, Oracz G, Norek A, Czerska K, Mazurczak T, Teisseyre M, Socha J, Zagulski M, Bal J. Analysis of CFTR, SPINK1, PRSS1 and AAT mutations in children with acute or chronic pancreatitis. *J Pediatr Gastroenterol Nutr* 2006; **43**: 299-306 [PMID: [16954950](#) DOI: [10.1097/01.mpg.0000232570.48773.df](#)]
- 86 **Mastoraki A**, Tzortzopoulou A, Tsela S, Danias N, Sakorafas G, Smyrniotis V, Arkadopoulos N. Hereditary pancreatitis: dilemmas in differential diagnosis and therapeutic approach. *J Gastrointest Cancer* 2014; **45**: 22-26 [PMID: [24242859](#) DOI: [10.1007/s12029-013-9559-6](#)]
- 87 **Simon P**, Weiss FU, Zimmer KP, Rand S, Brinkmann B, Domschke W, Lerch MM. Spontaneous and sporadic trypsinogen mutations in idiopathic pancreatitis. *JAMA* 2002; **288**: 2122 [PMID: [12413370](#) DOI: [10.1001/jama.288.17.2122](#)]
- 88 **Shelton CA**, Grubs RE, Umapathy C, Yadav D, Whitcomb DC. Impact of hereditary pancreatitis on patients and their families. *J Genet Couns* 2020; **29**: 971-982 [PMID: [32026589](#) DOI: [10.1002/jgc4.1221](#)]
- 89 **Kong X**, Sun T, Kong F, Du Y, Li Z. Chronic Pancreatitis and Pancreatic Cancer. *Gastrointest Tumors* 2014; **1**: 123-134 [PMID: [26674754](#) DOI: [10.1159/000365306](#)]
- 90 **Warren AJ**. Molecular basis of the human ribosomopathy Shwachman-Diamond syndrome. *Adv Biol Regul* 2018; **67**: 109-127 [PMID: [28942353](#) DOI: [10.1016/j.bior.2017.09.002](#)]
- 91 **Slæ M**, Wilschanski M. Cystic fibrosis: a gastrointestinal cancer syndrome. *Lancet Oncol* 2018; **19**: 719-720 [PMID: [29706381](#) DOI: [10.1016/S1470-2045\(18\)30250-X](#)]
- 92 **Yamada A**, Komaki Y, Komaki F, Micic D, Zullow S, Sakuraba A. Risk of gastrointestinal cancers in patients with cystic fibrosis: a systematic review and meta-analysis. *Lancet Oncol* 2018; **19**: 758-767 [PMID: [29706374](#) DOI: [10.1016/S1470-2045\(18\)30188-8](#)]
- 93 **Maisonneuve P**, Marshall BC, Lowenfels AB. Risk of pancreatic cancer in patients with cystic fibrosis. *Gut* 2007; **56**: 1327-1328 [PMID: [17698876](#) DOI: [10.1136/gut.2007.125278](#)]
- 94 **Weinstein LS**. G(s)alpha mutations in fibrous dysplasia and McCune-Albright syndrome. *J Bone Miner Res* 2006; **21** Suppl 2: P120-P124 [PMID: [17229000](#) DOI: [10.1359/jbmr.06s223](#)]
- 95 **Majoer BC**, Boyce AM, Bovée JV, Smit VT, Collins MT, Cleton-Jansen AM, Dekkers OM, Hamdy NA, Dijkstra PS, Appelman-Dijkstra NM. Increased Risk of Breast Cancer at a Young Age in Women with Fibrous Dysplasia. *J Bone Miner Res* 2018; **33**: 84-90 [PMID: [28856726](#) DOI: [10.1002/jbmr.3286](#)]
- 96 **Vortmeyer AO**, Gläsker S, Mehta GU, Abu-Asab MS, Smith JH, Zhuang Z, Collins MT, Oldfield EH. Somatic GNAS mutation causes widespread and diffuse pituitary disease in acromegalic patients with McCune-Albright syndrome. *J Clin Endocrinol Metab* 2012; **97**: 2404-2413 [PMID: [22564667](#) DOI: [10.1210/jc.2012-1274](#)]
- 97 **Collins MT**, Sarlis NJ, Merino MJ, Monroe J, Crawford SE, Krakoff JA, Guthrie LC, Bonat S, Robey PG, Shenker A. Thyroid carcinoma in the McCune-Albright syndrome: contributory role of activating Gs alpha mutations. *J Clin Endocrinol Metab* 2003; **88**: 4413-4417 [PMID: [12970318](#) DOI: [10.1210/jc.2002-021642](#)]
- 98 **Nishikawa G**, Sekine S, Ogawa R, Matsubara A, Mori T, Taniguchi H, Kushima R, Hiraoka N,

- Tsuta K, Tsuda H, Kanai Y. Frequent GNAS mutations in low-grade appendiceal mucinous neoplasms. *Br J Cancer* 2013; **108**: 951-958 [PMID: [23403822](#) DOI: [10.1038/bjc.2013.47](#)]
- 99 **Fragoso MC**, Latronico AC, Carvalho FM, Zerbini MC, Marcondes JA, Araujo LM, Lando VS, Frazzatto ET, Mendonca BB, Villares SM. Activating mutation of the stimulatory G protein (gsp) as a putative cause of ovarian and testicular human stromal Leydig cell tumors. *J Clin Endocrinol Metab* 1998; **83**: 2074-2078 [PMID: [9626141](#) DOI: [10.1210/jcem.83.6.4847](#)]
- 100 **Robinson C**, Estrada A, Zaheer A, Singh VK, Wolfgang CL, Goggins MG, Hruban RH, Wood LD, Noë M, Montgomery EA, Guthrie LC, Lennon AM, Boyce AM, Collins MT. Clinical and Radiographic Gastrointestinal Abnormalities in McCune-Albright Syndrome. *J Clin Endocrinol Metab* 2018; **103**: 4293-4303 [PMID: [30124968](#) DOI: [10.1210/jc.2018-01022](#)]
- 101 **Parvanescu A**, Cros J, Ronot M, Hentic O, Grybek V, Couvelard A, Levy P, Chanson P, Ruszniewski P, Sauvanet A, Gaujoux S. Lessons from McCune-Albright syndrome-associated intraductal papillary mucinous neoplasms: : GNAS-activating mutations in pancreatic carcinogenesis. *JAMA Surg* 2014; **149**: 858-862 [PMID: [24898823](#) DOI: [10.1001/jamasurg.2014.535](#)]
- 102 **Tanaka M**. Controversies in the management of pancreatic IPMN. *Nat Rev Gastroenterol Hepatol* 2011; **8**: 56-60 [PMID: [21212775](#) DOI: [10.1038/nrgastro.2010.193](#)]
- 103 **Pergolini I**, Sahara K, Ferrone CR, Morales-Oyarvide V, Wolpin BM, Mucci LA, Brugge WR, Mino-Kenudson M, Patino M, Sahani DV, Warshaw AL, Lillemoe KD, Fernández-Del Castillo C. Long-term Risk of Pancreatic Malignancy in Patients With Branch Duct Intraductal Papillary Mucinous Neoplasm in a Referral Center. *Gastroenterology* 2017; **153**: 1284-1294. e1 [PMID: [28739282](#) DOI: [10.1053/j.gastro.2017.07.019](#)]
- 104 **Alhakim M**, Elshimy G, Vinales K, Correa R. SUN-324 Should We Screen for Pancreatic Cancer in Patient with Mccune Albright Syndrome? *J Endocr Soc* 2019; **3**: SUN-324 [DOI: [10.1210/js.2019-SUN-324](#)]
- 105 **Kamilaris CDC**, Stratakis CA. Multiple Endocrine Neoplasia Type 1 (MEN1): An Update and the Significance of Early Genetic and Clinical Diagnosis. *Front Endocrinol (Lausanne)* 2019; **10**: 339 [PMID: [31263451](#) DOI: [10.3389/fendo.2019.00339](#)]
- 106 **Jensen RT**, Norton JA. Treatment of Pancreatic Neuroendocrine Tumors in Multiple Endocrine Neoplasia Type 1: Some Clarity But Continued Controversy. *Pancreas* 2017; **46**: 589-594 [PMID: [28426491](#) DOI: [10.1097/MPA.0000000000000825](#)]
- 107 **Concolino P**, Costella A, Capoluongo E. Multiple endocrine neoplasia type 1 (MEN1): An update of 208 new germline variants reported in the last nine years. *Cancer Genet* 2016; **209**: 36-41 [PMID: [26767918](#) DOI: [10.1016/j.cancergen.2015.12.002](#)]
- 108 **Hannan FM**, Nesbit MA, Christie PT, Fratter C, Dudley NE, Sadler GP, Thakker RV. Familial isolated primary hyperparathyroidism caused by mutations of the MEN1 gene. *Nat Clin Pract Endocrinol Metab* 2008; **4**: 53-58 [PMID: [18084346](#) DOI: [10.1038/ncpendmet0718](#)]
- 109 **Scarpa A**, Chang DK, Nones K, Corbo V, Patch AM, Bailey P, Lawlor RT, Johns AL, Miller DK, Mafficini A, Rusev B, Scardoni M, Antonello D, Barbi S, Sikora KO, Cingarlini S, Vicentini C, McKay S, Quinn MC, Bruxner TJ, Christ AN, Harliwong I, Idrisoglu S, McLean S, Nourse C, Nourbakhsh E, Wilson PJ, Anderson MJ, Fink JL, Newell F, Waddell N, Holmes O, Kazakoff SH, Leonard C, Wood S, Xu Q, Nagaraj SH, Amato E, Dalai I, Bersani S, Cataldo I, Dei Tos AP, Capelli P, Davi MV, Landoni L, Malpaga A, Miotto M, Whitehall VL, Leggett BA, Harris JL, Harris J, Jones MD, Humphris J, Chantrill LA, Chin V, Nagrial AM, Pajic M, Scarlett CJ, Pinho A, Rooman I, Toon C, Wu J, Pinese M, Cowley M, Barbour A, Mawson A, Humphrey ES, Colvin EK, Chou A, Lovell JA, Jamieson NB, Duthie F, Gingras MC, Fisher WE, Dagg RA, Lau LM, Lee M, Pickett HA, Reddel RR, Samra JS, Kench JG, Merrett ND, Epari K, Nguyen NQ, Zeps N, Falconi M, Simbolo M, Butturini G, Van Buren G, Partelli S, Fassan M; Australian Pancreatic Cancer Genome Initiative, Khanna KK, Gill AJ, Wheeler DA, Gibbs RA, Musgrove EA, Bassi C, Tortora G, Pederzoli P, Pearson JV, Biankin AV, Grimmond SM. Whole-genome landscape of pancreatic neuroendocrine tumours. *Nature* 2017; **543**: 65-71 [PMID: [28199314](#) DOI: [10.1038/nature21063](#)]
- 110 **Thakker RV**, Newey PJ, Walls GV, Bilezikian J, Dralle H, Ebeling PR, Melmed S, Sakurai A, Tonelli F, Brandi ML; Endocrine Society. Clinical practice guidelines for multiple endocrine neoplasia type 1 (MEN1). *J Clin Endocrinol Metab* 2012; **97**: 2990-3011 [PMID: [22723327](#) DOI: [10.1210/jc.2012-1230](#)]
- 111 **Findeis-Hosey JJ**, McMahon KQ, Findeis SK. Von Hippel-Lindau Disease. *J Pediatr Genet* 2016; **5**: 116-123 [PMID: [27617152](#) DOI: [10.1055/s-0036-1579757](#)]
- 112 **Maher ER**, Neumann HP, Richard S. von Hippel-Lindau disease: a clinical and scientific review. *Eur J Hum Genet* 2011; **19**: 617-623 [PMID: [21386872](#) DOI: [10.1038/ejhg.2010.175](#)]
- 113 **Lubensky IA**, Pack S, Ault D, Vortmeyer AO, Libutti SK, Choyke PL, Walther MM, Linehan WM, Zhuang Z. Multiple neuroendocrine tumors of the pancreas in von Hippel-Lindau disease patients: histopathological and molecular genetic analysis. *Am J Pathol* 1998; **153**: 223-231 [PMID: [9665483](#) DOI: [10.1016/S0002-9440\(10\)65563-0](#)]
- 114 **Blansfield JA**, Choyke L, Morita SY, Choyke PL, Pingpank JF, Alexander HR, Seidel G, Shutack Y, Yuldasheva N, Eugeni M, Bartlett DL, Glenn GM, Middleton L, Linehan WM, Libutti SK. Clinical, genetic and radiographic analysis of 108 patients with von Hippel-Lindau disease (VHL) manifested by pancreatic neuroendocrine neoplasms (PNETs). *Surgery* 2007; **142**: 814-818; discussion 818 [PMID: [18063061](#) DOI: [10.1016/j.surg.2007.09.012](#)]
- 115 **Tamura K**, Nishimori I, Ito T, Yamasaki I, Igarashi H, Shuin T. Diagnosis and management of

- pancreatic neuroendocrine tumor in von Hippel-Lindau disease. *World J Gastroenterol* 2010; **16**: 4515-4518 [PMID: [20857520](#) DOI: [10.3748/wjg.v16.i36.4515](#)]
- 116 **Fang S**, Krahe R, Lozano G, Han Y, Chen W, Post SM, Zhang B, Wilson CD, Bachinski LL, Strong LC, Amos CI. Effects of MDM2, MDM4 and TP53 codon 72 polymorphisms on cancer risk in a cohort study of carriers of TP53 germline mutations. *PLoS One* 2010; **5**: e10813 [PMID: [20520810](#) DOI: [10.1371/journal.pone.0010813](#)]
 - 117 **Hwang SJ**, Lozano G, Amos CI, Strong LC. Germline p53 mutations in a cohort with childhood sarcoma: sex differences in cancer risk. *Am J Hum Genet* 2003; **72**: 975-983 [PMID: [12610779](#) DOI: [10.1086/374567](#)]
 - 118 **Gonzalez KD**, Noltner KA, Buzin CH, Gu D, Wen-Fong CY, Nguyen VQ, Han JH, Lowstuter K, Longmate J, Sommer SS, Weitzel JN. Beyond Li Fraumeni Syndrome: clinical characteristics of families with p53 germline mutations. *J Clin Oncol* 2009; **27**: 1250-1256 [PMID: [19204208](#) DOI: [10.1200/JCO.2008.16.6959](#)]
 - 119 **Mai PL**, Malkin D, Garber JE, Schiffman JD, Weitzel JN, Strong LC, Wyss O, Locke L, Means V, Achatz MI, Hainaut P, Frebourg T, Evans DG, Bleiker E, Patenaude A, Schneider K, Wilfond B, Peters JA, Hwang PM, Ford J, Tabori U, Ognjanovic S, Dennis PA, Wentzensen IM, Greene MH, Fraumeni JF Jr, Savage SA. Li-Fraumeni syndrome: report of a clinical research workshop and creation of a research consortium. *Cancer Genet* 2012; **205**: 479-487 [PMID: [22939227](#) DOI: [10.1016/j.cancergen.2012.06.008](#)]
 - 120 **Birch JM**, Hartley AL, Tricker KJ, Prosser J, Condie A, Kelsey AM, Harris M, Jones PH, Binchy A, Crowther D. Prevalence and diversity of constitutional mutations in the p53 gene among 21 Li-Fraumeni families. *Cancer Res* 1994; **54**: 1298-1304 [PMID: [8118819](#)]
 - 121 **Solomon S**, Das S, Brand R, Whitcomb DC. Inherited pancreatic cancer syndromes. *Cancer J* 2012; **18**: 485-491 [PMID: [23187834](#) DOI: [10.1097/PPO.0b013e318278c4a6](#)]
 - 122 **Brand RE**, Lerch MM, Rubinstein WS, Neoptolemos JP, Whitcomb DC, Hruban RH, Brentnall TA, Lynch HT, Canto MI; Participants of the Fourth International Symposium of Inherited Diseases of the Pancreas. Advances in counselling and surveillance of patients at risk for pancreatic cancer. *Gut* 2007; **56**: 1460-1469 [PMID: [17872573](#) DOI: [10.1136/gut.2006.108456](#)]
 - 123 **Brose MS**, Rebbeck TR, Calzone KA, Stopfer JE, Nathanson KL, Weber BL. Cancer risk estimates for BRCA1 mutation carriers identified in a risk evaluation program. *J Natl Cancer Inst* 2002; **94**: 1365-1372 [PMID: [12237282](#) DOI: [10.1093/jnci/94.18.1365](#)]
 - 124 **Kanda M**, Matthaei H, Wu J, Hong SM, Yu J, Borges M, Hruban RH, Maitra A, Kinzler K, Vogelstein B, Goggins M. Presence of somatic mutations in most early-stage pancreatic intraepithelial neoplasia. *Gastroenterology* 2012; **142**: 730-733. e9 [PMID: [22226782](#) DOI: [10.1053/j.gastro.2011.12.042](#)]
 - 125 **Brune KA**, Lau B, Palmisano E, Canto M, Goggins MG, Hruban RH, Klein AP. Importance of age of onset in pancreatic cancer kindreds. *J Natl Cancer Inst* 2010; **102**: 119-126 [PMID: [20068195](#) DOI: [10.1093/jnci/djp466](#)]
 - 126 **Kastrinos F**, Mukherjee B, Tayob N, Wang F, Sparr J, Raymond VM, Bandipalliam P, Stoffel EM, Gruber SB, Syngal S. Risk of pancreatic cancer in families with Lynch syndrome. *JAMA* 2009; **302**: 1790-1795 [PMID: [19861671](#) DOI: [10.1001/jama.2009.1529](#)]
 - 127 **Geary J**, Sasieni P, Houlston R, Izatt L, Eeles R, Payne SJ, Fisher S, Hodgson SV. Gene-related cancer spectrum in families with hereditary non-polyposis colorectal cancer (HNPCC). *Fam Cancer* 2008; **7**: 163-172 [PMID: [17939062](#) DOI: [10.1007/s10689-007-9164-6](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

