World Journal of *Gastroenterology*

World J Gastroenterol 2021 June 28; 27(24): 3429-3692





Published by Baishideng Publishing Group Inc

W J G World Journal of Gastroenterology

Contents

Weekly Volume 27 Number 24 June 28, 2021

OPINION REVIEW

3429 Trial eligibility in advanced hepatocellular carcinoma: Does it support clinical practice in underrepresented subgroups?

Piñero F, da Fonseca LG

REVIEW

- 3440 Chronic intestinal failure and short bowel syndrome in Crohn's disease Aksan A, Farrag K, Blumenstein I, Schröder O, Dignass AU, Stein J
- 3466 Non-cirrhotic hepatocellular carcinoma in chronic viral hepatitis: Current insights and advancements Perisetti A, Goyal H, Yendala R, Thandassery RB, Giorgakis E
- 3483 Natural killer cells in pancreatic cancer stroma Fincham REA, Delvecchio FR, Goulart MR, Yeong JPS, Kocher HM
- 3502 COVID-19 and its effects on the digestive system Cao TT, Zhang GQ, Pellegrini E, Zhao Q, Li J, Luo LJ, Pan HQ
- 3516 Neuropilin-1: A feasible link between liver pathologies and COVID-19 Benedicto A, García-Kamiruaga I, Arteta B

MINIREVIEWS

- 3530 Hepatitis delta virus: From infection to new therapeutic strategies Niro GA, Ferro A, Cicerchia F, Brascugli I, Durazzo M
- 3543 Usefulness of artificial intelligence in gastric neoplasms Kim JH. Nam SJ. Park SC
- 3556 Current impact of viral hepatitis on liver cancer development: The challenge remains de Mattos ÂZ, Debes JD, Boonstra A, Yang JD, Balderramo DC, Sartori GDP, de Mattos AA
- 3568 Room for improvement in the treatment of pancreatic cancer: Novel opportunities from gene targeted therapy

Galanopoulos M, Doukatas A, Gkeros F, Viazis N, Liatsos C

ORIGINAL ARTICLE

Basic Study

3581 Fasudil prevents liver fibrosis via activating natural killer cells and suppressing hepatic stellate cells Han QJ, Mu YL, Zhao HJ, Zhao RR, Guo QJ, Su YH, Zhang J



O t	World Journal of Gastroenterology		
Conten	Weekly Volume 27 Number 24 June 28, 2021		
3595	Early genetic diagnosis of clarithromycin resistance in Helicobacter pylori		
	Li XH, Huang YY, Lu LM, Zhao LJ, Luo XK, Li RJ, Dai YY, Qin C, Huang YQ, Chen H		
3609	Case Control Study		
	Altered profiles of fecal bile acids correlate with gut microbiota and inflammatory responses in patients with ulcerative colitis		
	Yang ZH, Liu F, Zhu XR, Suo FY, Jia ZJ, Yao SK		
3630	Retrospective Study		
	High rate of complete histopathological response in hepatocellular carcinoma patients after combined transarterial chemoembolization and stereotactic body radiation therapy		
	Bauer U, Gerum S, Roeder F, Münch S, Combs SE, Philipp AB, De Toni EN, Kirstein MM, Vogel A, Mogler C, Haller B, Neumann J, Braren RF, Makowski MR, Paprottka P, Guba M, Geisler F, Schmid RM, Umgelter A, Ehmer U		
3643	Stem cell injection for complex anal fistula in Crohn's disease: A single-center experience		
	Schwandner O		
3654	Observational Study		
	Laparoscopic lateral lymph node dissection in two fascial spaces for locally advanced lower rectal cancer		
	Jiang HH, Liu HL, Li AJ, Wang WC, Lv L, Peng J, Pan ZH, Chang Y, Lin MB		
	SYSTEMATIC REVIEWS		
3668	Disorders of the brain-gut interaction and eating disorders		
	Stanculete MF, Chiarioni G, Dumitrascu DL, Dumitrascu DI, Popa SL		
3682	Weight loss interventions in living donor liver transplantation as a tool in expanding the donor pool: A systematic review and meta-analysis		

Trakroo S, Bhardwaj N, Garg R, Modaresi Esfeh J



Contents

Weekly Volume 27 Number 24 June 28, 2021

ABOUT COVER

Editorial Board Member of World Journal of Gastroenterology, Yasemin H Balaban, MD, Professor, Department of Gastroenterology, Faculty of Medicine, Hacettepe University, Ankara 06100, Turkey. ybalaban@hacettepe.edu.tr

AIMS AND SCOPE

The primary aim of World Journal of Gastroenterology (WJG, World J Gastroenterol) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2020 edition of Journal Citation Report[®] cites the 2019 impact factor (IF) for WJG as 3.665; IF without journal self cites: 3.534; 5-year IF: 4.048; Ranking: 35 among 88 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2019 is 7.1 and Scopus CiteScore rank 2019: Gastroenterology is 17/137.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Li-Li Wang, Production Department Director: Yu-Jie Ma; Editorial Office Director: Ze-Mao Gong.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Gastroenterology	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1007-9327 (print) ISSN 2219-2840 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
October 1, 1995	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Weekly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Andrzej S Tarnawski, Subrata Ghosh	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
http://www.wjgnet.com/1007-9327/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE June 28, 2021	STEPS FOR SUBMITTING MANUSCRIPTS https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2021 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2021 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



WJG

World Journal of Gastroenterology

Submit a Manuscript: https://www.f6publishing.com

World J Gastroenterol 2021 June 28; 27(24): 3556-3567

DOI: 10.3748/wjg.v27.i24.3556

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

MINIREVIEWS

Current impact of viral hepatitis on liver cancer development: The challenge remains

Ângelo Zambam de Mattos, Jose D Debes, Andre Boonstra, Ju-Dong Yang, Domingo C Balderramo, Giovana D P Sartori, Angelo Alves de Mattos

ORCID number: Ângelo Zambam de Mattos 0000-0002-3063-0199; Jose D Debes 0000-0002-1512-2604: Andre Boonstra 0000-0001-8607-1616; Ju-Dong Yang 0000-0001-7834-9825; Domingo C Balderramo 0000-0001-9598-2577; Giovana D P Sartori 0000-0002-2701-3499; Angelo Alves de Mattos 0000-0003-2417-9765.

Author contributions: All authors contributed to this paper with conception, literature review and analysis, drafting and critical revision of the manuscript, and approval of the final version of the paper.

Supported by European-Latin American ESCALON Consortium, funded by the EU Horizon 2020 Program, No. 825510; and Robert Wood Johnson Foundation, Harold Amos Medical Faculty Development Program (to Debes JD).

Conflict-of-interest statement: There are no conflicts of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to

Ângelo Zambam de Mattos, Angelo Alves de Mattos, Department of Gastroenterology and Hepatology, Federal University of Health Sciences of Porto Alegre, Porto Alegre 90020-090, Brazil

Ângelo Zambam de Mattos, Angelo Alves de Mattos, Gastroenterology and Hepatology Unit, Irmandade Santa Casa de Misericórdia de Porto Alegre, Porto Alegre 90050-170, Brazil

Jose D Debes, Department of Medicine, Division of Gastroenterology and Infectious Diseases, University of Minnesota, Minneapolis, MN 55455, United States

Jose D Debes, Andre Boonstra, Department of Gastroenterology and Hepatology, Erasmus Medical Center, Rotterdam NL-3015, Netherlands

Ju-Dong Yang, Division of Digestive and Liver Diseases, Department of Medicine, Cedars-Sinai Medical Center, Los Angeles, CA 90048, United States

Domingo C Balderramo, Department of Gastroenterology, Hospital Privado Universitario de Có rdoba, Córdoba 5016, Argentina

Domingo C Balderramo, Department of Medicine, Instituto Universitario de Ciencias Biomé dicas de Córdoba, Córdoba 5016, Argentina

Giovana D P Sartori, Department of Internal Medicine, Hospital Nossa Senhora da Conceição, Porto Alegre 91350-200, Brazil

Corresponding author: Angelo Zambam de Mattos, MD, MSc, PhD, Professor, Department of Gastroenterology and Hepatology, Federal University of Health Sciences of Porto Alegre, 154, Professor Annes Dias St., Office 1103, Porto Alegre 90020-090, Brazil. angmattos@hotmail.com

Abstract

Chronic infections due to hepatitis B and hepatitis C viruses are responsible for most cases of hepatocellular carcinoma (HCC) worldwide, and this association is likely to remain during the next decade. Moreover, viral hepatitis-related HCC imposes an important burden on public health in terms of disability-adjusted life years. In order to reduce such a burden, some major challenges must be faced. Universal vaccination against hepatitis B virus, especially in the neonatal period, is probably the most relevant primary preventive measure against the develo-



distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: htt p://creativecommons.org/License s/by-nc/4.0/

Manuscript source: Invited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: Brazil

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): 0 Grade D (Fair): D Grade E (Poor): 0

Received: January 29, 2021 Peer-review started: January 29, 2021 First decision: February 25, 2021 Revised: March 11, 2021 Accepted: April 21, 2021 Article in press: April 21, 2021 Published online: June 28, 2021

P-Reviewer: Zheng SJ S-Editor: Gao CC L-Editor: Webster JR P-Editor: Wang LL



pment of HCC. Moreover, considering the large adult population already infected with hepatitis B and C viruses, it is also imperative to identify these individuals to ensure their access to treatment. Both hepatitis B and C currently have highly effective therapies, which are able to diminish the risk of development of liver cancer. Finally, it is essential for individuals at high-risk of HCC to be included in surveillance programs, so that tumors are detected at an early stage. Patients with hepatitis B or C and advanced liver fibrosis or cirrhosis benefit from being followed in a surveillance program. As hepatitis B virus is oncogenic and capable of leading to liver cancer even in individuals with early stages of liver fibrosis, other high-risk groups of patients with hepatitis B are also candidates for surveillance. Considerable effort is required concerning these strategies in order to decrease the incidence and the mortality of viral hepatitis-related HCC.

Key Words: Hepatitis B virus; Hepatitis C virus; Hepatocellular carcinoma; Epidemiology; Vaccination; Surveillance

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Hepatitis B and C are associated with most cases of hepatocellular carcinoma, and it is estimated that this scenario will remain for the next decade. This review highlights the impact of viral hepatitis on the development of liver cancer, the characteristics of viral hepatitis-related hepatocellular carcinoma, and the challenges that must be faced in order to reduce their burden.

Citation: de Mattos ÂZ, Debes JD, Boonstra A, Yang JD, Balderramo DC, Sartori GDP, de Mattos AA. Current impact of viral hepatitis on liver cancer development: The challenge remains. World J Gastroenterol 2021; 27(24): 3556-3567

URL: https://www.wjgnet.com/1007-9327/full/v27/i24/3556.htm **DOI:** https://dx.doi.org/10.3748/wjg.v27.i24.3556

INTRODUCTION

Viral infections are the leading cause of hepatitis, with hepatitis B virus (HBV) and hepatitis C virus (HCV) being the most important causes of chronic viral hepatitis worldwide. It was estimated that nearly 292 million individuals had chronic HBV infection in 2016 (prevalence of 3.9%) and 71 million people had chronic HCV infection in 2015 (prevalence of 1%)[1-3]. Hepatitis D virus (HDV) is a defective virus, requiring HBV coexistence in order to replicate. It was estimated that HDV caused chronic infection in approximately 12 million individuals in 2016[4]. Most HBV, HCV and HDV infections are concentrated in low- and lower middle-income countries[3].

The incidence of liver cancer is the sixth highest of all malignant neoplasms (905677 new cases in 2020, an incidence of 11.6/100000 inhabitants). Moreover, it is the second cause of cancer-related death in the world (830180 deaths in 2020, mortality of 10.7/100000 inhabitants)[5]. Hepatocellular carcinoma (HCC) accounts for approximately 75%-85% of primary liver cancers[6,7]. HCC frequently develops in patients with cirrhosis, with an annual incidence of 1%-4%[8]. HBV and HCV are currently the two most important risk factors for HCC development worldwide. This is reflected by the geographical distribution of liver cancer, which coincides with those of HBV and HCV, predominating in transitioning countries, particularly in Asia and Africa[6,7,9]. In a systematic review of 260 studies involving 119006 patients with HCC, the seroprevalence of HBV and/or HCV was over 60% in most of the 50 countries which contributed with data[10]. Furthermore, it is noteworthy that, according to the Global Burden of Disease Study, HBV-related liver cancer was associated with 5.80 million disability-adjusted life years, and HCV-related liver cancer, with 2.88 million disability-adjusted life years in 2019[11]. The impact of HBV may be further increased by the fact that HCC usually develops at an earlier age in patients infected with this virus[12].



Despite vaccination against HBV and effective treatments for HBV and HCV, it is estimated that their impact on liver health will remain for the next decade at least. Using data from 195 countries or territories between 1990 and 2017, a recent study proposed Bayesian models to project primary liver cancer incidence through 2030. The authors estimated that the age-standardized incidence rate of liver cancer will increase globally from 11.80 in 2017 to 14.08 per 100000 inhabitants in 2030. According to the etiology of liver disease, HBV was responsible for 46.5% of cases of liver cancer in 1990 and 42.4% in 2017, and it is estimated that it will be associated with 40.7% of cases in 2030. With regard to HCV, it was associated with 25.2% of cases of primary liver cancer in 1990 and 27.0% in 2017, and it is projected that it will be responsible for 26.8% of cases in 2030[13]. The aim of this article is to review the current impact of viral hepatitis on the development of liver cancer, the characteristics of viral hepatitisrelated HCC and the challenges to reduce their burden (Figure 1).

IMMUNE ASPECTS OF VIRAL HEPATITIS IN HEPATOCARCINOGENESIS

In chronic viral hepatitis, the development of fibrosis, cirrhosis and HCC is linked to the activity of the immune response in the infected liver. Both HBV and HCV uniquely infect hepatocytes and replicate in a non-cytopathic manner. The ensuing liver damage that is observed during the acute and the chronic phase of infection is the consequence of virus-specific as well as non-specific immune activity within the inflamed liver. Histologically, active phases of chronic HBV and HCV infections are characterized by extensive infiltration around the portal tract areas consisting predominantly of CD4+ and CD8+ T cells, but these cells are incapable of clearing these viruses. Multiple mechanisms have been described that explain the weak activity of CD4+ and CD8+ T cells, and that are the main reasons for persistence of HBV and HCV in the liver, and the slowly progressing liver disease[14-16]. The mechanisms include impairment of dendritic cells and natural killer (NK) cells, increased production of immunosuppressive cytokines and an increase in the numbers of regulatory T cells. In addition, the continuous exposure of immune cells to high levels of viral antigens for many years further contributes and maintains the persistent infection due to exhaustion of HBV and HCV-specific T cells, which renders these cells functionally impaired mediated via triggering of exhaustion markers, such as programmed cell death protein 1 (PD-1), cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) and others[17].

In the liver, the balance between immune tolerance and immune activation is normally tightly controlled to ensure efficient elimination of pathogen products and transformed cells. Dysregulation of this balance due to viral infection has severe consequences for the process of immune surveillance that is highly efficient in the detection and elimination of transformed cells, and contributes to HCC development [18]. Also, alterations in lipid metabolism and the release of reactive oxygen species contribute to tumor initiation[19].

Interestingly, the status of tumor-infiltrating immune cells in HCC has a high degree of resemblance to that observed in an HBV or HCV inflamed liver: an increased T cell infiltrate is observed in HCC, but the CD8+ T cells are dysfunctional with low production of the cytolytic enzymes granzyme and perforin, low overall cytokine production, and relatively high expression of the exhaustion markers PD-1 and CTLA-4[20,21]. In addition, increased numbers of regulatory T cells and reduced numbers and functionality of NK cells have been reported by many groups[18,22,23]. These immune dysfunctions are generally more pronounced in the advanced HCC stages.

The importance of the immune system in controlling the development and progression to advanced stages of HCC is highlighted by observations that increased T cell infiltrates are associated with improved overall survival in HCC and lower tumor recurrence following resection[24-26]. Furthermore, the enrichment of exhausted T cells in HCC is associated with poorer survival, while higher numbers of infiltrating NK cells are associated with better survival. Using cytometry by time of flight and RNA sequencing, a recent study defined different immune subsets and showed that the tumor microenvironment in HBV-related HCC consisted of a more immunosuppressive and exhausted phenotype with an enrichment of regulatory T cells and resident memory T cells that differs from the non-viral-related HCC environment. The study further demonstrated that regulatory T cells are associated with poor prognosis, and memory T cells, with good prognosis[27].

The crucial importance of the patient's immune activity against the tumor has initiated the development of numerous immunotherapeutic approaches in recent years, all aimed at boosting anti-tumor immunity^[28]. The approaches are diverse and





Figure 1 Major challenges to reduce the burden of viral hepatitis-related hepatocellular carcinoma. HCC: Hepatocellular carcinoma; HBV: Hepatitis B virus; HCV: Hepatitis C virus.

> include strategies using peptide vaccines and cell-based therapies. Moreover, therapies aimed at neutralizing exhaustion markers using immune checkpoint inhibitors have shown promising results and PD-1 blockade has been approved as therapy for HCC. However, many studies are ongoing to improve therapy success by examining the therapeutic outcomes using different combinations of checkpoint inhibitors.

HEPATITIS B AND HCC

Implications on HCC

Advances in next-generation sequencing technology and integrative studies combining multiomic approaches, encompassing genomic and epigenomic characterization and transcriptome, proteomic, and metabolomic analysis of HCCs have shown that HCC is heterogeneous at the histomolecular level, with variable molecular features and clinical outcomes[29-31]. While ongoing hepatic inflammation, increased cell turnover, repeated cycles of cell death and regeneration, DNA instability leading to an alteration in DNA and dysregulation of CpG island methylation are common pathways leading to HCC regardless of the underlying etiology of liver disease[32], there are HBV specific pathogenesis pathways in HCC development. HBV promotes HCC through viral DNA integration into host genes that can provide a growth advantage to the host cell. HBV has been found to be integrated into or near cancerrelated genes[33,34]. For example, promoter mutation of telomerase reverse transcriptase, which is often integrated by HBV DNA, is the most frequent genetic event in HBV-associated HCC. The translated protein products of the viral HBV genome have also been implicated in promoting carcinogenesis. For instance, HBx protein, a regulatory protein required for viral replication, has been shown to regulate hepatocyte proliferation, regeneration and apoptosis by modulating the PI3K/AktmTOR, STAT3/Nanog, and Wnt/ β -catenin signaling pathways to promote the progression of HCC[35-37].

Occult HBV infection

Occult HBV infection is a condition where HBV DNA replication occurs in the liver in patients with negative serum HBsAg[38]. While detection of anti-HBc (hepatitis B core antibody) in the blood indicates prior HBV infection, 20% of patients with occult HBV infection are seronegative for anti-HBc. HBV DNA analysis should be further pursued to confirm the diagnosis[39]. Occult HBV infection increases the risk of HCC development in patients with or without chronic liver disease, which is thought to be mediated by HBV DNA integration into the host genome, the production of prooncogenic proteins such as HBx protein, and persistent low-grade necroinflammation leading to fibrosis progression[39]. Currently, HCC surveillance is not routinely

recommended in patients with occult HBV infection in the absence of advanced fibrosis/cirrhosis.

HCC surveillance in hepatitis B

HCC continues to be one of the leading causes of cancer-related death throughout the world and it frequently develops within the context of cirrhosis[40]. This has also led to the development of guidelines by all the major liver societies for HCC surveillance in patients with cirrhosis given its potential for early-stage cancer detection, which likely links to curative treatment[41,42]. Given its high oncogenic potential and its tendency to develop HCC in the absence of cirrhosis, screening for HCC is indicated in a broader population of HBV patients. All HBV patients with HBV cirrhosis (except for patients with severe liver dysfunction and not eligible for liver transplantation) are recommended to undergo HCC surveillance. Surveillance is also recommended in a selected high-risk population of non-cirrhotic HBV patients, as demonstrated in Table 1[43].

Eradication of hepatitis B through vaccination

HBV vaccination is an effective primary prevention method for HCC. The effectiveness of HBV vaccination in reducing HCC incidence was reported in a nationwide population-based study from Taiwan. The study used data from 2 Taiwanese HCC registry systems and showed that HCC incidence was four-fold higher in the HBV unvaccinated cohort than in the vaccinated at birth cohorts. Among the 1509 patients who were 6-26 years old diagnosed with HCC from 1983 through 2011, 1343 were born before, and 166 were born after the HBV vaccination program began. Moreover, the relative risks for HCC in patients vaccinated at 6-9 years old, 10-14 years old, 15-19 years old, and 20-26 years old in comparison to those who were not vaccinated were 0.26 [95% confidence interval (CI): 0.17-0.40], 0.34 (95%CI: 0.25-0.48), 0.37 (95%CI: 0.25-0.51), and 0.42 (95% CI: 0.32-0.56), respectively [44]. Although neonatal HBV vaccination is recommended in most countries, vaccine coverage is still less than 70% in most African countries where the incidence rates of HCC remain high[45], providing a clear window of opportunity for improvement in primary prevention of HCC[40].

Major challenges

Aside from identifying infected patients, offering effective treatment against HBV and surveying those at high risk for HCC, the major challenge in order to reduce HBVrelated HCC burden is widening the vaccination coverage against the virus. Despite the increase in vaccination coverage between 2010 and 2018 in 97% of the 194 member states of the World Health Organization (WHO), with an increment from 73% to 84% in the completion of the three-dose vaccination schedule and with an increase from 28% to 42% in the neonatal vaccination coverage in the 108 countries in which newborns are vaccinated, improvements are still necessary[46]. For instance, vaccination coverage requires improvement in countries such as Nigeria, India and Indonesia, which together are responsible for 50% of HBV infections in children under 5 years of age[3].

HEPATITIS C AND HCC

Implications on HCC

HCV is one of the most important risk factors for HCC. As previously mentioned, HCV is a non-cytopathic virus, which leads to liver damage through immunemediated mechanisms. However, the virus itself also plays a role in the development of HCC, which becomes clear when considering, for instance, that HCV genotype 3 is associated with a higher risk of HCC than other genotypes[47]. The mechanism through which HCV genotype 3 is involved in hepatocarcinogenesis is not yet completely understood, but the down-regulation of the phosphatase and tensin homolog (*PTEN*) gene, which has a tumor suppressor role, might be implicated[48].

Direct-acting antiviral (DAA) therapy in HCV-infected patients has been shown to reduce all-cause mortality (including liver-related and unrelated causes), as well as the incidence of HCC in patients with cirrhosis[49]. However, in the subgroup of patients with advanced fibrosis or cirrhosis, the risk of developing HCC after achieving sustained virologic response (SVR) remains high, ranging from 0.3% to 1.8% per year [50]. For this reason, different guidelines recommend lifelong surveillance for patients with advanced fibrosis or cirrhosis even after SVR[51]. Interestingly, recent studies have attempted to differentiate subgroups of patients with cirrhosis at the highest risk



Table 1 Indications for surveillance of hepatocellular carcinoma in patients with hepatitis B without cirrhosis

ndications for surveillance	
Asian males > 40 yr	
Asian females > 50 yr	
African ancestry	
First degree relative with HCC	
HDV co-infection	

HCC: Hepatocellular carcinoma; HDV: Hepatitis D virus.

for HCC after SVR. In an Italian study, the combination of clinical predictors (male sex and diabetes), albumin, and genetics identified patients at high risk for HCC[52]. Furthermore, it is suggested that changes in liver stiffness measured by transient elastography at 1 and 3 years after the end of HCV treatment could identify patients at low risk for HCC[53].

Population screening for hepatitis C

In 2017, the WHO Global Hepatitis Report stated that 71 million people worldwide were infected by HCV[54]. About 3.5 million to 5.0 million of these individuals are children and adolescents, and 2.3 million are co-infected with the human immunodeficiency virus (HIV)[51,54]. HCV infection causes approximately 400000 deaths annually, mainly from cirrhosis-related complications and HCC[54]. Despite the current availability of DAA therapies with high rates of SVR, the prevalence of HCV infection has not changed. One of the reasons that explain this situation is that only 20% of individuals with HCV infection worldwide are aware of their diagnosis. Another reason is that access to DAA treatments continues to be limited mainly by costs in several countries[54].

The WHO Global Health Sector Strategy on Viral Hepatitis recommended that an effort should be made in order to increase the HCV diagnosis rate to 30% by 2020 and to 90% by 2030[55]. Overall, HCV screening consists in a universal, one-time, and opt-out screening strategy in adults 18 years of age and older with an upper age limit of 79 years. However, periodic screening tests should be offered to individuals with behaviors, conditions or circumstances associated with an increased risk of exposure to HCV[51].

Vertical transmission could be present in about 5% of deliveries from mothers with HCV and accounts for the majority of HCV infections in the pediatric population[56]. Universal prenatal screening for HCV facilitates better identification of at-risk infants who require HCV testing[57]. This evaluation results in better detection of HCV infection in the pediatric population and enables early therapeutic interventions.

Eradication of hepatitis C

DAA regimens have a high cure rate for HCV infection[51]. Furthermore, new formulations have simplified the duration and administration of the treatment. However, in 2015, only 7% of people diagnosed with HCV had started antiviral treatment with DAA[54]. According to the strategy suggested by the WHO, access to DAA should reach 80% of eligible people by 2030[55]. After this declaration, access to treatment has improved in different countries. An example is the strategy in Australia, where more than 80% of the HCV-infected population was diagnosed during the last two decades and where there is currently an unrestricted DAA access program that permits prescriptions by any registered medical practitioner. This allowed for the initiation of DAA treatment in Australia for approximately 70% of the total population with HCV-related cirrhosis between 2014 and 2017[58].

Simultaneously with screening and therapy access, different preventive measures must be carried out in people capable of transmitting HCV (*e.g.*, people who inject drugs) in order to avoid new infections or reinfections in those who have been previously treated successfully. Global coverage of harm reduction programs for people who inject drugs, including needle and syringe programs, is currently less than 10% [54].

Major challenges

Currently, the major challenge in order to reduce HCV-related HCC burden is identifying infected patients. The silent course of the early stages of this disease makes this challenge even harder to overcome. Therefore, efforts should be made in order to improve awareness of the population and health care workers regarding the underdiagnosis of HCV[54]. It is also of great importance to shorten the pathway leading from the diagnosis to the extremely effective treatments against HCV, which could even allow for subgroups of patients to be treated in primary care[59].

Yet another challenge relates to the coronavirus disease 2019 pandemic. It is estimated that the impact of the pandemic on the global efforts towards HCV eradication could lead to an excess of 44800 cases of HCC[60].

CO-INFECTIONS AND HCC

As previously mentioned, HBV and HCV infections are the most common risk factors for HCC worldwide[61]. Individuals living with HIV are a high-risk population for developing HCC, mostly as a consequence of HCV and HBV co-infection[62,63]. Indeed, HCC has become a major clinical problem in HIV. Studies from Europe suggest that the incidence of HCC has risen in HIV-infected patients over the last 20 years, and in Spain alone HCC is the second cause of death in HIV-HCV co-infected patients with cirrhosis[64]. Studies from both the United States and Europe suggest that HIV may hasten the evolution of HBV-related HCC, resulting in an earlier age of HCC presentation in HIV co-infected patients[65,66].

Although the implementation of new treatments for HCV with DAA agents is expected to decrease HCC incidence, new cases will continue to emerge in the nearmedium term[67,68]. Moreover, a recent study that reported an HCC risk increase of 1% every year in HIV-HCV co-infected individuals in the HEPAVIR cohort in Europe included in their cohort 61% of individuals with cured HCV, indicating that this population remains at risk for HCC[63].

In HIV-HBV co-infection, a recent study from the EuroSIDA cohort showed a stable incidence of HCC in non-cirrhotic individuals under HBV therapy (tenofovir specifically), while increasing rates of HCC were seen over time in those not on tenofovir. This suggests a potential threshold to not survey the former group for HCC. However, the same study did show a continued increase in HCC risk among those that had cirrhosis, indicating the importance of surveillance in this population[69].

Most of the data reported on HIV and HCC originate in resource-rich settings. These dynamics related to poor outcomes are likely to be increased in resource-limited settings, where most of HIV infections occur. A small African study of 60 patients found that those infected with HIV developed HCC at a younger age (32 years) compared to those without HIV (49 years)[70], and a more recent study from South Africa showed the age of HCC development to be lower in those co-infected with HBV and HIV (mean age of 36 years *vs* 46 years in HIV-uninfected). Interestingly, this study showed that females were more impacted by HCC when co-infected with HIV, and the age of HCC presentation was much lower in females with HIV (mean age of 36 years *vs* 50 years in HIV-uninfected)[71]. In South America and Asia overall, data are scarce on the interrelation between HCC and HIV infection. The specific mechanisms underlying younger HCC occurrence in co-infected individuals with viral hepatitis and HIV is unclear. However, a large Swiss study found a direct association between CD4+ T cells and the risk of developing HCC, suggesting that impairment of the immune system could be implicated[72].

Surveillance for HCC represents a major challenge in co-infected populations not only due to the complex nature of dual disease, but also because most guidelines are tailored towards mono-infected individuals at risk for HCC. In addition, recent studies show that ultrasound surveillance has a low performance for HCC in HBV or HCV individuals co-infected with HIV, with a suboptimal 43% rate of early-stage diagnosis, compared to 63%-71% found in studies on HIV-uninfected cohorts[73,74].

Triple co-infection of HIV, HBV and HDV is rather uncommon, ranging from 1% to 20% depending on the geographical area[75]. However, studies performed in small cohorts have shown that the main impact of HDV on HBV-HIV disease is an accelerated pace to cirrhosis and hepatic decompensation[76]. In this context, a Swiss HIV cohort reported a 9-fold increase in HCC in those triple co-infected with HIV-HBV-HDV compared to HDV-negative individuals[77].

Zaishidene® WJG | https://www.wjgnet.com

CONCLUSION

Despite the advances in prevention and treatment of viral hepatitis, it is clear that many challenges remain in order to reduce the burden of viral hepatitis-related HCC. Universal neonatal vaccination against HBV, as well as vaccination of at-risk adult populations, screening for HBV and HCV, and access to highly-effective treatments against both viruses are instrumental measures that must be pursued with the objective of diminishing their impact on global health and particularly on the development of HCC. Finally, effective surveillance for HCC must be offered to patients who already have advanced fibrosis or cirrhosis and to high-risk HBVinfected individuals, so that HCC is detected at earlier stages, allowing for curative treatments and longer survival.

REFERENCES

- Polaris Observatory HCV Collaborators. Global prevalence and genotype distribution of hepatitis 1 C virus infection in 2015: a modelling study. Lancet Gastroenterol Hepatol 2017; 2: 161-176 [PMID: 28404132 DOI: 10.1016/S2468-1253(16)30181-9]
- Polaris Observatory Collaborators. Global prevalence, treatment, and prevention of hepatitis B 2 virus infection in 2016: a modelling study. Lancet Gastroenterol Hepatol 2018; 3: 383-403 [PMID: 29599078 DOI: 10.1016/S2468-1253(18)30056-6]
- Razavi H. Global Epidemiology of Viral Hepatitis. Gastroenterol Clin North Am 2020; 49: 179-189 [PMID: 32389357 DOI: 10.1016/j.gtc.2020.01.001]
- Stockdale AJ, Kreuels B, Henrion MYR, Giorgi E, Kyomuhangi I, de Martel C, Hutin Y, Geretti 4 AM. The global prevalence of hepatitis D virus infection: Systematic review and meta-analysis. J Hepatol 2020; 73: 523-532 [PMID: 32335166 DOI: 10.1016/j.jhep.2020.04.008]
- World Health Organization. Global Cancer Observatory GLOBOCAN 2020. [cited 17 January 5 2021]. In: World Health Organization [Internet]. Available from: https://gco.iarc.fr/
- McGlynn KA, Petrick JL, El-Serag HB. Epidemiology of Hepatocellular Carcinoma. Hepatology 6 2021; 73 Suppl 1: 4-13 [PMID: 32319693 DOI: 10.1002/hep.31288]
- Arnold M, Abnet CC, Neale RE, Vignat J, Giovannucci EL, McGlynn KA, Bray F. Global Burden of 5 Major Types of Gastrointestinal Cancer. Gastroenterology 2020; 159: 335-349. e15 [PMID: 32247694 DOI: 10.1053/j.gastro.2020.02.068]
- 8 Moon AM, Singal AG, Tapper EB. Contemporary Epidemiology of Chronic Liver Disease and Cirrhosis. Clin Gastroenterol Hepatol 2020; 18: 2650-2666 [PMID: 31401364 DOI: 10.1016/j.cgh.2019.07.060]
- 9 Debes JD, Chan AJ, Balderramo D, Kikuchi L, Gonzalez Ballerga E, Prieto JE, Tapias M, Idrovo V, Davalos MB, Cairo F, Barreyro FJ, Paredes S, Hernandez N, Avendaño K, Diaz Ferrer J, Yang JD, Carrera E, Garcia JA, Mattos AZ, Hirsch BS, Gonçalves PT, Carrilho FJ, Roberts LR. Hepatocellular carcinoma in South America: Evaluation of risk factors, demographics and therapy. Liver Int 2018; 38: 136-143 [PMID: 28640517 DOI: 10.1111/liv.13502]
- de Martel C, Maucort-Boulch D, Plummer M, Franceschi S. World-wide relative contribution of 10 hepatitis B and C viruses in hepatocellular carcinoma. Hepatology 2015; 62: 1190-1200 [PMID: 26146815 DOI: 10.1002/hep.27969]
- GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 11 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet 2020; 396: 1204-1222 [PMID: 33069326 DOI: 10.1016/S0140-6736(20)30925-9]
- Chan AJ, Balderramo D, Kikuchi L, Ballerga EG, Prieto JE, Tapias M, Idrovo V, Davalos MB, Cairo 12 F, Barreyro FJ, Paredes S, Hernandez N, Avendaño K, Ferrer JD, Yang JD, Carrera E, Mattos AZ, Hirsch BS, Gonçalves PT, Carrilho FJ, Roberts LR, Debes JD. Early Age Hepatocellular Carcinoma Associated With Hepatitis B Infection in South America. Clin Gastroenterol Hepatol 2017; 15: 1631-1632 [PMID: 28532694 DOI: 10.1016/j.cgh.2017.05.015]
- Liu Z, Xu K, Jiang Y, Cai N, Fan J, Mao X, Suo C, Jin L, Zhang T, Chen X. Global trend of 13 aetiology-based primary liver cancer incidence from 1990 to 2030: a modelling study. Int J Epidemiol 2021; 50: 128-142 [PMID: 33349860 DOI: 10.1093/ije/dyaa196]
- 14 Park SH, Rehermann B. Immune responses to HCV and other hepatitis viruses. *Immunity* 2014; 40: 13-24 [PMID: 24439265 DOI: 10.1016/j.immuni.2013.12.010]
- 15 Bertoletti A, Ferrari C. Adaptive immunity in HBV infection. J Hepatol 2016; 64: S71-S83 [PMID: 27084039 DOI: 10.1016/j.jhep.2016.01.026]
- Boeijen LL, Hoogeveen RC, Boonstra A, Lauer GM. Hepatitis B virus infection and the immune 16 response: The big questions. Best Pract Res Clin Gastroenterol 2017; 31: 265-272 [PMID: 28774408 DOI: 10.1016/j.bpg.2017.05.003]
- Wieland D, Hofmann M, Thimme R. Overcoming CD8+ T-Cell Exhaustion in Viral Hepatitis: 17 Lessons from the Mouse Model and Clinical Perspectives. Dig Dis 2017; 35: 334-338 [PMID: 28468011 DOI: 10.1159/000456584
- Ringelhan M, Pfister D, O'Connor T, Pikarsky E, Heikenwalder M. The immunology of 18



hepatocellular carcinoma. Nat Immunol 2018; 19: 222-232 [PMID: 29379119 DOI: 10.1038/s41590-018-0044-z]

- Farinati F, Piciocchi M, Lavezzo E, Bortolami M, Cardin R. Oxidative stress and inducible nitric 19 oxide synthase induction in carcinogenesis. Dig Dis 2010; 28: 579-584 [PMID: 21088405 DOI: 10.1159/000320052]
- 20 Unitt E, Rushbrook SM, Marshall A, Davies S, Gibbs P, Morris LS, Coleman N, Alexander GJ. Compromised lymphocytes infiltrate hepatocellular carcinoma: the role of T-regulatory cells. Hepatology 2005; 41: 722-730 [PMID: 15791620 DOI: 10.1002/hep.20644]
- 21 Zhou G, Sprengers D, Boor PPC, Doukas M, Schutz H, Mancham S, Pedroza-Gonzalez A, Polak WG, de Jonge J, Gaspersz M, Dong H, Thielemans K, Pan Q, IJzermans JNM, Bruno MJ, Kwekkeboom J. Antibodies Against Immune Checkpoint Molecules Restore Functions of Tumor-Infiltrating T Cells in Hepatocellular Carcinomas. Gastroenterology 2017; 153: 1107-1119. e10 [PMID: 28648905 DOI: 10.1053/j.gastro.2017.06.017]
- 22 Fu J, Xu D, Liu Z, Shi M, Zhao P, Fu B, Zhang Z, Yang H, Zhang H, Zhou C, Yao J, Jin L, Wang H, Yang Y, Fu YX, Wang FS. Increased regulatory T cells correlate with CD8 T-cell impairment and poor survival in hepatocellular carcinoma patients. Gastroenterology 2007; 132: 2328-2339 [PMID: 17570208 DOI: 10.1053/j.gastro.2007.03.102]
- Tian Z, Chen Y, Gao B. Natural killer cells in liver disease. Hepatology 2013; 57: 1654-1662 [PMID: 23 23111952 DOI: 10.1002/hep.26115]
- Unitt E, Marshall A, Gelson W, Rushbrook SM, Davies S, Vowler SL, Morris LS, Coleman N, 24 Alexander GJ. Tumour lymphocytic infiltrate and recurrence of hepatocellular carcinoma following liver transplantation. J Hepatol 2006; 45: 246-253 [PMID: 16580084 DOI: 10.1016/j.jhep.2005.12.027]
- 25 Flecken T, Schmidt N, Hild S, Gostick E, Drognitz O, Zeiser R, Schemmer P, Bruns H, Eiermann T, Price DA, Blum HE, Neumann-Haefelin C, Thimme R. Immunodominance and functional alterations of tumor-associated antigen-specific CD8+ T-cell responses in hepatocellular carcinoma. Hepatology 2014; 59: 1415-1426 [PMID: 24002931 DOI: 10.1002/hep.26731]
- Wu Y, Kuang DM, Pan WD, Wan YL, Lao XM, Wang D, Li XF, Zheng L. Monocyte/macrophage-26 elicited natural killer cell dysfunction in hepatocellular carcinoma is mediated by CD48/2B4 interactions. Hepatology 2013; 57: 1107-1116 [PMID: 23225218 DOI: 10.1002/hep.26192]
- 27 Lim CJ, Lee YH, Pan L, Lai L, Chua C, Wasser M, Lim TKH, Yeong J, Toh HC, Lee SY, Chan CY, Goh BK, Chung A, Heikenwälder M, Ng IO, Chow P, Albani S, Chew V. Multidimensional analyses reveal distinct immune microenvironment in hepatitis B virus-related hepatocellular carcinoma. Gut 2019; 68: 916-927 [PMID: 29970455 DOI: 10.1136/gutjnl-2018-316510]
- Pinato DJ, Guerra N, Fessas P, Murphy R, Mineo T, Mauri FA, Mukherjee SK, Thursz M, Wong CN, Sharma R, Rimassa L. Immune-based therapies for hepatocellular carcinoma. Oncogene 2020; 39: 3620-3637 [PMID: 32157213 DOI: 10.1038/s41388-020-1249-9]
- 29 Cancer Genome Atlas Research Network. Comprehensive and Integrative Genomic Characterization of Hepatocellular Carcinoma. Cell 2017; 169: 1327-1341. e23 [PMID: 28622513 DOI: 10.1016/j.cell.2017.05.046]
- Schulze K, Imbeaud S, Letouzé E, Alexandrov LB, Calderaro J, Rebouissou S, Couchy G, Meiller C, 30 Shinde J, Soysouvanh F, Calatayud AL, Pinyol R, Pelletier L, Balabaud C, Laurent A, Blanc JF, Mazzaferro V, Calvo F, Villanueva A, Nault JC, Bioulac-Sage P, Stratton MR, Llovet JM, Zucman-Rossi J. Exome sequencing of hepatocellular carcinomas identifies new mutational signatures and potential therapeutic targets. Nat Genet 2015; 47: 505-511 [PMID: 25822088 DOI: 10.1038/ng.3252]
- Hoshida Y, Nijman SM, Kobayashi M, Chan JA, Brunet JP, Chiang DY, Villanueva A, Newell P, 31 Ikeda K, Hashimoto M, Watanabe G, Gabriel S, Friedman SL, Kumada H, Llovet JM, Golub TR. Integrative transcriptome analysis reveals common molecular subclasses of human hepatocellular carcinoma. Cancer Res 2009; 69: 7385-7392 [PMID: 19723656 DOI: 10.1158/0008-5472.CAN-09-1089]
- Chisari FV. Rous-Whipple Award Lecture. Viruses, immunity, and cancer: lessons from hepatitis B. 32 Am J Pathol 2000; 156: 1117-1132 [PMID: 10751335 DOI: 10.1016/s0002-9440(10)64980-2]
- Sung WK, Zheng H, Li S, Chen R, Liu X, Li Y, Lee NP, Lee WH, Ariyaratne PN, Tennakoon C, 33 Mulawadi FH, Wong KF, Liu AM, Poon RT, Fan ST, Chan KL, Gong Z, Hu Y, Lin Z, Wang G, Zhang Q, Barber TD, Chou WC, Aggarwal A, Hao K, Zhou W, Zhang C, Hardwick J, Buser C, Xu J, Kan Z, Dai H, Mao M, Reinhard C, Wang J, Luk JM. Genome-wide survey of recurrent HBV integration in hepatocellular carcinoma. Nat Genet 2012; 44: 765-769 [PMID: 22634754 DOI: 10.1038/ng.2295
- Zhao LH, Liu X, Yan HX, Li WY, Zeng X, Yang Y, Zhao J, Liu SP, Zhuang XH, Lin C, Qin CJ, 34 Zhao Y, Pan ZY, Huang G, Liu H, Zhang J, Wang RY, Wen W, Lv GS, Zhang HL, Wu H, Huang S, Wang MD, Tang L, Cao HZ, Wang L, Lee TL, Jiang H, Tan YX, Yuan SX, Hou GJ, Tao QF, Xu QG, Zhang XQ, Wu MC, Xu X, Wang J, Yang HM, Zhou WP, Wang HY. Genomic and oncogenic preference of HBV integration in hepatocellular carcinoma. Nat Commun 2016; 7: 12992 [PMID: 27703150 DOI: 10.1038/ncomms12992]
- 35 Wang P, Guo QS, Wang ZW, Qian HX. HBx induces HepG-2 cells autophagy through PI3K/AktmTOR pathway. Mol Cell Biochem 2013; 372: 161-168 [PMID: 23001846 DOI: 10.1007/s11010-012-1457-x
- Ching RHH, Sze KMF, Lau EYT, Chiu YT, Lee JMF, Ng IOL, Lee TKW. C-terminal truncated 36 hepatitis B virus X protein regulates tumorigenicity, self-renewal and drug resistance via



STAT3/Nanog signaling pathway. Oncotarget 2017; 8: 23507-23516 [PMID: 28186991 DOI: 10.18632/oncotarget.15183]

- Chen Z, Tang J, Cai X, Huang Y, Gao Q, Liang L, Tian L, Yang Y, Zheng Y, Hu Y, Tang N. HBx 37 mutations promote hepatoma cell migration through the Wnt/β-catenin signaling pathway. Cancer Sci 2016; 107: 1380-1389 [PMID: 27420729 DOI: 10.1111/cas.13014]
- Raimondo G, Locarnini S, Pollicino T, Levrero M, Zoulim F, Lok AS; Taormina Workshop on 38 Occult HBV Infection Faculty Members. Update of the statements on biology and clinical impact of occult hepatitis B virus infection. J Hepatol 2019; 71: 397-408 [PMID: 31004683 DOI: 10.1016/j.jhep.2019.03.034]
- 39 Mak LY, Wong DK, Pollicino T, Raimondo G, Hollinger FB, Yuen MF. Occult hepatitis B infection and hepatocellular carcinoma: Epidemiology, virology, hepatocarcinogenesis and clinical significance. J Hepatol 2020; 73: 952-964 [PMID: 32504662 DOI: 10.1016/j.jhep.2020.05.042]
- Yang JD, Hainaut P, Gores GJ, Amadou A, Plymoth A, Roberts LR. A global view of hepatocellular carcinoma: trends, risk, prevention and management. Nat Rev Gastroenterol Hepatol 2019; 16: 589-604 [PMID: 31439937 DOI: 10.1038/s41575-019-0186-y]
- 41 Marrero JA, Kulik LM, Sirlin CB, Zhu AX, Finn RS, Abecassis MM, Roberts LR, Heimbach JK. Diagnosis, Staging, and Management of Hepatocellular Carcinoma: 2018 Practice Guidance by the American Association for the Study of Liver Diseases. Hepatology 2018; 68: 723-750 [PMID: 29624699 DOI: 10.1002/hep.29913]
- European Association for the Study of the Liver. EASL Clinical Practice Guidelines: Management 42 of hepatocellular carcinoma. J Hepatol 2018; 69: 182-236 [PMID: 29628281 DOI: 10.1016/i.ihep.2018.03.019
- 43 Terrault NA, Lok ASF, McMahon BJ, Chang KM, Hwang JP, Jonas MM, Brown RS Jr, Bzowej NH, Wong JB. Update on Prevention, Diagnosis, and Treatment of Chronic Hepatitis B: AASLD 2018 Hepatitis B Guidance. Clin Liver Dis (Hoboken) 2018; 12: 33-34 [PMID: 30988907 DOI: 10.1002/cld.728]
- 44 Chang MH, You SL, Chen CJ, Liu CJ, Lai MW, Wu TC, Wu SF, Lee CM, Yang SS, Chu HC, Wang TE, Chen BW, Chuang WL, Soon MS, Lin CY, Chiou ST, Kuo HS, Chen DS; Taiwan Hepatoma Study Group. Long-term Effects of Hepatitis B Immunization of Infants in Preventing Liver Cancer. Gastroenterology 2016; 151: 472-480. e1 [PMID: 27269245 DOI: 10.1053/j.gastro.2016.05.048]
- World Health Organization. WHO-UNICEF estimates of HepB3 coverage. [cited 17 January 45 2021]. In: World Health Organization [Internet]. Available from: http://apps.who.int/immunization_monitoring/globalsummary/timeseries/tswucoveragehepb3.html
- Peck M, Gacic-Dobo M, Diallo MS, Nedelec Y, Sodha SV, Wallace AS. Global Routine Vaccination 46 Coverage, 2018. MMWR Morb Mortal Wkly Rep 2019; 68: 937-942 [PMID: 31647786 DOI: 10.15585/mmwr.mm6842a1]
- Kanwal F, Kramer JR, Ilyas J, Duan Z, El-Serag HB. HCV genotype 3 is associated with an 47 increased risk of cirrhosis and hepatocellular cancer in a national sample of U.S. Veterans with HCV. Hepatology 2014; 60: 98-105 [PMID: 24615981 DOI: 10.1002/hep.27095]
- 48 Goossens N, Negro F. Is genotype 3 of the hepatitis C virus the new villain? Hepatology 2014; 59: 2403-2412 [PMID: 24155107 DOI: 10.1002/hep.26905]
- 49 Carrat F. Fontaine H. Dorival C. Simony M. Diallo A. Hezode C. De Ledinghen V. Larrey D. Haour G, Bronowicki JP, Zoulim F, Asselah T, Marcellin P, Thabut D, Leroy V, Tran A, Habersetzer F, Samuel D, Guyader D, Chazouilleres O, Mathurin P, Metivier S, Alric L, Riachi G, Gournay J, Abergel A, Cales P, Ganne N, Loustaud-Ratti V, D'Alteroche L, Causse X, Geist C, Minello A, Rosa I, Gelu-Simeon M, Portal I, Raffi F, Bourliere M, Pol S; French ANRS CO22 Hepather cohort. Clinical outcomes in patients with chronic hepatitis C after direct-acting antiviral treatment: a prospective cohort study. Lancet 2019; 393: 1453-1464 [PMID: 30765123 DOI: 10.1016/S0140-6736(18)32111-1
- Kanwal F, Kramer J, Asch SM, Chayanupatkul M, Cao Y, El-Serag HB. Risk of Hepatocellular 50 Cancer in HCV Patients Treated With Direct-Acting Antiviral Agents. Gastroenterology 2017; 153: 996-1005. e1 [PMID: 28642197 DOI: 10.1053/j.gastro.2017.06.012]
- Ghany MG, Morgan TR; AASLD-IDSA Hepatitis C Guidance Panel. Hepatitis C Guidance 2019 51 Update: American Association for the Study of Liver Diseases-Infectious Diseases Society of America Recommendations for Testing, Managing, and Treating Hepatitis C Virus Infection. Hepatology 2020; 71: 686-721 [PMID: 31816111 DOI: 10.1002/hep.31060]
- Degasperi E, Galmozzi E, Pelusi S, D'Ambrosio R, Soffredini R, Borghi M, Perbellini R, Facchetti F, 52 Iavarone M, Sangiovanni A, Valenti L, Lampertico P. Hepatic Fat-Genetic Risk Score Predicts Hepatocellular Carcinoma in Patients With Cirrhotic HCV Treated With DAAs. Hepatology 2020; 72: 1912-1923 [PMID: 32762045 DOI: 10.1002/hep.31500]
- 53 Alonso López S, Manzano ML, Gea F, Gutiérrez ML, Ahumada AM, Devesa MJ, Olveira A, Polo BA, Márquez L, Fernández I, Cobo JCR, Rayón L, Riado D, Izquierdo S, Usón C, Real Y, Rincón D, Fernández-Rodríguez CM, Bañares R. A Model Based on Noninvasive Markers Predicts Very Low Hepatocellular Carcinoma Risk After Viral Response in Hepatitis C Virus-Advanced Fibrosis. Hepatology 2020; 72: 1924-1934 [PMID: 33022803 DOI: 10.1002/hep.31588]
- 54 Centre for Disease Analysis. Web Annex B. WHO estimates of the prevalence and incidence of hepatitis C virus infection by World Health Organization region, 2015. Geneva: World Health Organization, 2018: 1-68
- 55 World Health Organization. Global Health Sector Strategy on Viral Hepatitis 2016-2021. Towards



Ending Viral Hepatitis. [cited 17 January 2021]. In: World Health Organization [Internet]. Available from: https://apps.who.int/iris/bitstream/handle/10665/246177/WHO-HIV-2016.06eng.pdf?sequence=1

- 56 Indolfi G, Azzari C, Resti M. Perinatal transmission of hepatitis C virus. J Pediatr 2013; 163: 1549-1552. e1 [PMID: 23919905 DOI: 10.1016/j.jpeds.2013.06.077]
- 57 Epstein RL, Sabharwal V, Wachman EM, Saia KA, Vellozzi C, Hariri S, Linas BP. Perinatal Transmission of Hepatitis C Virus: Defining the Cascade of Care. J Pediatr 2018; 203: 34-40. e1 [PMID: 30170857 DOI: 10.1016/j.jpeds.2018.07.006]
- Dore GJ, Hajarizadeh B. Elimination of Hepatitis C Virus in Australia: Laying the Foundation. Infect 58 Dis Clin North Am 2018; 32: 269-279 [PMID: 29778255 DOI: 10.1016/j.idc.2018.02.006]
- 59 Pawlotsky JM, Ramers CB, Dillon JF, Feld JJ, Lazarus JV. Simplification of Care for Chronic Hepatitis C Virus Infection. Semin Liver Dis 2020; 40: 392-402 [PMID: 32725611 DOI: 10.1055/s-0040-1713657
- Blach S, Kondili LA, Aghemo A, Cai Z, Dugan E, Estes C, Gamkrelidze I, Ma S, Pawlotsky JM, 60 Razavi-Shearer D, Razavi H, Waked I, Zeuzem S, Craxi A. Impact of COVID-19 on global HCV elimination efforts. J Hepatol 2021; 74: 31-36 [PMID: 32777322 DOI: 10.1016/j.jhep.2020.07.042]
- Debes JD, de Knegt RJ, Boonstra A. The Path to Cancer and Back: Immune Modulation During Hepatitis C Virus Infection, Progression to Fibrosis and Cancer, and Unexpected Roles of New Antivirals. Transplantation 2017; 101: 910-915 [PMID: 28045877 DOI: 10.1097/TP.000000000001623
- Merchante N, Merino E, Rodríguez-Arrondo F, Tural C, Muñoz J, Delgado-Fernández M, Jover F, Galindo MJ, Rivero A, López-Aldeguer J, Aguirrebengoa K, Romero-Palacios A, Martínez E, Pineda JA. HIV/hepatitis C virus-coinfected patients who achieved sustained virological response are still at risk of developing hepatocellular carcinoma. AIDS 2014; 28: 41-47 [PMID: 24056067 DOI: 10.1097/QAD.0000000000000005]
- 63 Merchante N, Rodríguez-Arrondo F, Revollo B, Merino E, Ibarra S, Galindo MJ, Montero M, García-Deltoro M, Rivero-Juárez A, Téllez F, Delgado-Fernández M, Ríos-Villegas MJ, García MA, Vera-Méndez FJ, Ojeda-Burgos G, López-Ruz MA, Metola L, Omar M, Alemán-Valls MR, Aguirrebengoa K, Portu J, Raffo M, Macías J, Pineda JA; GEHEP-002 Study Group. Hepatocellular carcinoma after sustained virological response with interferon-free regimens in HIV/hepatitis C viruscoinfected patients. AIDS 2018; 32: 1423-1430 [PMID: 29596108 DOI: 10.1097/QAD.000000000001809
- Merchante N, Merino E, López-Aldeguer J, Jover F, Delgado-Fernández M, Galindo MJ, Ortega E, Rivero A, Mínguez C, Romero-Palacios A, Padilla S, Márquez-Solero M, Amador C, Ríos-Villegas MJ, Téllez F, Portilla J, Pineda JA. Increasing incidence of hepatocellular carcinoma in HIV-infected patients in Spain. Clin Infect Dis 2013; 56: 143-150 [PMID: 22955438 DOI: 10.1093/cid/cis777]
- Sahasrabuddhe VV, Shiels MS, McGlynn KA, Engels EA. The risk of hepatocellular carcinoma 65 among individuals with acquired immunodeficiency syndrome in the United States. Cancer 2012; 118: 6226-6233 [PMID: 22736272 DOI: 10.1002/cncr.27694]
- Bräu N, Fox RK, Xiao P, Marks K, Naqvi Z, Taylor LE, Trikha A, Sherman M, Sulkowski MS, 66 Dieterich DT, Rigsby MO, Wright TL, Hernandez MD, Jain MK, Khatri GK, Sterling RK, Bonacini M, Martyn CA, Aytaman A, Llovet JM, Brown ST, Bini EJ; North American Liver Cancer in HIV Study Group. Presentation and outcome of hepatocellular carcinoma in HIV-infected patients: a U.S.-Canadian multicenter study. J Hepatol 2007; 47: 527-537 [PMID: 17692986 DOI: 10.1016/j.jhep.2007.06.010
- 67 Yu ML, Lin SM, Chuang WL, Dai CY, Wang JH, Lu SN, Sheen IS, Chang WY, Lee CM, Liaw YF. A sustained virological response to interferon or interferon/ribavirin reduces hepatocellular carcinoma and improves survival in chronic hepatitis C: a nationwide, multicentre study in Taiwan. Antivir Ther 2006; 11: 985-994 [PMID: 17302368]
- 68 Merchante N, Rodríguez-Fernández M, Pineda JA. Screening for Hepatocellular Carcinoma in HIV-Infected Patients: Current Evidence and Controversies. Curr HIV/AIDS Rep 2020; 17: 6-17 [PMID: 31933273 DOI: 10.1007/s11904-019-00475-0]
- Wandeler G, Mauron E, Atkinson A, Dufour JF, Kraus D, Reiss P, Peters L, Dabis F, Fehr J, 69 Bernasconi E, van der Valk M, Smit C, Gjærde LK, Rockstroh J, Neau D, Bonnet F, Rauch A; Swiss HIV Cohort Study, Athena Observational Cohort Study, EuroSIDA, ANRS CO3 Aquitaine Cohort. Incidence of hepatocellular carcinoma in HIV/HBV-coinfected patients on tenofovir therapy: Relevance for screening strategies. J Hepatol 2019; 71: 274-280 [PMID: 30965070 DOI: 10.1016/j.jhep.2019.03.032
- Tanon A, Jaquet A, Ekouevi DK, Akakpo J, Adoubi I, Diomande I, Houngbe F, Zannou MD, Sasco AJ, Eholie SP, Dabis F, Bissagnene E; IeDEA West Africa Collaboration. The spectrum of cancers in West Africa: associations with human immunodeficiency virus. PLoS One 2012; 7: e48108 [PMID: 23144732 DOI: 10.1371/journal.pone.0048108]
- 71 Maponga TG, Glashoff RH, Vermeulen H, Robertson B, Burmeister S, Bernon M, Omoshoro-Jones J, Ruff P, Neugut AI, Jacobson JS, Preiser W, Andersson MI. Hepatitis B virus-associated hepatocellular carcinoma in South Africa in the era of HIV. BMC Gastroenterol 2020; 20: 226 [PMID: 32660431 DOI: 10.1186/s12876-020-01372-2]
- 72 Clifford GM, Rickenbach M, Polesel J, Dal Maso L, Steffen I, Ledergerber B, Rauch A, Probst-Hensch NM, Bouchardy C, Levi F, Franceschi S; Swiss HIV Cohort. Influence of HIV-related immunodeficiency on the risk of hepatocellular carcinoma. AIDS 2008; 22: 2135-2141 [PMID:



18832877 DOI: 10.1097/QAD.0b013e32831103ad]

- 73 Singal AG, Pillai A, Tiro J. Early detection, curative treatment, and survival rates for hepatocellular carcinoma surveillance in patients with cirrhosis: a meta-analysis. PLoS Med 2014; 11: e1001624 [PMID: 24691105 DOI: 10.1371/journal.pmed.1001624]
- 74 Rodríguez de Lope C, Reig M, Matilla A, Ferrer MT, Dueñas E, Mínguez B, F Castroagudín J, Ortiz I, Pascual S, Lledó JL, Gallego A, Arenas JI, Aracil C, Forne M, Muñoz C, Pons F, Sala M, Iñarrairaegui M, Martin-Llahi M, Andreu V, Garre C, Rendón P, Fuentes J, Crespo J, Rodríguez M, Bruix J, Varela M; en representación del Grupo de Estudio de Cáncer Hepático (GECH). Clinical characteristics of hepatocellular carcinoma in Spain. Comparison with the 2008-2009 period and analysis of the causes of diagnosis out of screening programs. Analysis of 686 cases in 73 centers. Med Clin (Barc) 2017; 149: 61-71 [PMID: 28279536 DOI: 10.1016/j.medcli.2016.12.048]
- 75 Ferrante ND, Lo Re V 3rd. Epidemiology, Natural History, and Treatment of Hepatitis Delta Virus Infection in HIV/Hepatitis B Virus Coinfection. Curr HIV/AIDS Rep 2020; 17: 405-414 [PMID: 32607773 DOI: 10.1007/s11904-020-00508-z]
- Soriano V, Grint D, d'Arminio Monforte A, Horban A, Leen C, Poveda E, Antunes F, de Wit S, 76 Lundgren J, Rockstroh J, Peters L. Hepatitis delta in HIV-infected individuals in Europe. AIDS 2011; 25: 1987-1992 [PMID: 21857493 DOI: 10.1097/QAD.0b013e32834babb3]
- 77 Béguelin C, Moradpour D, Sahli R, Suter-Riniker F, Lüthi A, Cavassini M, Günthard HF, Battegay M, Bernasconi E, Schmid P, Calmy A, Braun DL, Furrer H, Rauch A, Wandeler G; Swiss HIV Cohort Study. Hepatitis delta-associated mortality in HIV/HBV-coinfected patients. J Hepatol 2017; 66: 297-303 [PMID: 27746337 DOI: 10.1016/j.jhep.2016.10.007]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

