

World Journal of *Diabetes*

World J Diabetes 2021 August 15; 12(8): 1141-1362



EDITORIAL

- 1141** Clinical effects of antidiabetic drugs on psoriasis: The perspective of evidence-based medicine
Zhang MX, Zheng BY, Chen HX, Chien CW

REVIEW

- 1146** Gut microbiota as a target for prevention and treatment of type 2 diabetes: Mechanisms and dietary natural products
Xia F, Wen LP, Ge BC, Li YX, Li FP, Zhou BJ
- 1164** Current progress in metabolomics of gestational diabetes mellitus
Wang QY, You LH, Xiang LL, Zhu YT, Zeng Y
- 1187** Progress in treatment of type 2 diabetes by bariatric surgery
Jin ZL, Liu W
- 1200** Genetics of macrovascular complications in type 2 diabetes
Tonyan ZN, Nasykhova YA, Danilova MM, Glotov AS

MINIREVIEWS

- 1220** Place of intravitreal dexamethasone implant in the treatment armamentarium of diabetic macular edema
Karti O, Saatci AO
- 1233** Cardiac changes in infants of diabetic mothers
Al-Biltagi M, El razaky O, El Amrousy D
- 1248** Effect of inflammatory bowel disease treatments on patients with diabetes mellitus
Bower JAJ, O'Flynn L, Kakad R, Aldulaimi D
- 1255** Perioperative challenges in management of diabetic patients undergoing non-cardiac surgery
Galway U, Chahar P, Schmidt MT, Araujo-Duran JA, Shivakumar J, Turan A, Ruetzler K

ORIGINAL ARTICLE

Basic Study

- 1267** Decabromodiphenyl ether causes insulin resistance and glucose and lipid metabolism disorders in mice
Alimu A, Abudureman H, Wang YZ, Li MY, Wang JS, Liu ZL

Case Control Study

- 1282** Clinical significance of serum miR-129-5p in patients with diabetes mellitus presenting macrovascular complications
He XY, Ou CL

Retrospective Study

- 1292** Association of β -cell function and insulin resistance with pediatric type 2 diabetes among Chinese children
Xu ZR, Du HW, Cui LW, Zheng RX, Li GM, Wei HY, Lu FY, Chen LL, Wu CS, Zhang SX, Zhang SL, Liu F, Zhang MY, Pei Z, Sun CJ, Wu J, Luo FH

Observational Study

- 1304** Polymorphisms in HIF-1 α gene are not associated with diabetic retinopathy in China
Liu YH, Guo C, Sun YQ, Li Q

SYSTEMATIC REVIEWS

- 1312** Which predictors could effect on remission of type 2 diabetes mellitus after the metabolic surgery: A general perspective of current studies?
Akkus G, Tetiker T
- 1325** What can we learn from β -cell failure biomarker application in diabetes in childhood? A systematic review
Calderón-Hernández MF, Altamirano-Bustamante NF, Revilla-Monsalve C, Mosquera-Andrade MB, Altamirano-Bustamante MM

ABOUT COVER

Editorial Board Member, Vaia D Raikou, PhD, Chief Doctor, Department of Nephrology, Doctors' Hospital, Athens 11257, Greece. vraikou@med.uoa.gr

AIMS AND SCOPE

The primary aim of *World Journal of Diabetes* (WJD, *World J Diabetes*) is to provide scholars and readers from various fields of diabetes with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of diabetes and covering a wide range of topics including risk factors for diabetes, diabetes complications, experimental diabetes mellitus, type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes, diabetic angiopathies, diabetic cardiomyopathies, diabetic coma, diabetic ketoacidosis, diabetic nephropathies, diabetic neuropathies, Donohue syndrome, fetal macrosomia, and prediabetic state.

INDEXING/ABSTRACTING

The WJD is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJD as 3.763; IF without journal self cites: 3.684; 5-year IF: 7.348; Journal Citation Indicator: 0.64□Ranking: 80 among 145 journals in endocrinology and metabolism; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Lin-YuTong Wang; Production Department Director: Yu-Jie Ma; Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Diabetes

ISSN

ISSN 1948-9358 (online)

LAUNCH DATE

June 15, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Timothy Koch

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1948-9358/editorialboard.htm>

PUBLICATION DATE

August 15, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Clinical effects of antidiabetic drugs on psoriasis: The perspective of evidence-based medicine

Mei-Xian Zhang, Bo-Yuan Zheng, Hai-Xiao Chen, Ching-Wen Chien

ORCID number: Mei-Xian Zhang 0000-0002-6538-7037; Bo-Yuan Zheng 0000-0003-2085-3513; Hai-Xiao Chen 0000-0002-4746-7500; Ching-Wen Chien 0000-0002-0601-4994.

Author contributions: Zhang MX, Zheng BY, Chen HX and Chien CW conducted the study and drafted the manuscript; Zhang MX and Chien CW participated in the design of the study; all of the authors read and approved the final manuscript.

Conflict-of-interest statement: The authors have no proprietary interest in any aspect of this study.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Mei-Xian Zhang, Evidence-based Medicine Center, Taizhou Hospital of Zhejiang Province, Wenzhou Medical University, Linhai 317000, Zhejiang Province, China

Mei-Xian Zhang, Public Laboratory, Taizhou Hospital of Zhejiang Province, Wenzhou Medical University, Linhai 317000, Zhejiang Province, China

Bo-Yuan Zheng, Ching-Wen Chien, Institute for Hospital Management, Tsing Hua University, Shenzhen Campus, Shenzhen 518055, Guangdong Province, China

Hai-Xiao Chen, Department of Orthopedics, Taizhou Hospital of Zhejiang Province, Wenzhou Medical University, Linhai 317000, Zhejiang Province, China

Corresponding author: Ching-Wen Chien, PhD, Professor, Institute for Hospital Management, Tsing Hua University, Shenzhen Campus, No. 2279 Lishui Street, Shenzhen 518055, Guangdong Province, China. ihca@sz.tsinghua.edu.tw

Abstract

Psoriasis and diabetes shared common underlying pathophysiological mechanisms. Emerging data suggested that antidiabetic medications may improve the psoriasis severity in patients with diabetes mellitus. Several hypoglycemic agents including thiazolidinediones, glucagon-like peptide-1 receptor agonists, dipeptidyl peptidase-4 inhibitors, and biguanides have been reported to make a remarkable reduction in the Psoriasis Area and Severity Index score from baseline. This antipsoriatic effect could be mediated not only by the glucose-lowering action of these agents but also *via* inhibition of keratinocyte over proliferation, increase expression of differentiation markers, suppression the immune inflammatory pathway, and blocking the calcium channels and mitogen-activated protein kinase signaling pathways. On the other hand, there was no significant increase in adverse reactions associated with the treatment of pioglitazone or metformin. However, previous studies often had the relatively short duration of the trials, and did not have enough power to assess recurrence of psoriasis. Potential bias in the study and missing data could undermine the reliability of the results. Therefore, the appropriately randomized controlled studies with large sample sizes and long-term durations in various psoriasis patients are warranted for further support.

Key Words: Psoriasis; Hypoglycemic agents; Thiazolidinediones; Glucagon-like peptide-1 receptor agonists; Dipeptidyl peptidase-4 inhibitors; Biguanides; Antipsoriatic effect;

Specialty type: Dermatology**Country/Territory of origin:** China**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

Received: February 3, 2021**Peer-review started:** February 3, 2021**First decision:** March 30, 2021**Revised:** April 8, 2021**Accepted:** July 14, 2021**Article in press:** July 14, 2021**Published online:** August 15, 2021**P-Reviewer:** Mirabelli M, Tung TH**S-Editor:** Liu M**L-Editor:** Filipodia**P-Editor:** Li X

Evidence-based medicine

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Several hypoglycemic agents including thiazolidinediones, glucagon-like peptide-1 receptor agonists, dipeptidyl peptidase-4 inhibitors, and biguanides have been reported to make a remarkable reduction in the Psoriasis Area and Severity Index score from baseline. The antipsoriatic effect could be mediated not only by the glucose-lowering action of these agents but also *via* inhibition of keratinocyte overproliferation, increase expression of differentiation markers, suppression the immune inflammatory pathway, and blocking the calcium channels and mitogen-activated protein kinase signaling pathways. Potential bias in the study and missing data could undermine the reliability of the results. Therefore, the appropriately randomized controlled studies with large sample sizes and long-term durations in various psoriasis patients are warranted for further support.

Citation: Zhang MX, Zheng BY, Chen HX, Chien CW. Clinical effects of antidiabetic drugs on psoriasis: The perspective of evidence-based medicine. *World J Diabetes* 2021; 12(8): 1141-1145

URL: <https://www.wjnet.com/1948-9358/full/v12/i8/1141.htm>

DOI: <https://dx.doi.org/10.4239/wjd.v12.i8.1141>

INTRODUCTION

Psoriasis is a relatively common disorder, though the prevalence varies among populations. An estimated 29.5 million adults worldwide suffered from psoriasis in 2017, equivalent to a lifetime prevalence of 0.59% (95% uncertainty interval 0.19% to 1.66%) diagnosed by physicians in the global adult population. The estimated prevalence of psoriasis was highest in high income countries such as Australasia (1.58%) and Western Europe (1.52%), and lowest in East Asia (0.11%). However, the highest number of adult populations with psoriasis lived in the United States, India, and China[1].

Psoriasis is an immune-mediated chronic inflammatory skin disease. The patients' quality of life was seriously affected by a variety of clinical manifestations including characteristic scaly, and itchy and reddish lesions of skin. Furthermore, psoriasis has also been identified as a systemic disorder that involves more than skin. Diabetes mellitus is one of common extra cutaneous disorders in individuals with psoriasis. Growing evidence supports that psoriasis and diabetes are comorbidities of each other and share common underlying pathophysiological mechanisms[2]. Type 1 diabetes is characterized by autoimmune destruction of the pancreatic beta cells. Type 2 diabetes is characterized by insulin resistance and a relative lack of insulin secretion from the beta cells. An altered adipose tissue microenvironment in obesity is responsible for insulin resistance and low-grade inflammation at either local or systemic level (*e.g.*, skin). That said, diabetes can relate with a wide spectrum of skin conditions, including but not limited to psoriasis, due to hyperglycemia-induced microvascular damage, altered immune functions, infections, genetic predisposition, *etc.* Inflammation can modulate the process of insulin resistance and beta cell damage, and play a critical role in the pathogenesis of type 1 and type 2 diabetes. Thus, antidiabetic drugs with anti-inflammatory properties may also have positive effects on treatment of psoriasis, which is the skin manifestation of an inflammatory response. Emerging data suggested that patients with psoriasis may also benefit significantly from antidiabetic medications due to the common immunological pathway and inflammatory signaling pathway[3-6].

As we know, psoriasis has no cure, so the main purpose of treatment is to control the disease rather than cure. There are many topical and systematic therapies for treatment of psoriatic skin manifestations. However, most of these treatments are expensive, unavailable or immune suppressive. Accumulating data support that hypoglycemic agents including thiazolidinediones (TZDs), glucagon-like peptide-1 (GLP-1) receptor agonists (RAs), dipeptidyl peptidase-4 (DPP-4) inhibitors, and

biguanides, take beneficial effects on the treatment of psoriasis.

ANTIDIABETIC DRUGS ON PSORIASIS

TZDs

The peroxisome proliferator-activated receptor- γ (PPAR- γ) agonist TZDs, like pioglitazone, are insulin sensitizers widely used for treatment of diabetes. In recent years, several double-blinded randomized controlled trial (RCT)s showed that TZDs are anti-inflammatory drugs with ameliorating effects on psoriasis[7-11]. In two meta-analysis of 6 RCTs in patients with psoriasis vulgaris, treatment with pioglitazone made a significant 75% or greater improvement in the Psoriasis Area and Severity Index score from baseline (PASI 75) response[5] and a remarkable reduction in the PASI score from baseline (weighted mean difference: 2.68)[6]; again, there was no increase in side effects associated with pioglitazone treatment such as elevated liver enzymes, weight gain, fatigue, and nausea. That suggested pioglitazone is an effective and safe drug in the treatment of patients with plaque psoriasis. We recently conducted a systemic review and meta-analysis including 10 randomized controlled studies to investigate a more comprehensive result among several kinds of antidiabetic drugs including GLP-1, analog TZDs and metformin for treatment of psoriasis[4]. Our study demonstrated that there was a significant increase in PASI 75 [risk difference = 0.42; 95% confidence interval (CI): 0.18-0.65] or a decrease in PASI score (mean difference = -3.82; 95%CI: -6.05 to -1.59) in psoriasis patients receiving pioglitazone than those intake placebo. But there was no significant effect for rosiglitazone and metformin. As for the different doses of pioglitazone in psoriasis, the daily 30 mg dose was significantly related to a greater therapeutic effect than that of the 15 mg dose per day with no significant change in the frequency of adverse reactions. This means that the amelioration from pioglitazone in psoriasis is dose-dependent[4,12].

There are several pathophysiological mechanisms underlying the antipsoriatic effects of TZDs. First, as PPAR γ -selective ligands, TZDs can suppress human keratinocyte over proliferation, increase the expression of epithelin and intermediate filament related proteins, and then promote skin keratinocytes differentiation[13]. Second, TZDs may be beneficial to psoriasis by reducing the inflammatory response of psoriasis through PPAR- γ . In psoriatic patients, pioglitazone can reduce inflammatory cell infiltration into the skin and reduce the expression of inflammatory cytokines such as C-reactive protein, tumor necrosis factor (TNF) - α and interleukin (IL) 2, and thus inhibit the local immune inflammatory response[14]. Additionally, TZDs could intensify the histology of psoriatic skin *via* interfering the calcium channels and mitogen-activated protein kinase signaling pathways, but not related to the PPAR effect.

GLP-1-based therapies

GLP-1-based therapies such as GLP-1 receptor agonists and inhibitors of DPP-4 are effective hypoglycemic agents. GLP-1 is an incretin hormone that regulates glucose metabolism. DPP-4 is a GLP-1 degrading enzyme and exerts its physiological functions particular in lowering blood glucose *via* the action of GLP-1. Previous study indicated the mechanisms of GLP-1 receptor agonists that determine their antihyperglycemic and anti-inflammatory effects[15]. DPP-4 inhibitors can hinder the DPP-4 to attenuate the degradation of endogenous GLP-1 and prolong the action of GLP-1. A growing body of data shows that GLP-1-based therapies had anti-inflammatory effects in several organs, tissues, and cells through reducing inflammatory cytokine production and immunocyte infiltration. Previous studies on the pathogenesis of psoriasis have focused on keratinocyte excess proliferation, but recently immune system dysfunction has been identified as a critical event. GLP-1 analogues have been reported to alleviate the clinical and histopathology severity of psoriasis through interacting with innate immune system, particularly with invariant natural killer T cells[16-18].

A few case reports showed an interest in using GLP-1 RAs as a potentially option in treatment of psoriasis[18-20]. In a prospective case-series study, seven psoriatic patients with type 2 diabetes following 16-20 wk of GLP-1 RA therapy, alleviated their PASI, and reduced dermal $\gamma\delta$ T-cell counts, and decreased IL-17 expression levels[21]. Such positive effects on psoriasis severity scores was also observed in other prospective case studies of patients receiving GLP-1 RAs for shorter periods[22,23], but the finding was not confirmed in subjects without diabetes[24]. A RCT of 25 patients with psoriasis and type 2 diabetes showed liraglutide treatment for 12 wk led

to a reduction in the PASI score and the relative expression levels of inflammatory cytokines such as IL-23, IL-17, and TNF- α in the damaged skin tissues[17]. However, current studies were small sample size or poorly controlled or short-term follow-up, large-scale multi-center randomized controlled trials with a long duration follow-up are still needed for further verification.

Biguanides

Metformin, a synthetic biguanide, is one of the most popular oral glucose-lowering medications approved for treating type 2 diabetes. Metformin has insulin-sensitizing effect, anti-inflammatory effects and proliferation inhibition in keratinocytes by acting adenosine monophosphate-activated protein kinase. Similar to other antidiabetic drugs, metformin use was also found to decrease the risk of psoriasis in a large case-control cohort study[25]. Clinical findings demonstrated that metformin can significantly improve psoriasis severity by modulating the immune response[14]. A 17-year population-based real-world cohort study focused on the safety of metformin in psoriatic patients with type 2 diabetes and found that metformin did not lead to increased mortality, severe psoriasis, and hospitalizations related to psoriasis. Further dose-response analysis confirmed that metformin is safe for psoriasis patients with diabetes[26]. However, there was only one RCT available for metformin as a potential option in the treatment of psoriasis patients with metabolic syndrome[11,27]. And the efficacy and safety of metformin in other psoriasis patients have not been determined. Further research in various psoriasis patients is clearly warranted to confirm the preliminary findings.

CONCLUSION

In conclusion, the current evidence demonstrates that pioglitazone but not rosiglitazone and metformin is effective in treating plaque psoriasis in diabetic patients. On the other hand, the antidiabetic agents do not appear to increase the incidence of adverse effects. However, the relatively short duration of the trials and no enough power to assess recurrence of psoriasis. Due to some biases such as loss to follow-up were observed in previous cohort studies, more RCTs in psoriasis patients with or without diabetes mellitus with a large sample and long observation time are needed for further support.

REFERENCES

- 1 **Parisi R**, Iskandar IYK, Kontopantelis E, Augustin M, Griffiths CEM, Ashcroft DM; Global Psoriasis Atlas. National, regional, and worldwide epidemiology of psoriasis: systematic analysis and modelling study. *BMJ* 2020; **369**: m1590 [PMID: 32467098 DOI: 10.1136/bmj.m1590]
- 2 **Abramczyk R**, Queller JN, Rachfal AW, Schwartz SS. Diabetes and Psoriasis: Different Sides of the Same Prism. *Diabetes Metab Syndr Obes* 2020; **13**: 3571-3577 [PMID: 33116708 DOI: 10.2147/DMSO.S273147]
- 3 **Ip W**, Kirchhof MG. Glycemic Control in the Treatment of Psoriasis. *Dermatology* 2017; **233**: 23-29 [PMID: 28538228 DOI: 10.1159/000472149]
- 4 **Chi CC**, Lee CY, Liu CY, Wang SH, Tien O'Donnell F, Tung TH. Effects of antidiabetic drugs on psoriasis: A meta-analysis. *Eur J Clin Invest* 2021; **51**: e13377 [PMID: 32914429 DOI: 10.1111/eci.13377]
- 5 **Chang G**, Wang J, Song J, Zhang Z, Zhang L. Efficacy and safety of pioglitazone for treatment of plaque psoriasis: a systematic review and meta-analysis of randomized controlled trials. *J Dermatolog Treat* 2020; **31**: 680-686 [PMID: 31116619 DOI: 10.1080/09546634.2019.1610552]
- 6 **Chen P**, Chen X, Lei L, Zhang Y, Xiang J, Zhou J, Lv J. The efficacy and safety of pioglitazone in psoriasis vulgaris: A meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2020; **99**: e21549 [PMID: 32769894 DOI: 10.1097/MD.00000000000021549]
- 7 **Ghiasi M**, Ebrahimi S, Lajevardi V, Taraz M, Azizpour A. Efficacy and safety of pioglitazone plus phototherapy vs phototherapy in patients with plaque type psoriasis: a Double Blinded Randomized Controlled Trial. *J Dermatolog Treat* 2019; **30**: 664-667 [PMID: 30394148 DOI: 10.1080/09546634.2018.1544702]
- 8 **Hafez VG**, Bosseila M, Abdel Halim MR, Shaker OG, Kamal M, Kareem HS. Clinical effects of "pioglitazone", an insulin sensitizing drug, on psoriasis vulgaris and its co-morbidities, a double blinded randomized controlled trial. *J Dermatolog Treat* 2015; **26**: 208-214 [PMID: 24902962 DOI: 10.3109/09546634.2014.932324]
- 9 **Lajevardi V**, Hallaji Z, Daklan S, Abedini R, Goodarzi A, Abdolreza M. The efficacy of methotrexate plus pioglitazone vs. methotrexate alone in the management of patients with plaque-type

- psoriasis: a single-blinded randomized controlled trial. *Int J Dermatol* 2015; **54**: 95-101 [PMID: 25209868 DOI: 10.1111/ijd.12585]
- 10 **Mittal R**, Malhotra S, Pandhi P, Kaur I, Dogra S. Efficacy and safety of combination Acitretin and Pioglitazone therapy in patients with moderate to severe chronic plaque-type psoriasis: a randomized, double-blind, placebo-controlled clinical trial. *Arch Dermatol* 2009; **145**: 387-393 [PMID: 19380660 DOI: 10.1001/archdermatol.2009.5]
 - 11 **Singh S**, Bhansali A. Randomized placebo control study of insulin sensitizers (Metformin and Pioglitazone) in psoriasis patients with metabolic syndrome (Topical Treatment Cohort). *BMC Dermatol* 2016; **16**: 12 [PMID: 27531132 DOI: 10.1186/s12895-016-0049-y]
 - 12 **Zhang JZ**, Ding Y, Xiang F, Yu SR, Zhang DZ, Guan MM, Kang XJ. Effectiveness and safety of different doses of pioglitazone in psoriasis: a meta-analysis of randomized controlled trials. *Chin Med J (Engl)* 2020; **133**: 444-451 [PMID: 31977550 DOI: 10.1097/CM9.0000000000000642]
 - 13 **Ramot Y**, Mastrofrancesco A, Camera E, Desreumaux P, Paus R, Picardo M. The role of PPAR γ -mediated signalling in skin biology and pathology: new targets and opportunities for clinical dermatology. *Exp Dermatol* 2015; **24**: 245-251 [PMID: 25644500 DOI: 10.1111/exd.12647]
 - 14 **El-Gharabawy RM**, Ahmed AS, Al-Najjar AH. Mechanism of action and effect of immune-modulating agents in the treatment of psoriasis. *Biomed Pharmacother* 2017; **85**: 141-147 [PMID: 27930978 DOI: 10.1016/j.biopha.2016.11.105]
 - 15 **Mirabelli M**, Chiefari E, Puccio L, Foti DP, Brunetti A. Potential Benefits and Harms of Novel Antidiabetic Drugs During COVID-19 Crisis. *Int J Environ Res Public Health* 2020; **17**: 3664 [PMID: 32456064 DOI: 10.3390/ijerph17103664]
 - 16 **Drucker DJ**, Rosen CF. Glucagon-like peptide-1 (GLP-1) receptor agonists, obesity and psoriasis: diabetes meets dermatology. *Diabetologia* 2011; **54**: 2741-2744 [PMID: 21892687 DOI: 10.1007/s00125-011-2297-z]
 - 17 **Lin L**, Xu X, Yu Y, Ye H, He X, Chen S, Chen X, Shao Z, Chen P. Glucagon-like peptide-1 receptor agonist liraglutide therapy for psoriasis patients with type 2 diabetes: a randomized-controlled trial. *J Dermatolog Treat* 2020; **1**-7 [PMID: 32962477 DOI: 10.1080/09546634.2020.1826392]
 - 18 **Faurschou A**, Knop FK, Thyssen JP, Zachariae C, Skov L, Vilsbøll T. Improvement in psoriasis after treatment with the glucagon-like peptide-1 receptor agonist liraglutide. *Acta Diabetol* 2014; **51**: 147-150 [PMID: 22160246 DOI: 10.1007/s00592-011-0359-9]
 - 19 **Hogan AE**, Tobin AM, Ahern T, Corrigan MA, Gaoatswe G, Jackson R, O'Reilly V, Lynch L, Doherty DG, Moynagh PN, Kirby B, O'Connell J, O'Shea D. Glucagon-like peptide-1 (GLP-1) and the regulation of human invariant natural killer T cells: lessons from obesity, diabetes and psoriasis. *Diabetologia* 2011; **54**: 2745-2754 [PMID: 21744074 DOI: 10.1007/s00125-011-2232-3]
 - 20 **Buysschaert M**, Tennstedt D, Preumont V. Improvement of psoriasis during exenatide treatment in a patient with diabetes. *Diabetes Metab* 2012; **38**: 86-88 [PMID: 22227407 DOI: 10.1016/j.diabet.2011.11.004]
 - 21 **Buysschaert M**, Baeck M, Preumont V, Marot L, Hendrickx E, Van Belle A, Dumoutier L. Improvement of psoriasis during glucagon-like peptide-1 analogue therapy in type 2 diabetes is associated with decreasing dermal $\gamma\delta$ T-cell number: a prospective case-series study. *Br J Dermatol* 2014; **171**: 155-161 [PMID: 24506139 DOI: 10.1111/bjd.12886]
 - 22 **Xu X**, Lin L, Chen P, Yu Y, Chen S, Chen X, Shao Z. Treatment with liraglutide, a glucagon-like peptide-1 analogue, improves effectively the skin lesions of psoriasis patients with type 2 diabetes: A prospective cohort study. *Diabetes Res Clin Pract* 2019; **150**: 167-173 [PMID: 30844468 DOI: 10.1016/j.diabres.2019.03.002]
 - 23 **Ahern T**, Tobin AM, Corrigan M, Hogan A, Sweeney C, Kirby B, O'Shea D. Glucagon-like peptide-1 analogue therapy for psoriasis patients with obesity and type 2 diabetes: a prospective cohort study. *J Eur Acad Dermatol Venereol* 2013; **27**: 1440-1443 [PMID: 22691169 DOI: 10.1111/j.1468-3083.2012.04609.x]
 - 24 **Faurschou A**, Gyldenløve M, Rohde U, Thyssen JP, Zachariae C, Skov L, Knop FK, Vilsbøll T. Lack of effect of the glucagon-like peptide-1 receptor agonist liraglutide on psoriasis in glucose-tolerant patients--a randomized placebo-controlled trial. *J Eur Acad Dermatol Venereol* 2015; **29**: 555-559 [PMID: 25139195 DOI: 10.1111/jdv.12629]
 - 25 **Brauchli YB**, Jick SS, Curtin F, Meier CR. Association between use of thiazolidinediones or other oral antidiabetics and psoriasis: A population based case-control study. *J Am Acad Dermatol* 2008; **58**: 421-429 [PMID: 18194825 DOI: 10.1016/j.jaad.2007.11.023]
 - 26 **Su YJ**, Chen TH, Hsu CY, Chiu WT, Lin YS, Chi CC. Safety of Metformin in Psoriasis Patients With Diabetes Mellitus: A 17-Year Population-Based Real-World Cohort Study. *J Clin Endocrinol Metab* 2019; **104**: 3279-3286 [PMID: 30779846 DOI: 10.1210/je.2018-02526]
 - 27 **Singh S**, Bhansali A. Randomized Placebo Control Study of Metformin in Psoriasis Patients with Metabolic Syndrome (Systemic Treatment Cohort). *Indian J Endocrinol Metab* 2017; **21**: 581-587 [PMID: 28670544 DOI: 10.4103/ijem.IJEM_46_17]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

