

国内版国际版

Microsoft Bing

Medication Adherence and quality of life among Type-2 diabetes n

ALLIMAGESVIDEOS

Add the Give

233,000 ResultsAny time ▾

Assessment of patient medication adherence among the ...
<https://www.sciencedirect.com/science/article/pii/S1658361217300185>
Apr 01, 2017 · In absolute numbers, in contrast to the estimated global prevalence of 220 million cases of diabetes by 2010, DPN is likely to affect as many as 110 million people worldwide. 7 In India, studies revealed that diabetic neuropathy occurs in 19%–27.5% of patients with type 2 diabetes. 8 Glycaemic control is crucial for individuals with diabetes to prevent the progression of neuropathy, and intensive ...
Cited by: 2 Author: Anu M. Samu, Palanisamy S. Amirthaling...
Publish Year: 2017

Search Tools
Turn off Hover Translation (关闭取词)

Factors affecting self-medication practices among people ...
<https://www.sciencedirect.com/science/article/pii/S2589936820300530>

84,500 Results

Any time ▾

Assessment of patient medication adherence among the type ...

<https://www.sciencedirect.com/science/article/pii/S1658361217300185>

Apr 01, 2017 · Assessment of patient **medication adherence among the type 2 diabetes mellitus** population with peripheral diabetic neuropathy in South **India** ... collected from all patients by means of the questionnaire during the interviews were thoroughly analysed in regard to **medication adherence** and the **quality of life** of the patients. ... **Adherence** to ...

Cited by: 2

Author: Anu M. Samu, Palanisamy S. Amirthalinga...

Publish Year: 2017

Self Care and Medication Adherence among Type 2 Diabetics ...

<https://europepmc.org/articles/PMC4064856> ▾

Apr 15, 2014 · Prevalence of **diabetes mellitus in India** varies from 5.6% in rural areas to 12.1% in major cities [2,3]. It has been projected that the greatest absolute increase which will occur in number of people with **diabetes**, will be in **India** . Due to sheer numbers, the economic burden caused by **diabetes in India** is among the highest in the world.

Estimated Reading Time: 9 mins

[PDF] Assessment of Factors Influencing Adherence to ...

<https://pdfs.semanticscholar.org/7971/68a5ff2b8bda...>

Assessment of Factors Influencing **Adherence** to Antidiabetic Drugs **Among** Patients with **Type 2 Diabetes Mellitus** at a Tertiary Care Hospital in **India** Nandini HC, Akshay Gali, Sushma Muraraiah
Department of Pharmacology, Bangalore Medical College and Research Institute, Bangalore, Karnataka

国内版

国际版

Medication adherence and quality of life among type-2 diabetes m



ALL

IMAGES

VIDEOS

320,000 Results

Any time ▾

Self Care and Medication Adherence among Type 2 Diabetics ...

<https://europepmc.org/articles/PMC4064856> ▾

Apr 15, 2014 · Prevalence of **diabetes mellitus in India** varies from 5.6% in rural areas to 12.1% in major cities [2,3]. It has been projected that the greatest absolute increase which will occur in number of people with **diabetes**, will be in **India** . Due to sheer numbers, the economic burden caused by **diabetes in India** is among the highest in the world.

Estimated Reading Time: 9 mins

Self Care and Medication Adherence among Type 2 Diabetics ...

<https://www.semanticscholar.org/paper/Self-Care...>

BACKGROUND Micro and macro-vascular complications of **Type 2 Diabetes mellitus (DM)** could be decreased by maintaining a good glycaemic control, which is dependent on **adherence to medication** and self care. AIM (1) To assess **medication adherence** and **adherence to self care among type 2** diabetics who were admitted to a tertiary care hospital (2) To identify factors which were associated ...

PEOPLE ALSO ASK

What causes poor medication adherence in type 2 diabetes? ▾

Is it true that diabetes patients prefer adherence? ▾

What are the effects of poor medication adherence? ▾

How does medication adherence affect patients with T2D? ▾

Feedback

Assessment of patient medication adherence among the ...

<https://www.sciencedirect.com/science/article/pii/S1658361217300185>

Apr 01, 2017 · In absolute numbers, in contrast to the estimated global prevalence of 220 million cases of **diabetes** by 2010, DPN is likely to affect as many as 110 million people worldwide. 7 In **India**, studies revealed that diabetic neuropathy occurs in 19%–27.5% of patients with **type 2 diabetes**. 8 Glycaemic control is crucial for individuals with **diabetes** to prevent the progression of neuropathy, and intensive ...

Cited by: 2

Author: Anu M. Samu, Palanisamy S. Amirthalingam