

3
Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 64724

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

**Long-term outcome of pancreatic function following oncological surgery in children:
Institutional experience and review of the literature**

Pediatric pancreatic function following oncological surgery

Abstract

BACKGROUND

Pancreatic neoplasms are uncommon in children and in most cases they are benign or at low malignant potential. Pancreatoblastoma and Solid Pseudopapillary Tumor are the most frequent types in early and late childhood respectively. Complete resection, although burdened by severe complications, is the only curative treatment for these

Match Overview

- | Match Number | Source | Words | Similarity |
|--------------|---|----------|------------|
| 1 | Crossref | 44 words | 2% |
| | Chiara Scandavini, Roberto Valente, Elena Rangelova, Ralf Se gersvärd et al. "Pancreatectomies for pancreatic neoplasms..." | | |
| 2 | Crossref | 44 words | 2% |
| | Erika B. Lindholm, Abdulaziz K. Alkattan, Sara J. Abramson, A nita P. Price et al. "Pancreaticoduodenectomy for pediatric a..." | | |
| 3 | Internet | 17 words | 1% |
| | crawled on 05-Apr-2021
f6publishing.blob.core.windows.net | | |

Long-term outcome of pancreatic function following oncological su



ALL

IMAGES

VIDEOS

2,310,000 Results

Any time ▾

Long-Term Pancreatic Functional Impairment after Surgery ...

<https://www.mdpi.com/2077-0383/8/10/1611/htm> ▾



Radical **surgery** represents the only curative treatment for **pancreatic** neuroendocrine neoplasms (PanNEN). The aim of this study was to evaluate the postoperative onset of diabetes mellitus (DM) and/or **pancreatic** exocrine insufficiency (PEI) in surgically treate...

Cited by: 6

Author: Valentina Andreasi, Stefano Partelli...

Publish Year: 2019

Long-term outcomes after supratotal resection of diffuse ...

<https://pubmed.ncbi.nlm.nih.gov/26530708>

Here, the goal is to analyze the **long-term** functional and **oncological outcomes following** supratotal resection for DLGG. Methods: Sixteen consecutive patients who underwent supratotal resection for a DLGG with a minimum follow-up of 8 years **after surgery** were included. The resection was continued up to functional cortical and subcortical ...

Cited by: 173

Author: Hugues Duffau

Publish Year: 2016

PEOPLE ALSO ASK

What is the prognosis for someone with pancreatic cancer? ▾

How long does it take to recover from pancreatic surgery? ▾



ALL

IMAGES

VIDEOS

231,000 Results

Any time ▾

Long-Term Pancreatic Functional Impairment after Surgery ...

<https://www.mdpi.com/2077-0383/8/10/1611/htm> ▾

Radical surgery represents the only curative treatment for **pancreatic** neuroendocrine neoplasms (PanNEN). The aim of this study was to evaluate the postoperative onset of diabetes mellitus (DM) and/...

Cited by: 5

Author: Valentina Andreasi, Stefano Partelli, Gabr...

Publish Year: 2019

What indication, morbidity and mortality for central ...

<https://www.sciencedirect.com/science/article/pii/S1743919115014314>

Apr 01, 2016 · Non-traditional **pancreatic** resections, like **pancreatic** enucleation, aim to reduce perioperative morbidity and to ensure good metabolic functions in **long-term outcome** , . Central...

Cited by: 13

Author: Michele Santangelo, Anna Esposito, Vinc...

Publish Year: 2016

(PDF) Pediatric pancreatic teratoma: A case report and ...

<https://www.researchgate.net/publication/337269248...>

Purpose: The purpose of this study was to analyze the **long-term outcomes**, such as nutritional status, **pancreatic function**, gastrointestinal (GI) function, and quality of life (QOL), in children ...

Operative vs Nonoperative Management for Blunt Pancreatic ...

<https://www.sciencedirect.com/science/article/pii/S1072751513011551>

Feb 01, 2014 · Excellent **long-term pancreatic function** has been reported in 9 patients managed nonoperatively, but the larger concern lies with those **children** undergoing **pancreatic** resection. 2...

Cited by: 67

Author: Corey W. Iqbal, Shawn D. St Peter, Kuoje...

Publish Year: 2014

Indications and outcomes of duodenum-preserving resection ...

<https://link.springer.com/article/10.1007/s00383-018-4410-6> ▾

Nov 01, 2018 · Duodenum-preserving resection of the **pancreatic** head (DPRPH) with Roux-en-Y pancreatojejunostomy is a procedure used to remove focal pathological lesions of the **pancreatic** head...

Cited by: 1

Author: Jiri Snajdauf, Michal Rygl, Ondrej Petru, J...


Publish Year: 2019

2,350,000 Results

Any time ▾

Long-Term Pancreatic Functional Impairment after Surgery ...

<https://www.mdpi.com/2077-0383/8/10/1611/htm> ▾



Radical **surgery** represents the only curative treatment for **pancreatic** neuroendocrine neoplasms (PanNEN). The aim of this study was to evaluate the postoperative onset of diabetes mellitus (DM) and/or **pancreatic** exocrine insufficiency (PEI) in surgically treate...

Cited by: 6

Author: Valentina Andreasi, Stefano Partelli...

Publish Year: 2019

II: 89 (32)

Long-term outcomes after supratotal resection of diffuse ...

<https://pubmed.ncbi.nlm.nih.gov/26530708>

Here, the goal is to analyze the **long-term** functional and **oncological outcomes following** supratotal resection for DLGG. Methods: Sixteen consecutive patients who underwent supratotal resection for a DLGG with a minimum follow-up of 8 years **after surgery** were included. The resection was continued up to functional cortical and subcortical ...

Cited by: 174

Author: Hugues Duffau, Hugues Duffau

Publish Year: 2016

PEOPLE ALSO ASK

What is the prognosis for someone with pancreatic cancer?

▾

Are there any long term complications from pancreatic surgery?

▾

What to expect after surgery for pancreatic cancer?

▾

What are the risk factors for developing pancreatic cancer?

▾

Feedback

Ovarian torsion in children: management and outcomes

<https://pubmed.ncbi.nlm.nih.gov/24074673>

Background: The purpose of this study is to evaluate the clinical symptoms, diagnosis, management, and **outcomes in children** with ovarian torsion. Methods: The charts of 50 patients with 53 cases of ovarian torsion treated between January 1989 and March 2012 were reviewed retrospectively. **Long** term follow up was available for 20 girls who had their ovaries left in the abdominal cavity **after**