

World Journal of *Gastroenterology*

World J Gastroenterol 2021 July 7; 27(25): 3693-3950



OPINION REVIEW

- 3693** Approach to medical therapy in perianal Crohn's disease
Vasudevan A, Bruining DH, Loftus EV Jr, Faubion W, Ehman EC, Raffals L

REVIEW

- 3705** Incorporating mucosal-associated invariant T cells into the pathogenesis of chronic liver disease
Czaja AJ
- 3734** Artificial intelligence in small intestinal diseases: Application and prospects
Yang Y, Li YX, Yao RQ, Du XH, Ren C
- 3748** Impact of the COVID-19 pandemic on inflammatory bowel disease patients: A review of the current evidence
Kumric M, Ticinovic Kurir T, Martinovic D, Zivkovic PM, Bozic J
- 3762** Management of hepatitis B virus infection in patients with inflammatory bowel disease under immunosuppressive treatment
Axiaris G, Zampeli E, Michopoulos S, Bamias G

MINIREVIEWS

- 3780** Worldwide management of hepatocellular carcinoma during the COVID-19 pandemic
Inchingolo R, Acquafredda F, Tedeschi M, Laera L, Surico G, Surgo A, Fiorentino A, Spiliopoulos S, de'Angelis N, Memeo R
- 3790** Human immune repertoire in hepatitis B virus infection
Zhan Q, Xu JH, Yu YY, Lo KK E, Felicianna, El-Nezami H, Zeng Z
- 3802** Emerging applications of radiomics in rectal cancer: State of the art and future perspectives
Hou M, Sun JH
- 3815** Advances in paediatric nonalcoholic fatty liver disease: Role of lipidomics
Di Sessa A, Riccio S, Pirozzi E, Verde M, Passaro AP, Umano GR, Guarino S, Miraglia del Giudice E, Marzuillo P
- 3825** Autoimmune pancreatitis and pancreatic cancer: Epidemiological aspects and immunological considerations
Poddighe D
- 3837** Gut microbiota in obesity
Liu BN, Liu XT, Liang ZH, Wang JH

ORIGINAL ARTICLE**Basic Study**

- 3851** Zinc oxide nanoparticles reduce the chemoresistance of gastric cancer by inhibiting autophagy

Miao YH, Mao LP, Cai XJ, Mo XY, Zhu QQ, Yang FT, Wang MH

- 3863** PPARGC1A rs8192678 G>A polymorphism affects the severity of hepatic histological features and nonalcoholic steatohepatitis in patients with nonalcoholic fatty liver disease

Zhang RN, Shen F, Pan Q, Cao HX, Chen GY, Fan JG

Retrospective Cohort Study

- 3877** Does endoscopic intervention prevent subsequent gastrointestinal bleeding in patients with left ventricular assist devices? A retrospective study

Palchoudhuri S, Dhawan I, Parsikia A, Birati EY, Wald J, Siddique SM, Fisher LR

Retrospective Study

- 3888** Diverse expression patterns of mucin 2 in colorectal cancer indicates its mechanism related to the intestinal mucosal barrier

Gan GL, Wu HT, Chen WJ, Li CL, Ye QQ, Zheng YF, Liu J

- 3901** Clinical characteristics of patients in their forties who underwent surgical resection for colorectal cancer in Korea

Lee CS, Baek SJ, Kwak JM, Kim J, Kim SH

Observational Study

- 3913** Effect of gastric microbiota on quadruple *Helicobacter pylori* eradication therapy containing bismuth

Niu ZY, Li SZ, Shi YY, Xue Y

META-ANALYSIS

- 3925** Endoscopic submucosal dissection *vs* endoscopic mucosal resection for colorectal polyps: A meta-analysis and meta-regression with single arm analysis

Lim XC, Nistala KRY, Ng CH, Lin SY, Tan DJH, Ho KY, Chong CS, Muthiah M

CASE REPORT

- 3940** Gastric schwannoma treated by endoscopic full-thickness resection and endoscopic purse-string suture: A case report

Lu ZY, Zhao DY

LETTER TO THE EDITOR

- 3948** Gastrointestinal cytomegalovirus disease secondary to measles in an immunocompetent infant

Hung CM, Lee PH, Lee HM, Chiu CC

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Paola De Nardi, MD, FASCRS, Doctor, Surgeon, Surgical Oncologist, Division of Gastrointestinal Surgery, IRCCS San Raffaele Scientific Institute, via Olgettina 60, Milan 20132, Italy. denardi.paola@hsr.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2021 edition of Journal Citation Report® cites the 2020 impact factor (IF) for WJG as 5.742; Journal Citation Indicator: 0.79; IF without journal self cites: 5.590; 5-year IF: 5.044; Ranking: 28 among 92 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2020 is 6.9 and Scopus CiteScore rank 2020: Gastroenterology is 19/136.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ying-Yi Yuan; Production Department Director: Xiang Li; Editorial Office Director: Ze-Mao Gong.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, Subrata Ghosh

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

July 7, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Endoscopic submucosal dissection vs endoscopic mucosal resection for colorectal polyps: A meta-analysis and meta-regression with single arm analysis

Xiong Chang Lim, Kameswara Rishi Yeshayahu Nistala, Cheng Han Ng, Snow Yunni Lin, Darren Jun Hao Tan, Khek-Yu Ho, Choon-Seng Chong, Mark Muthiah

ORCID number: Xiong Chang Lim 0000-0002-0812-891X; Kameswara Rishi Yeshayahu Nistala 0000-0003-1299-8712; Cheng Han Ng 0000-0002-8297-1569; Snow Yunni Lin 0000-0002-2759-2662; Darren Jun Hao Tan 0000-0001-5358-524X; Khek-Yu Ho 0000-0002-2932-1962; Choon-Seng Chong 0000-0003-0669-7307; Mark Muthiah 0000-0002-9724-4743.

Author contributions: Lim XC, Nistala KRY, Ng CH, Ho KY, Muthiah M and Chong CS were involved in the conceptualization of the paper; Lim XC and Nistala KRY were involved in the acquisition of data; Lim XC, Nistala KRY, Ng CH, Lin SY, Tan DJH, Ho KY, Muthiah M and Chong CS were involved in the data analysis and interpretation; Lim XC, Nistala KRY and Ng CH were involved in writing of first draft; Ho KY, Muthiah M, Chong CS were involved in the reviewing of the drafts; all eight authors approved the final manuscript.

Conflict-of-interest statement: The authors deny any conflict of interest.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript

Xiong Chang Lim, Kameswara Rishi Yeshayahu Nistala, Cheng Han Ng, Snow Yunni Lin, Darren Jun Hao Tan, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117597, Singapore

Khek-Yu Ho, Mark Muthiah, Department of Medicine, National University Hospital, Singapore 119074, Singapore

Choon-Seng Chong, Department of Surgery, National University Hospital, Singapore 119074, Singapore

Corresponding author: Mark Muthiah, MBBS, MRCP, MMed, Assistant Professor, Department of Medicine, National University Hospital, 5 Lower Kent Ridge Road, Singapore 119074, Singapore. mdcmdm@nus.edu.sg

Abstract

BACKGROUND

Endoscopic submucosal dissection (ESD) has shown to be effective in management of colorectal neoplasm in the Asian countries, while its implementation in Western countries where endoscopic mucosal resection (EMR) is preferred is still debatable.

AIM

To compare the surgical, histological, and oncological outcomes between ESD and EMR in the treatment of colorectal polyps, with subgroup analysis comparing the efficacy of ESD and EMR between Japan and the rest of the world.

METHODS

Embase and Medline databases were searched from inception to October 2020 in accordance with PRISMA guidelines for studies comparing *en bloc*, complete resection, margin involvement, resection time, need for additional surgery, complications, and recurrence rate of ESD with EMR.

RESULTS

Of 281344 colorectal polyps from 21 studies were included. When compared to EMR, the pooled analysis revealed ESD was associated with higher *en bloc* and complete resection rate, and lower lateral margin involvement and recurrence.

was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: Singapore

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

Received: February 24, 2021

Peer-review started: February 24, 2021

First decision: April 18, 2021

Revised: April 21, 2021

Accepted: June 1, 2021

Article in press: June 1, 2021

Published online: July 7, 2021

P-Reviewer: Protopapas AA

S-Editor: Fan JR

L-Editor: A

P-Editor: Wang LL



ESD led to increased procedural time, need for additional surgery, and perforation risk. No significant difference in bleeding risk was found between the two groups. Meta-regression analysis suggested only right colonic polyps correlated with an increased perforation risk in ESD. Confounders including polyp size and invasion depth did not significantly influence the *en bloc* and complete resection rate, bleeding risk and recurrence. In subgroup analysis, Japan performed better than the rest of the world in both ESD and EMR with perforation risk of 4% and 0.0002%, respectively, as compared to perforation risk of 8% and 1%, respectively, in reports coming from rest of the world.

CONCLUSION

ESD resulted in better resection outcomes and lower recurrence compared to EMR. With appropriate training, ESD is preferred over EMR as the first-line therapy for resection of colorectal polyps, without restricting to lesions greater than 20 mm and those with high suspicion of submucosal invasion.

Key Words: Endoscopic mucosal resection; Endoscopy; Colonic polyps; Colorectal neoplasm; Colonoscopy

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The present study is the most extensive meta-analysis evaluating the surgical, histological, and oncological outcomes of endoscopic submucosal dissection (ESD) in comparison to endoscopic mucosal resection (EMR) in the treatment of colorectal polyps. Our analysis also showed the increased proficiency in performing ESD and EMR in Japan as compared to the rest of the world.

Citation: Lim XC, Nistala KRY, Ng CH, Lin SY, Tan DJH, Ho KY, Chong CS, Muthiah M. Endoscopic submucosal dissection vs endoscopic mucosal resection for colorectal polyps: A meta-analysis and meta-regression with single arm analysis. *World J Gastroenterol* 2021; 27(25): 3925-3939

URL: <https://www.wjgnet.com/1007-9327/full/v27/i25/3925.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v27.i25.3925>

INTRODUCTION

In recent years, the incidence of colorectal polyps has increased drastically with the widespread implementation of national colorectal cancer (CRC) screening programs. CRC screening with colonoscopy provides the opportunity to identify and remove any precursor lesions[1] and polypectomy has been shown to reduce the incidence and mortality of CRC[1-3]. Currently, 2 major forms of polypectomy are performed, namely endoscopic submucosal dissection (ESD) and endoscopic mucosal resection (EMR) with the latter procedure favored by western countries.

ESD was first proposed in 1999 as an endoscopic resection technique for safe *en bloc* removal of superficial lesions in the upper gastrointestinal tract[4]. Although colorectal ESD is widely practiced in the Asian countries, the implementation of ESD in Western countries where endoscopic mucosal resection is preferred is still debatable. According to the 2015 European Society of Gastrointestinal Endoscopy (ESGE), ESD is strongly recommended when there is high suspicion of submucosal invasion as determined by morphologic factors and advanced surface pattern, especially in lesions greater than 20 mm[5]. Despite the technical difficulties faced, ESD is designed for *en bloc* resection of lesions in an attempt to achieve complete resection, reliable histopathological analysis, and reduced recurrence rate.

Previous meta-analyses and systematic reviews comparing ESD with EMR mainly focused on data reported from Asian countries with limited data coming from outside of Japan, without controlling for confounders or lesions < 20 mm[6,7]. Since then, numerous studies comparing between ESD and EMR have been reported from the West and other countries[8-16]. We sought to perform an expanded meta-analysis to include lesions ≤ 20 mm, and incorporate regression analysis to control for

confounders.

MATERIALS AND METHODS

Search strategy and inclusion criteria

A search was conducted with reference to the PRISMA methodology on October 4, 2020 in Medline and Embase[17]. Keywords and Mesh terms were employed in the search strategy relating to the 'Colorectal', 'ESD' and 'EMR', the full list of terms included in the search can be found in the [Supplementary Material 1](#). References were managed with Endnote X9. Two authors (XCL, KRYN) were involved in the sieving of the abstracts. Articles that compared between EMR and ESD for colorectal polyps were included in the search. Study designs included both retrospective and prospective cohorts, and case-controlled studies. Editorials, conferences abstracts, and commentaries were excluded in the review.

Data extraction

Based on the pre-determined inclusion criterion, article selection was performed by two authors with consensus from an independent third author should discrepancies arise. Information pertaining to the demographics (country, sample size, age, gender, polyp location, polyp size, polyp macroscopic type and depth of cancer invasion) and outcomes were extracted. Outcomes were limited to *en bloc* resection, complete resection, lateral and vertical margin involvement, lymphovascular invasion, mean operation time, additional surgery required, perforation, bleeding, and recurrence. *En bloc* resection was defined as the resection of the lesion in a single specimen, and complete resection was defined as the resection of the lesion with no marginal involvement of neoplastic tissue, as assessed by the pathologist. Additional surgical operations performed were considered in cases with intra-operative complications as a result of ESD or EMR or those with technical difficulties. Bleeding was considered both intra-operative and post-operative, and perforation was considered when diagnosed endoscopically during resection or radiologically by the presence of free air. Recurrence was defined as the detection of local or secondary primary tumors on interval colonoscopy. When estimates of mean and standard deviation (SD) were not available for continuous data, well established formulae were used to estimate mean and SD from median and range[18].

Statistical analysis and quality assessment

Three type of analysis were performed in this review. Firstly, a meta-analysis of proportions was used to pool results from the individual arms (ESD and EMR) using a freeman turkey double arsine transformation[19]. The freeman turkey double arsine transformation is recommended for single arm meta-analysis and necessary to stabilize the variance[20]. Next, comparative meta-analysis after a 0.5 continuity correction was to compare the difference in ESD and EMR, using risk ratios (RR) as the outcome of interest for dichotomous variables and weighted mean differences for continuous variables[21]. A 0.5 continuity correction was considered appropriate to incorporate 0 events into the meta-analysis. Regardless of heterogeneity measures (I^2 , tau, Cochran Q test), all analysis was performed in random effects with the Dersimonian and Laird model[22,23]. Region (Japan *vs* rest of the world) was defined with refences to the manuscript country of origin. Publication bias was assessed with harbord test and egger regression for dichotomous and continuous variables respectively when > 10 studies were available. A univariate meta regression was also conducted with in random effects using the restricted maximum likelihood model with Kapp variance estimator to adjust for confounders in the outcomes[24]. A subgroup analysis was considered to compare between the articles originating from Japan and outside of Japan and a sensitivity analysis was performed to only include studies that have polyps size of ≥ 20 mm. All articles were graded with the Newcastle Ottawa scale for cohort studies, grading articles on the domains of study group selection, study group comparability, and ascertainment of outcomes of interest[25]. The statistical methods of this study were reviewed by Cheng Han Ng (ORCID ID: 0000-0002-8297-1569) from Yong Loo Lin School of Medicine, National University of Singapore.

RESULTS

Study selection

A systematic search of the literature using our search strategy yielded 1180 articles. After removal of duplicates, 895 were excluded based on study title and abstract, and 212 underwent full text review, of which 21 articles comparing ESD with EMR in resection of colorectal polyps were subsequently included in the meta-analysis (Figure 1)[8-16,26-37]. In total, 12 originated from Japan[26-37], five from European[10-13,16], three from South Korea[8,9,14] and one from China[15]. Of the 21 studies, there were 17 retrospective cohort studies, two prospective cohort studies and two retrospective case-control studies. In total, 281344 colorectal polyps were resected, of which 19573 underwent ESD while 261771 underwent EMR. Quality assessment of those included studies mostly scored 6-7. A summary of the study characteristics of the included articles can be found in the [Supplementary Table 1](#).

Overall results

The summary of the results is found in [Table 1](#). The pooled analysis showed a higher rate of *en bloc* resection (RR = 1.837; 95%CI: 1.464 to 2.305; $P < 0.001$), and a lower frequency of positive lateral margin involvement (RR = 0.292; 95%CI: 0.089 to 0.995; $P = 0.042$) in the ESD group than the EMR group. Publication bias was significant for *en bloc* resection rate ($P = 0.0025$). No significant difference in the rates of positive vertical margin involvement was observed between ESD and EMR groups (RR = 4.368; 95%CI: 0.409 to 46.710; $P = 0.223$). However, the rate of complete resection was higher in the ESD compared to EMR groups (RR = 1.504; 95%CI: 1.041 to 2.174; $P < 0.03$). Significantly, the time taken for ESD was longer than EMR (RR = 72.709; 95%CI: 54.487 to 90.931; $P < 0.001$). ESD group required more additional surgical operations relative to that of EMR group (RR = 3.139; 95%CI: 1.360 to 7.243; $P = 0.007$).

Compared to EMR, ESD shows a significant increased risk of perforation (RR = 7.597; 95%CI: 4.281 to 13.479; $P < 0.001$) ([Figure 2](#)), but no significant difference in the bleeding risk was observed between the two groups (RR = 1.277; 95%CI: 0.896 to 1.820; $P = 0.175$) ([Figure 3](#)). The rate of recurrence in the ESD group was lower than that of the EMR group (RR = 0.269; 95%CI: 0.112 to 0.648; $P = 0.003$) ([Figure 4](#)).

Meta regression

The influence of the included covariates on *en bloc* resection, complete resection, risk of perforation and bleeding, and rate of recurrence evaluated by meta-regression are summarized in [Table 2](#). Meta-regression analysis for risk of perforation suggested that right colonic polyps ($\beta = 7.731$; 95%CI: 4.965 to 10.497; $P < 0.001$) correlated with an increased risk in perforation in ESD ([Figure 5](#)). Other factors including polyp size and depth of invasion, did not influence the *en bloc* resection rate, complete resection rate, the risk of bleeding, and recurrence rate.

Sensitivity analysis for articles ≥ 20 mm

The results of the sensitivity analysis are summarized in [Supplementary Table 2](#) and results were largely unchanged after a sensitivity analysis. When a sensitivity analysis was performed to include ≥ 20 mm colorectal polyps only, the pooled analysis revealed higher rate of *en bloc* resection (RR = 1.932; 95%CI: 1.389 to 2.688; $P < 0.001$), longer operation time (RR = 3.247; 95%CI: 59.249 to 87.245; $P < 0.001$), higher risk of perforation (RR = 4.513; 95%CI: 2.531 to 8.046; $P < 0.001$) and lower recurrence rate (RR = 0.191; 95%CI: 0.085 to 0.431; $P < 0.001$) in ESD compared to EMR groups. Furthermore, in the two studies included, more additional surgical operations were required (RR = 3.139; 95%CI: 1.360 to 7.243; $P = 0.007$) in the ESD than in the EMR groups.

Subgroup analysis by region

The results of the subgroup analysis comparing outcomes of ESD with EMR in Japan vs the rest of the world are presented in [Table 3](#).

ESD in Japan had an *en bloc* resection rate of 89% (95%CI: 0.77 to 0.97), perforation risk of 4% (95%CI: 0.01 to 0.07), bleeding risk of 2.4% (95%CI: 0.01 to 0.04) and recurrence rate of 1% (95%CI: 0.01 to 0.02) while its EMR had an *en bloc* resection rate of 53% (95%CI: 0.38 to 0.67), perforation risk of 0.0002% (95%CI: 0.00 to 0.00081), bleeding risk of 2.1% (95%CI: 0.01 to 0.03) and recurrence rate of 7% (95%CI: 0.02 to 0.15). Comparing between the two techniques in Japan, ESD had higher *en bloc* resection rate (RR = 1.658, 95%CI: 1.270 to 2.165, $P < 0.001$), perforation risk (RR = 9.586, 95%CI: 4.425 to 20.768, $P < 0.001$) and bleeding risk (RR = 1.267, 95%CI: 1.174 to

Table 1 Pooled proportions and comparative meta-analysis of endoscopic submucosal dissection and endoscopic mucosal resection

	Total papers	Sample size (ESD)	Pooled proportions	Sample size (EMR)	Pooled proportions	RR (CI)	P value	Publication bias
<i>En bloc</i> resection	11	1641	89% (0.83-0.94)	1411	47% (0.36-0.59)	1.837 (1.464-2.305)	< 0.001	0.0025
Positive lateral margin	2	123	3% (0.01-0.06)	187	14% (0.09-0.19)	0.292 (0.089-0.995)	0.042	-
Positive vertical margin	1	38	5% (0.00-0.17)	83	1% (0.00-0.07)	4.368 (0.409-46.710)	0.223	-
Complete resection	8	918	82% (0.74-0.88)	1012	56% (0.34-0.77)	1.504 (1.041-2.174)	0.03	-
Lymphovascular invasion	1	54	6% (0.03-0.13)	23	0% (0.00-0.04)	4.352 (0.248-76.483)	0.315	-
Mean procedural time	8	1087	-	838	-	72.709 (54.487-90.931)	< 0.001	-
Additional surgery	2	99	13% (0.07-0.21)	153	5% (0.02-0.09)	3.139 (1.360-7.243)	0.007	-
Perforation	18	19470	5% (0.03-0.09)	260901	0% (0.00-0.01)	7.597 (4.281-13.479)	< 0.001	0.301
Bleeding	14	20048	3% (0.02-0.05)	257065	3% (0.02-0.04)	1.277 (0.896-1.820)	0.175	0.139
Recurrences	12	1822	2% (0.01-0.03)	37721	10% (0.04-0.17)	0.269 (0.112-0.648)	0.003	0.725

ESD: Endoscopic submucosal dissection; EMR: Endoscopic mucosal resection; CI: Confidence interval.

1.367, $P < 0.001$) than EMR. Following sensitivity analysis, similar results were obtained, except ESD had lower recurrence rate than EMR (RR = 0.204, 95%CI: 0.097 to 0.429, $P < 0.001$), and there was no difference in bleeding risk (RR = 0.895, 95%CI: 0.438 to 1.829, $P = 0.762$) between the two techniques.

With regards to studies from the rest of the world, ESD had an *en bloc* resection rate of 90% (95%CI: 0.85 to 0.94), perforation risk of 8% (95%CI: 0.05 to 0.12), bleeding risk of 5% (95%CI: 0.02 to 0.11) and recurrence rate of 3% (95%CI: 0.00 to 0.08) while EMR had an *en bloc* resection rate of 41% (95%CI: 0.22 to 0.61), perforation risk of 1% (95%CI: 0.00 to 0.02), bleeding risk of 3% (95%CI: 0.01 to 0.06) and recurrence rate of 16% (95%CI: 0.04 to 0.32). Comparing between the two techniques, ESD had higher *en bloc* resection rate (RR = 2.201, 95%CI: 1.411 to 3.433, $P = 0.001$), perforation rate (RR = 4.602, 95%CI: 2.729 to 7.759, $P < 0.001$) and lower recurrence rate (RR = 0.245, 95%CI: 0.073 to 0.819, $P = 0.022$) than EMR. Following sensitivity analysis, similar results were obtained.

DISCUSSION

The advancements in endoscopic resection techniques have resulted in the shift from radical surgery to minimally invasive and organ-sparing endoscopic resection techniques, such as ESD and EMR, for the treatment of colorectal lesions. With reference to 2015 ESGE guidelines, ESD should be considered for colorectal lesions larger than 20 mm, with high suspicion of submucosal invasion, or those where *en bloc* resection by EMR are not feasible[5]. Previous meta-analyses comparing ESD and EMR for colorectal polyps primarily reported data from Asian countries, with 72.7% of the published studies from Japan[6]. Since then, several retrospective and prospective studies comparing ESD and EMR for the treatment of colorectal polyps have been published outside of Japan. The present study is the most extensive meta-analysis evaluating the surgical, histological, and oncological outcomes of ESD in comparison to EMR in the treatment of colorectal polyps. Nine out of 21 studies (42.9%) on colorectal polyps included in this meta-analysis were conducted in countries outside of Japan. While ESD has been known to provide significantly better resection outcomes and lower recurrence rate, our analysis found that polyp size and depth of invasion did not significantly influence the *en bloc* and complete resection rate,

Table 2 Meta-regression

	En bloc resection			Complete resection			Perforation		Bleeding		Recurrence				
	n	Beta (CI)	P value	n	Beta (CI)	P value	n	Beta (CI)	P value	n	Beta (CI)	P value	n	Beta (CI)	P value
Age	8	-0.017 (-0.168-0.133)	0.785	5	-0.010 (-0.426-0.406)	0.944	9	0.073 (-0.191-0.336)	0.535	8	-0.201 (-0.540-0.138)	0.197	8	0.254 (-0.039-0.547)	0.078
Gender	6	-6.023 (-23.152-11.106)	0.384	4	-20.662 (-108.758-67.434)	0.419	8	-7.868 (-22.078-6.342)	0.224	7	-4.446 (-27.915-19.022)	0.647	6	-9.096 (-62.174-43.982)	0.659
Polyp size	10	0.242 (-0.037-0.086)	0.390	6	-0.034 (-0.098-0.030)	0.217	10	-0.034 (-0.117-0.049)	0.378	9	-0.041 (-0.177-0.095)	0.498	8	0.077 (-0.118-0.272)	0.369
Lateral spreading tumor	10	1.262 (-1.072-3.596)	0.248	7	-0.481 (-4.655-3.693)	0.779	11	-2.324 (-5.536-0.888)	0.136	9	-0.680 (-5.586-4.226)	0.753	9	3.827 (-2.323-9.977)	0.185
Right colon	8	1.247 (-2.024-4.517)	0.387	5	2.373 (-0.881-5.626)	0.103	8	7.731 (4.965-10.497)	< 0.001	7	1.688 (-12.442-15.817)	0.771	6	-8.739 (-26.616-9.139)	0.246
Colon	7	-1.803 (-4.486-0.880)	0.145	4	-1.914 (-7.213-3.385)	0.26	8	0.759 (-3.184-4.701)	0.654	6	6.649 (-5.182-18.480)	0.194	7	-1.102 (-4.754-2.551)	0.473
Rectum	7	1.803 (-0.880-4.486)	0.145	4	1.914 (-3.385-7.213)	0.26	8	-0.759 (-4.701-3.184)	0.654	6	-6.649 (-18.480-5.182)	0.194	7	1.102 (-2.551-4.754)	0.473
Submucosal invasion	8	-0.004 (-0.014-0.007)	0.431	5	-0.006 (-0.020-0.008)	0.253	9	0.001 (-0.049-0.051)	0.956	8	-0.016 (-0.033-0.001)	0.061	7	-6.508 (-27.976-14.960)	0.471
Muscularis propria invasion	8	-5.870 (-17.738-5.998)	0.272	5	-0.157 (-21.715-21.401)	0.983	9	6.731 (-33.772-47.234)	0.706	8	-109.836 (-1040.943-821.272)	0.783	-	-	-

CI: Confidence interval.

bleeding and perforation risk, and recurrence rate in colorectal polyps that was not previously reported. Additionally, previous reviews were confined only to sessile lesions larger than 20 mm[6,7]. Our analysis also showed the increased proficiency in performing ESD and EMR in Japan as compared to the rest of the world.

Consistent with previous studies, ESD showed benefits in the technical, histological, and oncological outcomes. Pooled analysis showed higher rates of *en bloc* resection and complete resection in ESD than in EMR albeit significant publication bias ($P = 0.0025$). *En bloc* resection offers the technical advantage of removing the entire pathologic specimen, thus allowing for detailed histologic evaluation. This results in an increase in the complete resection rate which in turn reduces the recurrence rate. Therefore, *en bloc* resection with ESD is favored as it provides curative treatment without the need for surgery for lesions with significant likelihood of submucosal invasion[38]. However, the advantages of ESD come at the expense of longer procedural time, additional surgical operations, and increased perforation risk compared to EMR[6]. The high rate of additional surgical operations for ESD is presumed to be due to the aggressive selection of ESD for T1 Lesions. Although the perforation risk was higher with ESD, most perforations in the studies included were treated conservatively or endoscopically using endoclips[9,12,31,33,34,37].

Meta-regression was performed to assess the risk factors that affect surgical, histological, and oncological outcomes. Our present analysis showed that polyp size did not affect the risk of perforation, which was reported otherwise in studies by Kim

Table 3 Subgroup analysis (Japan and rest of the world) of pooled proportions and comparative meta-analysis of endoscopic submucosal dissection and endoscopic mucosal resection

	Japan papers	Pooled proportions (ESD)	Pooled proportions (EMR)	RR (CI)	P value	Rest of the world papers	Pooled proportions (ESD)	Pooled proportions (EMR)	RR (CI)	P value
Overall results										
<i>En bloc</i> resection	6	89% (0.77-0.97)	53% (0.38-0.67)	1.658 (1.270-2.165)	< 0.001	5	90% (0.85-0.94)	41% (0.22-0.61)	2.201 (1.411-3.433)	0.001
Positive lateral margin	2	3% (0.00-0.06)	14% (0.09-0.19)	0.292 (0.089-0.955)	0.042	-	-	-	-	-
Positive vertical margin	1	5% (0.01-0.17)	1% (0.00-0.05)	4.368 (0.409-46.710)	0.223	-	-	-	-	-
Complete resection	2	79% (0.73-0.84)	53% (0.48-0.58)	1.452 (1.303-1.618)	< 0.001	6	85% (0.78-0.91)	59% (0.27-0.88)	1.562 (0.921-2.650)	0.098
Lymphovascular invasion	1	6% (0.03-0.13)	0% (0.00-0.04)	4.352 (0.248-76.483)	0.315	-	-	-	-	-
Mean procedural time	4	-	-	72.106 (48.831-95.382)	< 0.001	4	-	-	73.916 (36.075-111.757)	< 0.001
Additional surgery	2	13% (0.07-0.21)	5% (0.02-0.09)	3.139 (1.360-7.243)	0.007	-	-	-	-	-
Perforation	11	4% (0.01-0.07)	0.0002% (0.00-0.00081)	9.586 (4.425-20.768)	< 0.001	7	8% (0.05-0.12)	1% (0.00-0.02)	4.602 (2.729-7.759)	< 0.001
Bleeding	8	2.4% (0.01-0.04)	2.1% (0.01-0.03)	1.267 (1.174-1.367)	< 0.001	6	5% (0.02-0.11)	3% (0.01-0.06)	1.986 (0.716-5.508)	0.188
Recurrences	7	1% (0.01-0.02)	7% (0.02-0.15)	0.274 (0.071-1.054)	0.06	5	3% (0.00-0.08)	16% (0.04-0.32)	0.245 (0.073-0.819)	0.022
Sensitivity analysis (≥ 20 mm only)										
<i>En bloc</i> resection	4	82% (0.72-0.91)	50% (0.33-0.67)	1.645 (1.174-2.306)	0.004	2	91% (0.88-0.93)	33% (0.28-0.37)	2.668 (1.752-4.063)	< 0.001
Positive lateral margin	2	3% (0.00-0.06)	14% (0.09-0.19)	0.292 (0.089-0.955)	0.042	-	-	-	-	-
Positive vertical margin	1	5% (0.01-0.17)	1% (0.00-0.07)	4.368 (0.409-46.710)	0.223	-	-	-	-	-
Complete resection	2	79% (0.73-0.84)	53% (0.48-0.58)	1.452 (1.303-1.618)	< 0.001	2	90% (0.87-0.93)	77% (0.73-0.82)	1.613 (0.209-12.452)	0.647
Lymphovascular invasion	2	6% (0.03-0.13)	0% (0.00-0.04)	4.352 (0.248-76.483)	0.315	-	-	-	-	-
Mean procedural time	2	-	-	69.167 (48.446-89.888)	< 0.001	1	-	-	82.700 (67.578-97.822)	< 0.001
Additional surgery	2	13% (0.07-0.21)	5% (0.02-0.09)	3.139 (1.360-7.243)	0.007	-	-	-	-	-
Perforation	5	4% (0.01-0.07)	0% (0.00-0.01)	5.235 (2.123-12.910)	< 0.001	3	7% (0.05-0.10)	1% (0.00-0.03)	4.546 (1.674-12.346)	0.003

Bleeding	4	3% (0.00-0.07)	3% (0.01-0.05)	0.895 (0.438- 1.829)	0.762	3	3% (0.00-0.08)	1% (0.00-0.04)	2.233 (0.489- 10.197)	0.300
Recurrences	5	1% (0.00-0.02)	7% (0.02-0.15)	0.204 (0.097- 0.429)	< 0.001	3	3% (0.00-0.10)	27% (0.10-0.20)	0.179 (0.032- 0.990)	0.049

ESD: Endoscopic submucosal dissection; EMR: Endoscopic mucosal resection; RR: Relative risk; CI: Confidence interval.




Figure 1 PRISMA flow diagram of included articles. ESD: Endoscopic submucosal dissection; EMR: Endoscopic mucosal resection.

et al[39] and Hong *et al*[40] Polyp size and depth of invasion were also not associated with significant change in *en bloc* and complete resection rate, risk of bleeding, and recurrence between ESD and EMR. Furthermore, the *en bloc* resection rate (before RR = 1.837, 95%CI: 1.464 to 2.305, $P < 0.001$; after RR = 1.932, 95%CI: 1.389 to 2.688, $P < 0.001$) and recurrence rate (before RR = 0.269, 95%CI: 0.112 to 0.648, $P = 0.003$; after RR = 0.191, 95%CI: 0.085 to 0.431; $P < 0.001$) appeared comparable before and after the sensitivity analysis to ≥ 20 mm polyps. Instead, the risk of perforation was increased in patients with right colonic polyps and this was consistent with previous study which identified the technical difficulty and proficient endoscopic skills required to remove polyps from right colon safely[41]. As such, training should ensure endoscopists achieve procedural proficiency in left sided lesions before proceeding to attempt right sided lesions. Other factors including polyp size and depth of invasion are less important criteria when deciding between EMR and ESD in skilled tertiary centers.

Despite the advantages of ESD, the data regarding the efficacy of colorectal ESD have been inconsistent and vary between Japan and the rest of the world. One of the reasons is the limitations to the implementation of ESD in other countries, which are in part due to the lack of expertise and training centers. To date, no meta-analysis comparing ESD and EMR between Japan and the rest of the world have been performed. Our single arm meta-analysis found that Japan performed better than the rest of the world in ESD and EMR. Significantly, perforation is a major concern in ESD. The perforation risk of ESD and EMR was 4% and 0.0002%, respectively in Japan, and 8% and 1%, respectively in the rest of the world. While there is an observed increase risk of perforation from Japan only studies (RR = 9.586) compared to the rest of the world (RR = 4.602), even after sensitivity analysis, the risk of perforation for ESD was only statistically higher in Japan only studies due to the very low risk of perforation in EMR in Japan. In addition, the challenges of doing ESD in difficult could have resulted in the higher perforation rate in ESD compared with EMR in Japan. Another important factor to consider is the recurrence rate of ESD and EMR which was 1% and 7%,




Figure 2 Forest plot for risk of perforation. RR: Relative risk; CI: Confidence interval.

respectively in Japan and 3% and 27%, respectively in the rest of the world following sensitivity analysis. The observed increase in recurrence rate from Japan only studies (RR = 0.204) compared to the rest of the world (RR = 0.179) was due to the much lower recurrence rate of EMR in Japan as compared to the rest of the world. Overall, our results seem to favor studies from Japan and are in tandem with a single arm analysis of ESD only procedures with subgroup on region efficacy[42]. However, most of the studies originating from the rest of the world should be interpreted with the understanding that these studies are mainly from tertiary centers and the results may not be generalized to non-tertiary centers.

The potential of ESD resection is limited by the difficulty in conducting the procedure as the length of procedure for ESD even when performed by experienced endoscopists can be three times longer than that for EMR[43]. However, the advancement in endoscopic resection equipment has been shown to shorten the procedure time despite a relatively short training duration[44]. Using the cumulative sum method, Miyakawa *et al*[44] recently reported the use of Stag-Beetle Knife Jr for ESD in a Japan single-center study generated good learning curve to achieve satisfactory resection speed (min/cm²), which allowed the acquisition of proficient and safe skills within 120 cases[44]. Other alternatives to ESD do exist, such as hybrid ESD and pre-cut EMR. This hybrid approach has been shown to have lower *en bloc* resection rate (68.4% *vs* 91.0%) and complete resection rate (60.6% *vs* 82.9%) than conventional ESD[42]. Currently, underwater EMR has been thought to be a safe and effective method with higher rate of *en bloc* resection and lower rate of recurrence[45], but no head-to-head comparisons have been done between UMER and ESD.

The inclusion of 21 studies with a total of 281344 polyps based on our search strategy and inclusion criteria represents the most extensive meta-analysis on this issue. However, as no randomized controlled trials comparing the performance between EMR and ESD have yet been conducted, this highlights the need for a randomized study to better understand the efficacy and safety of these techniques in the management of colorectal polyps. The evaluation of heterogeneity allowed us to compare the significant differences in the performance of ESD and EMR between




Figure 3 Forest plot for risk of bleeding. RR: Relative risk; CI: Confidence interval.

Japan and the rest of the world.

Our meta-analysis has some limitations. While we aimed to decrease heterogeneity, the included articles used a variety of EMR techniques including standard EMR, piecemeal EMR, EMR with small incision, EMR-precutting and EMR-circumferential incision. This, however, was an acceptable confounding factor in previous meta-analysis analysis. Also, a major factor that we were unable to regress for was the procedural skills of each centers. ESD and EMR are largely operator dependent and we were only able to account for it in a subgroup analysis comparing between studies conducted in Japan and the rest of the world.

CONCLUSION

Evidence from this meta-analysis suggests that with appropriate training, ESD is preferred over EMR as the first-line therapy for resection of colorectal polyps, without restricting to lesions greater than 20 mm and those with high suspicion of submucosal invasion. Our overall findings are consistent with previous meta-analyses showing ESD is associated with higher rate of *en bloc* and complete resection, and lower recurrence compared to EMR, but at the cost of increased procedural time, need for additional surgical operations and perforation risk. This is coupled with the new finding that confounders including polyp size and invasion depth did not influence the rates of *en bloc* and complete resection, bleeding risk, and recurrence.




Figure 4 Forest plot for recurrence rate. RR: Relative risk; CI: Confidence interval.




Figure 5 Meta regression of perforation and proportion of right colon polyps. CI: Confidence interval.

ARTICLE HIGHLIGHTS

Research background

Endoscopic submucosal dissection (ESD) has shown to be effective in management of colorectal neoplasm in the Asian countries, while its implementation in Western countries where endoscopic mucosal resection (EMR) is preferred is still debatable.

Research motivation

Previous meta-analyses and systematic reviews comparing ESD with EMR included studies mainly conducted in Asia, with limited data coming from outside of Japan and did not control for confounders or lesions smaller than 20 mm.

Research objectives

To compare the outcomes of ESD and EMR in the treatment of colorectal polyps, with subgroup analysis comparing the efficacy of these two techniques between Japan and the rest of the world.

Research methods

Embase and Medline databases were searched in accordance with PRISMA guidelines for studies comparing *en bloc*, complete resection, margin involvement, resection time, need for additional surgery, complications, and recurrence rate of ESD with EMR in patients with colorectal lesions.

Research results

ESD was associated with better resection outcomes and lower recurrence rate when compared to EMR. Meta-regression analysis suggested only right colonic polyps correlated with an increased perforation risk in ESD, while confounders including polyp size and invasion depth did not significantly influence the resection outcomes, bleeding risk and recurrence. Subgroup analysis showed that Japan performed better than the rest of the world in both ESD and EMR with lower perforation risk.

Research conclusions

This meta-analysis suggests that with appropriate training, ESD is preferred over EMR as the first-line therapy for resection of colorectal polyps, without restricting to lesions greater than 20 mm and those with high suspicion of submucosal invasion. Increased proficiency in performing ESD and EMR was shown in Japan as compared with the rest of the world.

Research perspectives

This highlights the need to establish adequate training programs for colorectal ESD to be performed effectively. A randomized controlled trial is necessary to better understand the efficacy and safety of these techniques in the management of colorectal polyps.

ACKNOWLEDGEMENTS

We would like to thank Dr. Hiroaki Ikematsu of the Division of Science and Technology for Endoscopy, Exploratory Oncology Research & Clinical Trial Center, National Cancer Center for his advice on the writing of the manuscript.

REFERENCES

- 1 Winawer SJ, Zauber AG, Ho MN, O'Brien MJ, Gottlieb LS, Sternberg SS, Wayne JD, Schapiro M, Bond JH, Panish JF. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyp Study Workgroup. *N Engl J Med* 1993; **329**: 1977-1981 [PMID: 8247072 DOI: 10.1056/NEJM199312303292701]
- 2 Zauber AG, Winawer SJ, O'Brien MJ, Lansdorp-Vogelaar I, van Ballegooijen M, Hankey BF, Shi W, Bond JH, Schapiro M, Panish JF, Stewart ET, Wayne JD. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med* 2012; **366**: 687-696 [PMID: 22356322 DOI: 10.1056/NEJMoa1100370]
- 3 Siegel RL, Miller KD, Goding Sauer A, Fedewa SA, Butterly LF, Anderson JC, Cercek A, Smith RA,

- Jemal A. Colorectal cancer statistics, 2020. *CA Cancer J Clin* 2020; **70**: 145-164 [PMID: [32133645](#) DOI: [10.3322/caac.21601](#)]
- 4 Yamamoto H, Yube T, Isoda N, Sato Y, Sekine Y, Higashizawa T, Ido K, Kimura K, Kanai N. A novel method of endoscopic mucosal resection using sodium hyaluronate. *Gastrointest Endosc* 1999; **50**: 251-256 [PMID: [10425422](#) DOI: [10.1016/s0016-5107\(99\)70234-8](#)]
- 5 Pimentel-Nunes P, Dinis-Ribeiro M, Ponchon T, Repici A, Vieth M, De Ceglie A, Amato A, Berr F, Bhandari P, Bialek A, Conio M, Haringsma J, Langner C, Meisner S, Messmann H, Morino M, Neuhaus H, Piessevaux H, Rugge M, Saunders BP, Robaszkiewicz M, Seewald S, Kashin S, Dumonceau JM, Hassan C, Deprez PH. Endoscopic submucosal dissection: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 2015; **47**: 829-854 [PMID: [26317585](#) DOI: [10.1055/s-0034-1392882](#)]
- 6 Arezzo A, Passera R, Marchese N, Galloro G, Manta R, Cirocchi R. Systematic review and meta-analysis of endoscopic submucosal dissection vs endoscopic mucosal resection for colorectal lesions. *United European Gastroenterol J* 2016; **4**: 18-29 [PMID: [26966519](#) DOI: [10.1177/2050640615585470](#)]
- 7 De Ceglie A, Hassan C, Mangiavillano B, Matsuda T, Saito Y, Ridola L, Bhandari P, Boeri F, Conio M. Endoscopic mucosal resection and endoscopic submucosal dissection for colorectal lesions: A systematic review. *Crit Rev Oncol Hematol* 2016; **104**: 138-155 [PMID: [27370173](#) DOI: [10.1016/j.critrevonc.2016.06.008](#)]
- 8 Lee EJ, Lee JB, Lee SH, Youk EG. Endoscopic treatment of large colorectal tumors: comparison of endoscopic mucosal resection, endoscopic mucosal resection-precutting, and endoscopic submucosal dissection. *Surg Endosc* 2012; **26**: 2220-2230 [PMID: [22278105](#) DOI: [10.1007/s00464-012-2164-0](#)]
- 9 Kim YJ, Kim ES, Cho KB, Park KS, Jang BK, Chung WJ, Hwang JS. Comparison of clinical outcomes among different endoscopic resection methods for treating colorectal neoplasia. *Dig Dis Sci* 2013; **58**: 1727-1736 [PMID: [23385636](#) DOI: [10.1007/s10620-013-2560-x](#)]
- 10 Acquistapace F, Maternini F, Snider L, Bellini O, Moglia P, Capretti P. Endoscopic treatment of superficial colorectal neoplasms. Retrospective analysis of a single center technique and results. *G Chir* 2015; **36**: 247-250 [PMID: [26888699](#) DOI: [10.11138/gchir/2015.36.6.247](#)]
- 11 Emmanuel A, Gulati S, Burt M, Hayee B, Haji A. Combining eastern and western practices for safe and effective endoscopic resection of large complex colorectal lesions. *Eur J Gastroenterol Hepatol* 2018; **30**: 506-513 [PMID: [29406437](#) DOI: [10.1097/MEG.0000000000001086](#)]
- 12 Soliman H, Briau B, Guillaumot MA, Leblanc S, Barret M, Camus M, Dior M, Terris B, Coriat R, Prat F, Chaussade S. Invasive pit pattern, macronodule and depression are predictive factors of submucosal invasion in colorectal laterally spreading tumours from a Western population. *United European Gastroenterol J* 2018; **6**: 1569-1577 [PMID: [30574328](#) DOI: [10.1177/2050640618804713](#)]
- 13 Urban O, Falt P, Fojtik P, Andelova R. Comparison of endoscopic mucosal resection and endoscopic submucosal dissection in the treatment of flat neoplastic lesions in the rectum. *Gastroenterol Hepatol* 2018; **72**: 193-198 [DOI: [10.14735/amgh2018193](#)]
- 14 Jung JS, Hong JY, Oh HH, Kweon SS, Lee J, Kim SW, Seo GS, Kim HS, Joo YE. Clinical outcomes of endoscopic resection for colorectal laterally spreading tumors with advanced histology. *Surg Endosc* 2019; **33**: 2562-2571 [PMID: [30350100](#) DOI: [10.1007/s00464-018-6550-0](#)]
- 15 Liu C, Wu R, Sun X, Tao C, Liu Z. Risk factors for delayed hemorrhage after colonoscopic polypectomy: Polyp size and operative modality. *JGH Open* 2019; **3**: 61-64 [PMID: [30834342](#) DOI: [10.1002/jgh3.12106](#)]
- 16 Benites Goñi H, Palacios Salas F, Marin Calderón L, Bardalez Cruz P, Vásquez Quiroga J, Alva Alva E, Calixto Aguilar L, Alferez Andía J, Dávalos Moscol M. Closure of colonic deep mural injury and perforation with endoclips. *Rev Esp Enferm Dig* 2020; **112**: 772-777 [PMID: [32954773](#) DOI: [10.17235/reed.2020.6880/2020](#)]
- 17 Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097 [PMID: [19621072](#) DOI: [10.1371/journal.pmed.1000097](#)]
- 18 Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; **14**: 135 [PMID: [25524443](#) DOI: [10.1186/1471-2288-14-135](#)]
- 19 Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Arch Public Health* 2014; **72**: 39 [PMID: [25810908](#) DOI: [10.1186/2049-3258-72-39](#)]
- 20 Rousseau MJ, Evans JC. Key statistical assumptions and methods in one-arm meta-analyses with binary endpoints and low event rates, including a real-life example in the area of endoscopic colonic stenting. *Cogent Med* 2017; **4**: 1334318 [DOI: [10.1080/2331205X.2017.1334318](#)]
- 21 Harris R, Bradburn M, Deeks J, Harbord R, Altman D, Sterne J. Metan: Fixed- and random-effects meta-analysis. *Stata J* 2008; **8**: 3-28 [DOI: [10.1177/1536867X0800800102](#)]
- 22 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**: 177-188 [PMID: [3802833](#) DOI: [10.1016/0197-2456\(86\)90046-2](#)]
- 23 Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629-634 [PMID: [9310563](#) DOI: [10.1136/bmj.315.7109.629](#)]
- 24 Harbord RM, Higgins JPT. Meta-regression in stata. *Stata J* 2008; **8**: 493-519 [DOI: [10.1177/1536867X0800800403](#)]
- 25 Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses, 2012. [cited 10

- January 2021]. Available from: http://www.ohri.ca/Programs/clinical_epidemiology/oxford.asp
- 26 **Taku K**, Sano Y, Fu KI, Saito Y, Matsuda T, Uraoka T, Yoshino T, Yamaguchi Y, Fujita M, Hattori S, Ishikawa T, Saito D, Fujii T, Kaneko E, Yoshida S. Iatrogenic perforation associated with therapeutic colonoscopy: a multicenter study in Japan. *J Gastroenterol Hepatol* 2007; **22**: 1409-1414 [PMID: [17593224](#) DOI: [10.1111/j.1440-1746.2007.05022.x](#)]
 - 27 **Tamegai Y**, Saito Y, Masaki N, Hinohara C, Oshima T, Kogure E, Liu Y, Uemura N, Saito K. Endoscopic submucosal dissection: a safe technique for colorectal tumors. *Endoscopy* 2007; **39**: 418-422 [PMID: [17516348](#) DOI: [10.1055/s-2007-966427](#)]
 - 28 **Iizuka H**, Okamura S, Onozato Y, Ishihara H, Kakizaki S, Mori M. Endoscopic submucosal dissection for colorectal tumors. *Gastroenterol Clin Biol* 2009; **33**: 1004-1011 [PMID: [19762190](#) DOI: [10.1016/j.gcb.2009.02.039](#)]
 - 29 **Toyonaga T**, Man-I M, Morita Y, Sanuki T, Yoshida M, Kutsumi H, Inokuchi H, Azuma T. The new resources of treatment for early stage colorectal tumors: EMR with small incision and simplified endoscopic submucosal dissection. *Dig Endosc* 2009; **21** Suppl 1: S31-S37 [PMID: [19691730](#) DOI: [10.1111/j.1443-1661.2009.00872.x](#)]
 - 30 **Oka S**, Tanaka S, Kanao H, Ishikawa H, Watanabe T, Igarashi M, Saito Y, Ikematsu H, Kobayashi K, Inoue Y, Yahagi N, Tsuda S, Simizu S, Iishi H, Yamano H, Kudo SE, Tsuruta O, Tamura S, Cho E, Fujii T, Sano Y, Nakamura H, Sugihara K, Muto T. Current status in the occurrence of postoperative bleeding, perforation and residual/local recurrence during colonoscopic treatment in Japan. *Dig Endosc* 2010; **22**: 376-380 [PMID: [21175503](#) DOI: [10.1111/j.1443-1661.2010.01016.x](#)]
 - 31 **Saito Y**, Fukuzawa M, Matsuda T, Fukunaga S, Sakamoto T, Uraoka T, Nakajima T, Ikehara H, Fu KI, Itoi T, Fujii T. Clinical outcome of endoscopic submucosal dissection vs endoscopic mucosal resection of large colorectal tumors as determined by curative resection. *Surg Endosc* 2010; **24**: 343-352 [PMID: [19517168](#) DOI: [10.1007/s00464-009-0562-8](#)]
 - 32 **Tajika M**, Niwa Y, Bhatia V, Kondo S, Tanaka T, Mizuno N, Hara K, Hijioka S, Imaoka H, Ogura T, Haba S, Yamao K. Comparison of endoscopic submucosal dissection and endoscopic mucosal resection for large colorectal tumors. *Eur J Gastroenterol Hepatol* 2011; **23**: 1042-1049 [PMID: [21869682](#) DOI: [10.1097/MEG.0b013e32834aa47b](#)]
 - 33 **Kobayashi N**, Yoshitake N, Hirahara Y, Konishi J, Saito Y, Matsuda T, Ishikawa T, Sekiguchi R, Fujimori T. Matched case-control study comparing endoscopic submucosal dissection and endoscopic mucosal resection for colorectal tumors. *J Gastroenterol Hepatol* 2012; **27**: 728-733 [PMID: [22004124](#) DOI: [10.1111/j.1440-1746.2011.06942.x](#)]
 - 34 **Sagawa T**, Kakizaki S, Iizuka H, Onozato Y, Soharu N, Okamura S, Mori M. Analysis of colonoscopic perforations at a local clinic and a tertiary hospital. *World J Gastroenterol* 2012; **18**: 4898-4904 [PMID: [23002362](#) DOI: [10.3748/wjg.v18.i35.4898](#)]
 - 35 **Terasaki M**, Tanaka S, Oka S, Nakadoi K, Takata S, Kanao H, Yoshida S, Chayama K. Clinical outcomes of endoscopic submucosal dissection and endoscopic mucosal resection for laterally spreading tumors larger than 20 mm. *J Gastroenterol Hepatol* 2012; **27**: 734-740 [PMID: [22098630](#) DOI: [10.1111/j.1440-1746.2011.06977.x](#)]
 - 36 **Niikura R**, Yasunaga H, Yamada A, Matsui H, Fushimi K, Hirata Y, Koike K. Factors predicting adverse events associated with therapeutic colonoscopy for colorectal neoplasia: a retrospective nationwide study in Japan. *Gastrointest Endosc* 2016; **84**: 971-982.e6 [PMID: [27189658](#) DOI: [10.1016/j.gie.2016.05.013](#)]
 - 37 **Sato T**, Fukuzawa M, Gotoda T, Moriyasu F. Comparison of clinical outcomes between colorectal EMR and ESD. *J Tokyo Med Univer* 2016; **74**: 154-162
 - 38 **Tanaka S**, Kashida H, Saito Y, Yahagi N, Yamano H, Saito S, Hisabe T, Yao T, Watanabe M, Yoshida M, Kudo SE, Tsuruta O, Sugihara KI, Watanabe T, Saitoh Y, Igarashi M, Toyonaga T, Ajioka Y, Ichinose M, Matsui T, Sugita A, Sugano K, Fujimoto K, Tajiri H. JGES guidelines for colorectal endoscopic submucosal dissection/endoscopic mucosal resection. *Dig Endosc* 2015; **27**: 417-434 [PMID: [25652022](#) DOI: [10.1111/den.12456](#)]
 - 39 **Kim ES**, Cho KB, Park KS, Lee KI, Jang BK, Chung WJ, Hwang JS. Factors predictive of perforation during endoscopic submucosal dissection for the treatment of colorectal tumors. *Endoscopy* 2011; **43**: 573-578 [PMID: [21448852](#) DOI: [10.1055/s-0030-1256339](#)]
 - 40 **Hong SN**, Byeon JS, Lee BI, Yang DH, Kim J, Cho KB, Cho JW, Jang HJ, Jeon SW, Jung SA, Chang DK. Prediction model and risk score for perforation in patients undergoing colorectal endoscopic submucosal dissection. *Gastrointest Endosc* 2016; **84**: 98-108 [PMID: [26708921](#) DOI: [10.1016/j.gie.2015.12.011](#)]
 - 41 **Nishiyama H**, Isomoto H, Yamaguchi N, Fukuda E, Ikeda K, Ohnita K, Mizuta Y, Nakamura T, Nakao K, Kohno S, Shikuwa S. Endoscopic submucosal dissection for colorectal epithelial neoplasms. *Dis Colon Rectum* 2010; **53**: 161-168 [PMID: [20087091](#) DOI: [10.1007/DCR.0b013e3281b78cb6](#)]
 - 42 **Fuccio L**, Hassan C, Ponchon T, Mandolesi D, Farioli A, Cucchetti A, Frazzoni L, Bhandari P, Bellisario C, Bazzoli F, Repici A. Clinical outcomes after endoscopic submucosal dissection for colorectal neoplasia: a systematic review and meta-analysis. *Gastrointest Endosc* 2017; **86**: 74-86.e17 [PMID: [28254526](#) DOI: [10.1016/j.gie.2017.02.024](#)]
 - 43 **Fujiya M**, Tanaka K, Dokoshi T, Tominaga M, Ueno N, Inaba Y, Ito T, Moriichi K, Kohgo Y. Efficacy and adverse events of EMR and endoscopic submucosal dissection for the treatment of colon neoplasms: a meta-analysis of studies comparing EMR and endoscopic submucosal dissection. *Gastrointest Endosc* 2015; **81**: 583-595 [PMID: [25592748](#) DOI: [10.1016/j.gie.2014.07.034](#)]

- 44 **Miyakawa A**, Kuwai T, Sakuma Y, Kubota M, Nakamura A, Itobayashi E, Shimura H, Suzuki Y, Shimura K. Learning curve for endoscopic submucosal dissection of early colorectal neoplasms with a monopolar scissor-type knife: use of the cumulative sum method. *Scand J Gastroenterol* 2020; **55**: 1234-1242 [PMID: [32853052](#) DOI: [10.1080/00365521.2020.1807597](#)]
- 45 **Choi AY**, Moosvi Z, Shah S, Roccato MK, Wang AY, Hamerski CM, Samarasena JB. Underwater vs conventional EMR for colorectal polyps: systematic review and meta-analysis. *Gastrointest Endosc* 2021; **93**: 378-389 [PMID: [33068608](#) DOI: [10.1016/j.gie.2020.10.009](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

