



Match Overview

1	Crossref 122 words Beyann Alzoubi, Abish Kharel, Fauzia Osman, Fahad Aziz et al. "Incidence, risk factors, and outcomes of post-transpl...	3%
2	Internet 37 words crawled on 22-Jul-2020 www.tandfonline.com	1%
3	Internet 30 words crawled on 20-Sep-2019 onlinelibrary.wiley.com	1%
4	Internet 14 words crawled on 28-Nov-2016 pdfs.semanticscholar.org	<1%
5	Crossref 13 words Demetrios V. Vlahakos, Katerina P. Marathias, Basil Agroy... nnis, Nicolaos E. Madias. "Posttransplant erythrocytosis", Ki	<1%
6	Internet 12 words crawled on 22-Dec-2019 www.science.gov	<1%

Name of Journal: *World Journal of Transplantation*

Manuscript NO: 64973

Manuscript Type: MINIREVIEWS

Post-transplant erythrocytosis after kidney transplantation: A review

Alzoubi B *et al.* Post-transplant erythrocytosis

Beyann Alzoubi, Abish Kharel, Rushad Machhi, Fahad Aziz, Kurtis J Swanson, Sandesh Parajuli

Abstract

Post-Transplant Erythrocytosis: A Review



ALL

IMAGES

VIDEOS

310,000 Results

Any time

(PDF) Post-transplant erythrocytosis: a review of 47 renal ...

<https://www.researchgate.net/publication/6195393...>



Post-transplant erythrocytosis (PTE) is a well known phenomenon occurring in 5-17% of renal transplant recipients (RTR). In this retrospective study, we studied 47 RTR which included 39 males and...

[PDF] Postrenal transplant erythrocytosis: A review of 53 patients

[https://www.kidney-international.org/article/S0085-2538\(15\)32956-2/pdf](https://www.kidney-international.org/article/S0085-2538(15)32956-2/pdf)

Postrenal transplant erythrocytosis: A review of 53 patients. Review of a large renal transplant experience revealed a 17.3% incidence of post-transplant erythrocytosis. The influence of kidney sourc...

Related searches

- post transplant erythrocytosis **treatment**
- post transplant erythrocytosis **guidelines**
- work up for** erythrocytosis
- uptodate** erythrocytosis
- secondary** erythrocytosis **treatment list**
- erythrocytosis **differential diagnosis**

Search Tools

Turn off Hover Translation (关闭取词)

PEOPLE ALSO ASK



ALL

IMAGES

VIDEOS

203,000 Results

Any time ▾

Posttransplant erythrocytosis - Kidney International

[https://www.kidney-international.org/article/S0085-2538\(15\)48996-3/fulltext](https://www.kidney-international.org/article/S0085-2538(15)48996-3/fulltext)

Posttransplant erythrocytosis (PTE) is defined as a persistently elevated hematocrit to a level greater than 51% after renal transplantation. It occurs in 10% to 15% of graft recipients and usually develops 8 to 24 months after engraftment.

Posttransplant erythrocytosis - PubMed

<https://pubmed.ncbi.nlm.nih.gov/12631334>

Posttransplant erythrocytosis (PTE) is defined as a persistently elevated hematocrit to a level greater than 51% **after renal transplantation**. It occurs in 10% to 15% of graft recipients and usually develops 8 to 24 months after engraftment. Spontaneous remission of established **PTE** is observed in one fourth of the patients within 2 years from onset, whereas in the remaining three fourths it persists for several years, ...

[PDF] Posttransplant erythrocytosis - Kidney International

[https://www.kidney-international.org/article/S0085-2538\(15\)48996-3/pdf](https://www.kidney-international.org/article/S0085-2538(15)48996-3/pdf)

Posttransplant erythrocytosis (PTE) ropoietin secretion peaks within 3 days, reticulocytosis is defined as a persistently elevated hematocrit to a level greater than 51% **after renal transplantation**. It occurs in 10% to 15% of patients over 3 months. Thereafter, erythropoietin levels

Kidney transplantation in adults: Posttransplant ...

<https://www.uptodate.com/contents/kidney-transplantation-in-adults>

See results for

Polycythemia (Medical Condition)

A rare blood cancer in which bone marrow produces too many red blood cells. This causes itching, bruising, bloating, abdominal pain, fatigue, and weight loss.

Search Tools

Turn on Hover Translation (开启取词)

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

624,000 Results

Any time ▾

Posttransplant erythrocytosis (PTE) is defined as a persistently elevated hematocrit to a level greater than 51% after **renal transplantation**. It occurs in 10% to 15% of graft recipients and usually develops 8 to 24 months after engraftment.

Posttransplant erythrocytosis - Kidney International

[www.kidney-international.org/article/S0085-2538\(15\)48996-3/fulltext](http://www.kidney-international.org/article/S0085-2538(15)48996-3/fulltext)

Was this helpful?  

PEOPLE ALSO ASK

What is posttransplant erythrocytosis? ▾

Is erythrocytosis related to renal artery stenosis? ▾

Can a kidney transplant cause headache and dizziness? ▾

Can erythrocytosis cause polycythemia? ▾

Feedback

Posttransplant erythrocytosis - Kidney International

[https://www.kidney-international.org/article/S0085-2538\(15\)48996-3/fulltext](https://www.kidney-international.org/article/S0085-2538(15)48996-3/fulltext)

Posttransplant erythrocytosis (PTE) is defined as a persistently elevated hematocrit to a level greater than 51% after renal transplantation. It occurs in 10% to 15% of graft recipients and usually develops 8 to 24...

Posttransplant erythrocytosis - PubMed

<https://pubmed.ncbi.nlm.nih.gov/12631334>

Posttransplant erythrocytosis (PTE) is defined as a persistently elevated hematocrit to a level greater than 51% after **renal transplantation**. It occurs in 10% to 15% of graft recipients and usually develops 8 t...

[PDF] Post-Transplant Erythrocytosis - Proceedings

<https://proceedings.med.ucla.edu/wp-content/...>

See results for

Polycythemia (Medical Condition)

A rare blood cancer in which bone marrow produces too many red blood cells. This causes itching, bruising, bloating, abdominal pain, fatigue, hea...