

TO WHOM IT MAY CONCERN:

Lowered paraoxonase 1 activities may explain the comorbidities between temporal lobe epilepsy or mesial temporal sclerosis and depression, anxiety and psychosis.

This to confirm that the text has been revised by Prof. Dr. Michael Maes, M.D., Ph.D.

Prof. Dr. Michael Maes, M.D., Ph.D.

Impact Research Center

Department Psychiatry, Deakin University

Geelong, Vic, Australia

<https://scholar.google.co.th/citations?user=1wzMZ7UAAAAJ&hl=th&oi=ao>



Highly cited: 2020

Expert worldwide ranking:

#1 in CFS

#1 in oxidative stress

#1 in encephalomyelitis

#1 in nitrosative stress

#1 in tryptophan

#2 in stress, physiological

#3 in inflammation

#2 in neuroimmune

#7 in depression

#5 in fatigue

#7 in psychiatry

#11 in melancholia

Bibliometric Website Expertscape, objective ranking of medical expertise.

top 1 scientist in psychiatry and neurology in Asia-Australasia

top 3 scientist in psychiatry and neurology in Europe

top 22 scientist worldwide in psychiatry and neurology

top 0.01% of all scientists worldwide

standardized citation data annotated for multiple impact indicators