

[Screening for sepsis in general hospitalized patients: a ...](#)

<https://www.sciencedirect.com/science/article/pii/S019567011730275X>

Aug 01, 2017 · The search resulted in 14,771 citations retrieved from six search engines and manual searching. After eliminating duplicates, 8456 citations including titles and abstracts were screened for eligibility. Six articles met the inclusion criteria, including two **prospective observational** pilot studies, one **prospective observational study**, two pre-post studies and one retrospective cohort **study**

Cited by: 9 **Author:** L. Alberto, A.P. Marshall, A.P. Marsha...

Publish Year: 2017

[Noninvasive ventilation failure in patients with hypoxemic ...](#)

<https://pubmed.ncbi.nlm.nih.gov/31722614>

Background: **Sepsis** and septic shock are common in noninvasive ventilation (NIV) patients. However, studies on the association between **sepsis** and NIV failure are lacking. Methods: A **prospective** multi-center **observational study** was performed in 16 Chinese intensive care units (ICUs). Patients who used NIV due to hypoxemic respiratory failure were enrolled.

Cited by: 2 **Author:** Jun Duan, Lijuan Chen, Guopeng Lia...

Publish Year: 2019

PEOPLE ALSO ASK

- How is sepsis diagnosed in intensive care units? ▾
- What is the window between the onset and identification of sepsis? ▾
- Which is a useful predictor of outcome in sepsis? ▾
- When is a patient screen positive for severe sepsis? ▾

Feedback

[\(PDF\) Risk modifiers of acute respiratory distress ...](#)

<https://www.researchgate.net/publication/338525533...>

International normalized ratio to albumin ratio - IJGM | IJGM

[https://www.dovepress.com/international-normalized...](https://www.dovepress.com/international-normalized-ratio-to-albumin-ratio) ▾

Objective: Our **study** aimed to validate the usefulness of the prothrombin time–**international normalized ratio** (PT-INR) to albumin **ratio** (PTAR) in predicting the mortality of patients with **sepsis**. Methods: Data on a total of 4536 patients, obtained from the Multiparameter Intelligent Monitoring in Intensive Care III database, were included in ...

Author: Shaoxiong Wang, Shuizi Ding, Hong Luo... **Publish Year:** 2021

The effect of vitamin K on prothrombin time in critically ill patients ...

<https://jintensivecare.biomedcentral.com/articles/10.1186/s40560-020-00517-5> ▾

Jan 18, 2021 · Previous studies have indicated that vitamin K deficiency is common in non-bleeding critically ill patients with slightly prolonged prothrombin time–**international normalized ratio** (PT-INR). It has never been investigated thoroughly whether the administration of vitamin K to these patients could affect their PT-INR. Therefore, the aim of this registry **study** was to evaluate changes in PT-INR in ...

Author: Sofia Dahlberg, Ulf Schött, Thomas Kan... **Publish Year:** 2021

PEOPLE ALSO ASK

Why is the international normalised ratio raised after liver resection? ▾

What is the SOFA score for sepsis patients? ▾

What is the international consensus definition of sepsis? ▾


When to use FFP for a prolonged PT / INR? ▾

Feedback

Role of thromboelastography in the evaluation of septic shock ...

<https://www.nature.com/articles/s41598-021-91221-3>

Jun 04, 2021 · After excluding overt-disseminated intravascular coagulation (DIC) defined by scoring system, we included patients with a normal value of **international normalized ratio** ≤ 1.3 and aPTT ≤ 34 s.



Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 67488

Manuscript Type: ORIGINAL ARTICLE

Observational Study

The role of international normalised ratio (INR) in nonpulmonary sepsis screening: a prospective observational study

INR in nonpulmonary sepsis screening

Jing Zhang, Huimin Du, Mingxiang Cheng, Faming He, Bailin Niu

Match Overview

Match #	Source	Similarity
1	Internet 84 words Crawled on 21-Sep-2020 journals.lww.com	2%
2	Internet 30 words Crawled on 27-Jun-2018 www.sjgnet.com	1%
3	Internet 16 words www.ncbi.nlm.nih.gov	<1%
4	Internet 15 words Crawled on 18-Feb-2020 easgtrnmgagp.springeropen.com	<1%
5	Internet 14 words Crawled on 22-Mar-2019 www.sciencedirect.com	<1%
6	Internet 13 words Crawled on 09-Aug-2020 workwellscience.org	<1%
7	Internet 13 words Crawled on 09-Sep-2020 www.frontiersin.org	<1%
8	Crossref 12 words Q. Hsieh: "The effect of economic status on height, insulin-like growth factor (IGF-1) and IGF binding protein-3 cor..."	<1%
9	Crossref 11 words Nashida, T.A., "Influence of Toll Like Receptor-4, CD14, Tumor Necrosis Factor, and Interleukin-10 Gene Polymor..."	<1%

国内版

国际版

Role of international normalized ratio in nonpulmonary sepsis scre



ALL

IMAGES

VIDEOS

40,600 Results

Any time ▾

[An observational study of international normalized ratio ...](#)

<https://pubmed.ncbi.nlm.nih.gov/31774502>

Aims : In patients with non-valvular atrial fibrillation prescribed warfarin, the UK National Institute of Health and Care Excellence (NICE) defines poor anticoagulation as a time in therapeutic range (TTR) of <65%, any two **international normalized ratios** (INRs) within a 6-month period of ≤ 1.5 ('low'), two INRs ≥ 5 within 6 months, or any INR ≥ 8 ('high').

Cited by: 2

Author: Daniel E Harris, Daniel E Harris, Daniel T...

Publish Year: 2021

[Routine coagulation tests on ICU admission are associated ...](#)

<https://pubmed.ncbi.nlm.nih.gov/28681428>

The purpose of this retrospective single-centre study was to investigate whether activated partial thromboplastin time (APTT) or prothrombin time - **international normalized ratio** (PT-INR) measured on admission to the ICU in patients with severe sepsis or septic shock may be associated with mortality independent of SAPS 3 score.

Cited by: 7

Author: S. Benediktsson, A. Frigyesi, T. Kander

Publish Year: 2017

PEOPLE ALSO ASK

What are the inclusion criteria for sepsis diagnosis? ▾

Are there any competing interests in sepsis research? ▾

Why are emergency departments so slow to diagnose sepsis? ▾

What are the diagnostic codes for sepsis in Norway? ▾

Feedback

[International normalized ratio to albumin ratio - IJGM | IJGM](#)

<https://www.dovepress.com/international-normalized...> ▾

International Normalized Ratio to Albumin Ratio (PTAR): An Objective Risk Stratification Tool in Patients with Sepsis. Background: Sepsis is a life-threatening multiple-organ dysfunction caused by