

ALL IMAGES VIDEOS

13 Results Any time ▾

[A phase II trial of induction NAB-paclitaxel and cisplatin ...](#)<https://www.researchgate.net/publication/316837767...>

Request PDF | A phase II trial of induction NAB-paclitaxel and cisplatin followed by concurrent chemoradiotherapy in patients with locally advanced nasopharyngeal carcinoma | Objectives ...

[Secondary resectability in locally advanced pancreatic ...](#)<https://www.researchgate.net/publication/330095869...>

Home; Endocrine Gland Neoplasms; Neoplasms

[154361-50-9 | MFCD00930626 | 5'-Deoxy-5-fluoro-N ...](#)<https://www.aablocks.com/prod/154361-50-9> ▾

Title: Phase II clinical trial of induction chemotherapy with fixed dose rate gemcitabine and cisplatin followed by concurrent chemoradiotherapy with capecitabine for locally advanced pancreatic cancer. Journal: Cancer chemotherapy and pharmacology 20120901

Offer Count: 4 Price Range: \$5 - \$35

[www.science.gov](#)<https://www.science.gov/topicpages/a/advanced+inoperable+squamous.html> ▾

May 01, 2018 · Parenteral Nutrition for Patients Treated for Locally Advanced Inoperable Tumors of the Head and Neck. ClinicalTrials.gov. 2018-03-28. Squamous Cell Carcinoma of the Hypopharynx S

[Search result for 'advanced dermal fillers'](#)[https://www.dovepress.com/search\\_results.php?...](https://www.dovepress.com/search_results.php?...) ▾

Encouraging Pathological Complete Response Rate from Neoadjuvant Chemotherapy with Albumin-Bound Paclitaxel Plus Cisplatin and Capecitabine for Locally Advanced Esophageal Squamous Carcinoma: Preliminary Outcome of a Retrospective Study. Zhang W, Li Y, Xue L, Qu D, Jiang Z, Wang Z, Yang Z, Zhou A. Cancer Management and Research 2021, 13:2163-2170

[clinicaltrials.gov](#)<https://clinicaltrials.gov/ct2/results?displayxml=true&count=1000&start=54001>

Recruiting Advanced Breast Carcinoma; Anatomic Stage III Breast Cancer AJCC v8; Anatomic Stage IIIA Breast Cancer AJCC v8; Anatomic Stage IIIB Breast Cancer AJCC v8; Anatomic Stage

国内版 国际版

Microsoft Bing

Induction chemotherapy with albumin-bound paclitaxel plus lobap



All Images Videos

1,620 Results Any time ▾

### Liposome-paclitaxel and carboplatin combination ...

<https://www.sciencedirect.com/science/article/pii/S127832182100024X>

Jul 01, 2021 · Although the FP regimen is the standard regimen for CCRT for esophageal cancer, paclitaxel plus platinum agent has also been used as one of the main options for nearly a decade. In 2014, Tang et al. treated patients with locally advanced ESCC with 3-week schedule paclitaxel and cisplatin-based definitive CCRT. The results showed that the ...

Author: J. Zeng, X. Cui, L. Cheng, Y. Chen, X. ... Publish Year: 2021

Male: 72

### multi-institutional randomized phase III study comparing ...

<https://academic.oup.com/jjco/article/51/5/836/6174321> ▾

Mar 17, 2021 · A multi-institutional randomized phase III study comparing weekly carboplatin plus nab-paclitaxel and daily low-dose carboplatin as regimens for concurrent chemoradiotherapy in elderly patients with unresectable locally advanced non-small cell lung cancer: Japan Clinical Oncology Group Study JCOG1914

Author: Ryo Shimoyama, Shota Omori, Shog... Publish Year: 2021

### Phase II study of neoadjuvant therapy with nab-paclitaxel ...

<https://europepmc.org/article/MED/27244882> ▾

Aug 01, 2016 · Background We carried out a phase II study to evaluate the efficiency and safety of the combination of nanoparticle albumin bound-paclitaxel (nab-paclitaxel) and cisplatin as preoperative chemotherapy for locally advanced esophageal squamous cell carcinoma (ESCC)Results: From Oct 2011 to Dec 2012, 35 patients were enrolled and received neoadjuvant chemotherapy.

Cited by: 2 Author: Yun Fan, Youhua Jiang, Xinming Zhou, Qi...

Publish Year: 2016

### The antitumor effect of lobaplatin against Ishikawa ...


<https://www.sciencedirect.com/science/article/pii/S0753332219301647>

Jun 01, 2019 · The CCK-8 assay was used to evaluate anti-proliferative effect of lobaplatin on Ishikawa cells. After the treatment of lobaplatin for 48 h, the proliferation of Ishikawa cells was markedly attenuated and represented a dose-dependent manner (Fig. 1A). the inhibition rates were  $34.73 \pm 0.79\%$  and  $56.3 \pm 0.63\%$  respectively at 2ug and 4ug/mL dosage, which led to a significant inhibition of ...

20-Aug-2021 11:14AM

406 words • 20 matches • 8 sources

FAQ

 iThenticate

68661\_Auto\_Edited.docx

Quotes Excluded  
Bibliography Excluded

12%

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 68661

**Manuscript Type:** ORIGINAL ARTICLE

*Clinical Trials Study*

**Induction chemotherapy with albumin-bound paclitaxel plus lobaplatin followed by concurrent radiochemotherapy for locally advanced esophageal cancer**

Yan MH *et al.* Induction chemotherapy for locally advanced esophageal cancer

Mao-Hui Yan, Fang Liu, Bao-Lin Qu, Bo-Ning Cai, Wei Yu, Xiang-Kun Dai

**Abstract**

BACKGROUND

Match Overview

1

Internet 69 words  
created on 08-May-2021  
[www.sagepub.com](#)

3%

2

Internet 62 words  
created on 16-Feb-2021  
[wordedsciences.org](#)

2%

3

Internet 37 words  
created on 21-Sep-2020  
[www.frontiersin.org](#)

1%

4

Crossref 33 words  
Zhenhua Zhan, Yixue Wen, Dongbiao Liao, Jidong Miao *et al.* "Single-Agent Versus Double-Agent Chemotherapy in Co

1%

5

Internet 28 words  
[hdi.handle.net](#)

1%

6

Internet 22 words  
created on 01-Aug-2020  
[www.karger.com](#)

1%

7

Internet 20 words  
created on 09-Dec-2019  
[rs-journal.biomedcentral.com](#)

1%

8

Internet 17 words  
created on 18-Sep-2020  
[www.dovepress.com](#)

1%

PDF: 1 OF 10

Print Only Report

国内版

国际版

Induction chemotherapy with albumin-bound paclitaxel plus lobap



ALL

IMAGES

VIDEOS

1,590 Results

Any time ▾

### Phase II study of neoadjuvant therapy with nab-paclitaxel ...

<https://europepmc.org/article/MED/27244882> ▾

Aug 01, 2016 · Background We carried out a phase II study to evaluate the efficiency and safety of the combination of nanoparticle albumin bound-paclitaxel (nab-paclitaxel) and cisplatin as preoperative chemotherapy for locally advanced esophageal squamous cell carcinoma (ESCC) Results: From Oct 2011 to Dec 2012, 35 patients were enrolled and received neoadjuvant chemotherapy.

Cited by: 2

Author: Yun Fan, Youhua Jiang, Xinming Zhou, Qi...

Publish Year: 2016

### Liposome-paclitaxel and carboplatin combination ...

<https://www.sciencedirect.com/science/article/pii/S127832182100024X>

Jul 01, 2021 · Although the FP regimen is the standard regimen for CCRT for esophageal cancer, paclitaxel plus platinum agent has also been used as one of the main options for nearly a decade. In 2014, Tang et al. treated patients with locally advanced ESCC with 3-week schedule paclitaxel and cisplatin-based definitive CCRT. The results showed that the ...

Author: J. Zeng, X. Cui, L. Cheng, Y. Chen, X. ... Publish Year: 2021

Male: 72

### multi-institutional randomized phase III study comparing ...

<https://academic.oup.com/jjco/article/51/5/836/6174321> ▾

Mar 17, 2021 · A multi-institutional randomized phase III study comparing weekly carboplatin plus nab-paclitaxel and daily low-dose carboplatin as regimens for concurrent chemoradiotherapy in elderly patients with unresectable locally advanced non-small cell lung cancer: Japan Clinical Oncology Group Study JCOG1914

Author: Ryo Shimoyama, Shota Omori, Shog... Publish Year: 2021

### The antitumor effect of lobaplatin against Ishikawa ...

<https://www.sciencedirect.com/science/article/pii/S0753332219301647>

Jun 01, 2019 · The CCK-8 assay was used to evaluate anti-proliferative effect of lobaplatin on Ishikawa cells. After the treatment of lobaplatin for 48 h, the proliferation of Ishikawa cells was markedly attenuated and represented a dose-dependent manner (Fig. 1A). the inhibition rates were  $34.73 \pm 0.79\%$  and  $56.3 \pm 0.63\%$  respectively at 2ug and 4ug/mL dosage, which led to a significant inhibition of ...

Author: Jingchao He, Haijun Zhang Publish Year: 2019