

Supplementary Table 1 Baseline characteristics of statin user and non-user of 2016 HIRA-NPS data

Variable	Statin user	Statin non-user	<i>P</i>
	(<i>n</i> = 17413)	(<i>n</i> = 17267)	
Gender (male)	8,435 (48.44%)	10,075 (58.35%)	<0.0001
Age group (10-yr interval)			0.0770
30-39	319 (1.83%)	416 (2.41%)	
40-49	1,718 (9.87%)	1,853 (10.73%)	
50-59	4,559 (26.18%)	4,270 (24.73%)	
60-69	5,332 (30.62%)	4,677 (27.09%)	
≥ 70	5,485 (31.50%)	6,051 (35.04%)	

Supplementary Table 2 Baseline characteristics of statin user and non-user of 2017 HIRA-NPS data

Variable	Statin user	Statin non-user	P
	(n = 18707)	(n = 16882)	
Gender (male)	9,105 (48.67%)	9,958 (58.99%)	< 0.0001
Age group (10-year interval)			0.0631
30-39	415 (2.22%)	442 (2.62%)	
40-49	2,034 (10.87%)	1,946 (11.53%)	
50-59	5,024 (26.86%)	4,314 (25.55%)	
60-69	5,522 (29.52%)	4,559 (27.01%)	
≥ 70	5,712 (30.53%)	5,621 (33.30%)	

Supplementary Table 3 Baseline characteristics of metformin user and non-user of 2016 HIRA-NPS data

Variable	Metformin	Metformin	non-	<i>P</i>
	user	user		
	(<i>n</i> = 30683)	(<i>n</i> = 3922)		
Gender (male)	16,497 (53.77%)	1,950 (49.72%)		<0.000
				1
Age group (10-year interval)				<0.000
				1
30-39	647 (2.11%)	73 (1.86%)		
40-49	3,307 (10.78%)	246 (6.27%)		
50-59	8,113 (26.44%)	705 (17.98%)		
60-69	8,894 (28.99%)	1,093 (27.87%)		
≥ 70	9,722 (31.69%)	1,805 (46.02%)		

Supplementary Table 4 Baseline characteristics of metformin user and non-user of 2017 HIRA-NPS data

Variable	Metformin user (n = 31624)	Metformin user (n = 4004)	non- user	P
Gender (male)	17,117 (54.13%)	1,963 (49.03%)	<	0.0001
Age group (10-year interval)			<	0.0001
30-39	789 (2.49%)	71 (1.77%)		
40-49	3,706 (11.72%)	279 (6.97%)		
50-59	8,508 (26.90%)	848 (21.18%)		
60-69	9,049 (28.61%)	1,046 (26.12%)		
≥ 70	9,572 (30.27%)	1,760 (43.96%)		

Supplementary Table 5 Baseline characteristics of statin + metformin user and non-user of 2016 HIRA-NPS data

Variable	Statin	+	Statin	&	P
	Metformin user (n = 14524)	Metformin user (n = 2175)	Metformin non- user	Metformin non- user	
Gender (male)	7,080 (48.75%)	1,166 (53.61%)			<0.000 1
Age group (10-year interval)					<0.000 1
30-39	288 (1.98%)	48 (2.21%)			
40-49	1,587 (10.93%)	155 (7.13%)			
50-59	4,026 (27.72%)	359 (16.51%)			
60-69	4,432 (30.52%)	558 (25.66%)			
≥ 70	4,191 (28.86%)	1,055 (48.51%)			

Supplementary Table 6 Baseline characteristics of statin + metformin user and non-user of 2017 HIRA-NPS data

Variable	Statin	+	Statin	&	P
	Metformin user (n = 16096)	Metformin user (n = 2040)	Metformin non- user	Metformin non- user	
Gender (male)	7,869 (48.89%)	1,090 (53.43%)	<		0.0001
Age group (10-year interval)			<		0.0001
30-39	396 (2.46%)	50 (2.45%)			
40-49	1,880 (11.68%)	144 (7.06%)			
50-59	4,547 (28.25%)	442 (21.67%)			
60-69	4,748 (29.50%)	476 (23.33%)			
≥ 70	4,525 (28.11%)	928 (45.49%)			