

World Journal of *Clinical Cases*

World J Clin Cases 2022 April 6; 10(10): 2976-3320



Contents

Thrice Monthly Volume 10 Number 10 April 6, 2022

REVIEW

- 2976 Gut microbiota in gastrointestinal diseases during pregnancy
Liu ZZ, Sun JH, Wang WJ
- 2990 Targeting metabolism: A potential strategy for hematological cancer therapy
Tang X, Chen F, Xie LC, Liu SX, Mai HR

MINIREVIEWS

- 3005 Elevated intra-abdominal pressure: A review of current knowledge
Łagosz P, Sokolski M, Biegus J, Tycinska A, Zymlinski R

ORIGINAL ARTICLE

Case Control Study

- 3014 Changes in corneal nerve morphology and function in patients with dry eyes having type 2 diabetes
Fang W, Lin ZX, Yang HQ, Zhao L, Liu DC, Pan ZQ
- 3027 Combined sevoflurane-dexmedetomidine and nerve blockade on post-surgical serum oxidative stress biomarker levels in thyroid cancer patients
Du D, Qiao Q, Guan Z, Gao YF, Wang Q

Retrospective Cohort Study

- 3035 Early warning prevention and control strategies to reduce perioperative venous thromboembolism in patients with gastrointestinal cancer
Lu Y, Chen FY, Cai L, Huang CX, Shen XF, Cai LQ, Li XT, Fu YY, Wei J
- 3047 Dose-response relationship between risk factors and incidence of COVID-19 in 325 hospitalized patients: A multicenter retrospective cohort study
Zhao SC, Yu XQ, Lai XF, Duan R, Guo DL, Zhu Q

Retrospective Study

- 3060 Preventive online and offline health management intervention in polycystic ovary syndrome
Liu R, Li M, Wang P, Yu M, Wang Z, Zhang GZ
- 3069 Evidence-based intervention on postoperative fear, compliance, and self-efficacy in elderly patients with hip fracture
Fu Y, Zhu LJ, Li DC, Yan JL, Zhang HT, Xuan YH, Meng CL, Sun YH
- 3078 Significance of dysplasia in bile duct resection margin in patients with extrahepatic cholangiocarcinoma: A retrospective analysis
Choe JW, Kim HJ, Kim JS

- 3088** Diagnostic value and safety of medical thoracoscopy for pleural effusion of different causes

Liu XT, Dong XL, Zhang Y, Fang P, Shi HY, Ming ZJ

Observational Study

- 3101** Oxaliplatin-induced neuropathy and colo-rectal cancer patient's quality of life: Practical lessons from a prospective cross-sectional, real-world study

Prutianu I, Alexa-Stratulat T, Cristea EO, Nicolau A, Moisuc DC, Covrig AA, Ivanov K, Croitoru AE, Miron MI, Dinu MI, Ivanov AV, Marinca MV, Radu I, Gafton B

- 3113** Breast-conserving surgery and sentinel lymph node biopsy for breast cancer and their correlation with the expression of polyligand proteoglycan-1

Li FM, Xu DY, Xu Q, Yuan Y

SYSTEMATIC REVIEWS

- 3121** Clinical significance of aberrant left hepatic artery during gastrectomy: A systematic review

Tao W, Peng D, Cheng YX, Zhang W

META-ANALYSIS

- 3131** Betel quid chewing and oral potential malignant disorders and the impact of smoking and drinking: A meta-analysis

Lin HJ, Wang XL, Tian MY, Li XL, Tan HZ

- 3143** Effects of physical exercise on the quality-of-life of patients with haematological malignancies and thrombocytopenia: A systematic review and meta-analysis

Yang YP, Pan SJ, Qiu SL, Tung TH

CASE REPORT

- 3156** Primary malignant peritoneal mesothelioma mimicking tuberculous peritonitis: A case report

Lin LC, Kuan WY, Shiu BH, Wang YT, Chao WR, Wang CC

- 3164** Endoscopic submucosal dissection combined with adjuvant chemotherapy for early-stage neuroendocrine carcinoma of the esophagus: A case report

Tang N, Feng Z

- 3170** Lymph-node-first presentation of Kawasaki disease in a 12-year-old girl with cervical lymphadenitis caused by *Mycoplasma pneumoniae*: A case report

Kim N, Choi YJ, Na JY, Oh JW

- 3178** Tuberculosis-associated hemophagocytic lymphohistiocytosis misdiagnosed as systemic lupus erythematosus: A case report

Chen WT, Liu ZC, Li MS, Zhou Y, Liang SJ, Yang Y

- 3188** Migration of a Hem-o-Lok clip to the renal pelvis after laparoscopic partial nephrectomy: A case report

Sun J, Zhao LW, Wang XL, Huang JG, Fan Y

- 3194** Ectopic intrauterine device in the bladder causing cystolithiasis: A case report
Yu HT, Chen Y, Xie YP, Gan TB, Gou X
- 3200** Giant tumor resection under ultrasound-guided nerve block in a patient with severe asthma: A case report
Liu Q, Zhong Q, Zhou NN, Ye L
- 3206** Myomatous erythrocytosis syndrome: A case report
Shu XY, Chen N, Chen BY, Yang HX, Bi H
- 3213** Middle thyroid vein tumor thrombus in metastatic papillary thyroid microcarcinoma: A case report and review of literature
Gui Y, Wang JY, Wei XD
- 3222** Severe pneumonia and acute myocardial infarction complicated with pericarditis after percutaneous coronary intervention: A case report
Liu WC, Li SB, Zhang CF, Cui XH
- 3232** IgA nephropathy treatment with traditional Chinese medicine: A case report
Zhang YY, Chen YL, Yi L, Gao K
- 3241** Appendico-vesicocolonic fistula: A case report and review of literature
Yan H, Wu YC, Wang X, Liu YC, Zuo S, Wang PY
- 3251** *Scedosporium apiospermum* infection of the lumbar vertebrae: A case report
Shi XW, Li ST, Lou JP, Xu B, Wang J, Wang X, Liu H, Li SK, Zhen P, Zhang T
- 3261** Woman diagnosed with obsessive-compulsive disorder became delusional after childbirth: A case report
Lin SS, Gao JF
- 3268** Emphysematous pyelonephritis: Six case reports and review of literature
Ma LP, Zhou N, Fu Y, Liu Y, Wang C, Zhao B
- 3278** Atypical infantile-onset Pompe disease with good prognosis from mainland China: A case report
Zhang Y, Zhang C, Shu JB, Zhang F
- 3284** *Mycobacterium tuberculosis* bacteremia in a human immunodeficiency virus-negative patient with liver cirrhosis: A case report
Lin ZZ, Chen D, Liu S, Yu JH, Liu SR, Zhu ML
- 3291** Cervical aortic arch with aneurysm formation and an anomalous right subclavian artery and left vertebral artery: A case report
Wu YK, Mao Q, Zhou MT, Liu N, Yu X, Peng JC, Tao YY, Gong XQ, Yang L, Zhang XM
- 3297** Dedifferentiated chondrosarcoma of the middle finger arising from a solitary enchondroma: A case report
Yonezawa H, Yamamoto N, Hayashi K, Takeuchi A, Miwa S, Igarashi K, Morinaga S, Asano Y, Saito S, Tome Y, Ikeda H, Nojima T, Tsuchiya H

- 3306** Endoscopic-catheter-directed infusion of diluted (-)-noradrenaline for atypical hemobilia caused by liver abscess: A case report
Zou H, Wen Y, Pang Y, Zhang H, Zhang L, Tang LJ, Wu H
- 3313** *Pneumocystis jiroveci* pneumonia after total hip arthroplasty in a dermatomyositis patient: A case report
Hong M, Zhang ZY, Sun XW, Wang WG, Zhang QD, Guo WS

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Hui-Jeong Hwang, MD, PhD, Associate Professor, Department of Cardiology, Kyung Hee University Hospital at Gangdong, Kyung Hee University College of Medicine, Seoul 05278, South Korea. neonic7749@hanmail.net

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lai Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

April 6, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Betel quid chewing and oral potential malignant disorders and the impact of smoking and drinking: A meta-analysis

Hui-Jun Lin, Xiao-Lei Wang, Meng-Yuan Tian, Xing-Li Li, Hong-Zhuan Tan

Specialty type: Public, environmental and occupational health

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Jeng J

Received: July 12, 2021

Peer-review started: July 12, 2021

First decision: October 22, 2021

Revised: October 26, 2021

Accepted: February 23, 2022

Article in press: February 23, 2022

Published online: April 6, 2022



Hui-Jun Lin, Xiao-Lei Wang, Meng-Yuan Tian, Xing-Li Li, Hong-Zhuan Tan, Department of Epidemiology and Health Statistics, Xiangya School of Public Health, Central South University, Changsha 410008, Hunan Province, China

Corresponding author: Hong-Zhuan Tan, MD, Full Professor, Department of Epidemiology and Health Statistics, Xiangya School of Public Health, Central South University, No. 110 Xiangya Road, Changsha 410008, Hunan Province, China. tanhz99@qq.com

Abstract

BACKGROUND

Oral potential malignant disorders (OPMDs) are a precancerous condition of oral disease. Several studies have found that betel quid chewing, smoking and alcohol drinking might be the risk factors of OPMDs. But the relationships of them, especially their interaction are still inconclusive.

AIM

To evaluate the relationship between betel quid chewing and OPMDs and to explore the interaction of smoking and alcohol drinking on the relationship.

METHODS

We searched PubMed, Web of Science, Embase and the Cochrane Library databases with items complete until January 2021 for relevant studies. The research data were extracted according to the inclusion criteria. The pooled odds ratios (ORs) and 95% confidence intervals (CIs) were used to evaluate the effect size. Subgroup analysis was performed to assess interactions between exposures and OPMDs. Relative excess risk of interaction (RERI) was used to estimate the size of interaction.

RESULTS

Nine articles were selected in the final meta-analysis. The results showed that betel quid chewing (pooled OR: 8.70, 95% CI: 5.18-14.61), alcohol consumption (pooled OR: 1.95, 95% CI: 1.5-2.55), and smoking (pooled OR: 4.35, 95% CI: 3.06-6.2) could significantly increase the risk of OPMDs compared to individuals without these behaviors. Smoking and alcohol drinking synergistically increased the association between betel quid chewing and OPMDs (pooled OR_(BQ+SM): 14.38, 95% CI: 7.14-28.95; pooled OR_(BQ+DK): 11.12, 95% CI: 8.00-15.45, respectively). The RERI_(BQ+SM) and RERI_(BQ+DK) were 2.33 and 1.47, respectively.

CONCLUSION

The synergistic effects between smoking/drinking and betel quid highlights the importance of focusing on individuals with multiple exposures. Further study should be conducted to confirm these interactions.

Key Words: Oral potential malignant disorders; Betel quid chewing; Smoking; Drinking; Synergistic effect

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Betel quid chewing, smoking and drinking could significantly increase the risk of oral potential malignant disorders (OPMDs). And smoking and drinking synergistically increased the association between betel quid chewing and OPMDs.

Citation: Lin HJ, Wang XL, Tian MY, Li XL, Tan HZ. Betel quid chewing and oral potential malignant disorders and the impact of smoking and drinking: A meta-analysis. *World J Clin Cases* 2022; 10(10): 3131-3142

URL: <https://www.wjgnet.com/2307-8960/full/v10/i10/3131.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i10.3131>

INTRODUCTION

Oral cancer is the sixth most common cancer worldwide[1], occurring predominantly in developing countries. It has a high incidence in young adults creating a major economic burden to families and society. Diagnosis time is the most important factor affecting the prognosis; therefore, early diagnosis is crucial. Oral squamous cell carcinoma (OSCC) is the most common type of oral cancer and the majority of OSCCs are transformed from oral potential malignant disorders (OPMDs). OPMDs are a precancerous condition of oral disease which can be transformed and cured. Hence, controlling the disease in a precancerous state is the most effective prevention strategy. A meta-analysis found that the global prevalence of OPMDs was 4.47% higher in Asia and South America[2]. Leukoplakia, erythroplakia, oral submucous fibrosis (OSF) and oral lichen planus are four common categories of OPMDs focused upon in the study. Among them, OSF and leukoplakia, especially erythroplakia, have a higher risk of malignant transformation[3-5].

A number of studies have shown that betel quid and areca nut chewing is associated with OSF, leukoplakia and other premalignant disorders[6-8]. In contrast, the effects of smoking and alcohol consumption on OPMDs were mixed in studies, with some suggesting that smoking or alcohol increases the risk of diseases[9-11], while others reporting the opposite conclusion[10,12-13]. At present, about 600 million people in the world have the habit of betel quid chewing[14], accounting for about 10% of the global population with a younger age trend[15,16].

On the other hand, they generally have multiple exposures, such as most areca nut chewers are smokers[17-19]. If there is a joint effect among multiple exposure factors, the incidence of disease will greatly increase when multiple exposure factors exist. Lee *et al*[12] found that when smoking and betel quid were combined, the risks of OPMDs were greater than the sum of each of them alone, indicating the potential interaction between smoking and betel quid. However, Ray *et al*[20] did not observe such an interaction. Similarly, the same situation was observed in studies of betel nut and alcohol consumption exposure. Hence, whether smoking and alcohol consumption play a role in the relationship between betel quid and OPMDs is uncertain. The purpose of this meta-analysis was to evaluate the relationship between betel quid chewing and OPMDs and to explore the effects of smoking and alcohol drinking on this process.

MATERIALS AND METHODS

Search strategy and inclusion criteria

We systematically searched four databases: PubMed, Web of Science, Embase, and the Cochrane Library. The search terms were: (1) Oral precancerous lesions OR oral premalignant lesions OR oral potentially malignant disorders OR oral leukoplakia OR oral submucous fibrosis OR oral lichen planus OR oral erythema OR erythroplakia; (2) Betel nut OR betel quid OR areca nut OR pan chewing; (3) Alcohol OR drinking OR alcoholic beverage OR ethanol; and (4) Smoking OR cigarettes OR tobacco. The specific search strategy is shown in Table 1. The study language was limited to English. The data retrieval process was completed in January 2021.

Table 1 Search strategy

Number	Contents
1	Oral potentially malignant disorders OR oral precancerous lesions OR oral premalignant lesions OR oral leukoplakia OR oral submucous fibrosis OR oral lichen planus OR oral erythema OR erythroplakia
2	Betel quid OR betel nut OR areca nut OR pan chewing
3	Alcohol OR drinking OR alcoholic beverage OR ethanol
4	Tobacco OR smoking OR cigarettes
5	(#1) AND (#2) AND (#3)
6	(#1) AND (#2) AND (#4)
7	(#5) OR (#6)

The study inclusion criteria were: (1) The diagnosis of oral potentially malignant disorders is confirmed by a clinician and histological biopsy; (2) The exposure factors included both betel quid chewing and smoking, or both betel quid chewing and alcohol drinking; and (3) The study provided ORs and 95% confidence intervals (CIs) or sufficient information to calculate them.

Data extraction and quality assessment

After using Endnote software and manually to remove duplicates, literature screening was carried out. Firstly, two authors (Hui-Jun Lin and Xiao-Lei Wang) conducted preliminary screening by reading the titles and abstracts respectively, and then carefully read the full text to select the final literatures. Finally, the third author (Meng-Yuan Tian) was consulted to make the decision when the conclusions between the two authors were inconsistent.

The outcome indicators were oral potential malignant disorders, including oral leukoplakia, oral erythema, OSF and OLP. Though the specific mechanism of betel quid to every precancerous lesion were different, we believed that betel quid could cause all these diseases just with an inconsistent degree of damage after reading literatures. Therefore, it was reasonable to include the four precancerous lesions in the meta-analysis. Several precancerous lesions might be present in a study, or only one of them. The exposure categories were divided into five types: betel quid chewing (BQ), smoking (SM), alcohol drinking (DK), betel quid chewing and smoking (BQ + SM), betel quid chewing and alcohol drinking (BQ + DK). The presence of all three exposures is rarely mentioned in studies, this case was not considered in this meta-analysis. BQ included exposure to betel quid, betel nut, areca nut and other products containing betel quid; SM referred to exposure to cigarettes, bidi and other tobacco products; and DK stood for exposure to all alcoholic beverages, including beer, white wine, *etc.* BQ + SM means exposure to both SM and BQ. BQ + DK means exposure to both BQ and DK. All exposures were defined as having lasted at least 1 year. The odds ratios (ORs) and 95% confidence intervals (CIs) were extracted because the number of cases/controls in each exposure category were not specified. In the data extraction of BQ, SM, and DK, we chose the OR and 95%CI of the single exposure category, such as smoking only. If the single exposure category data were not provided in the article, the adjusted total population exposure data were selected. Other information, such as study design and types of oral precancerous lesions, were also collected in a standardized manner.

Newcastle-Ottawa Scale (NOS) criteria was used for the quality assessment of case-control studies respectively. NOS scale is a tool for evaluating the quality of case-control study with ten questions, consisting of three main components: Selection, Comparability and Exposure. The quality of cross-sectional study was assessed using the Agency for Healthcare Research and Quality (AHRQ) guidelines (11 questions) recommended by AHRQ. Each study was evaluated independently by two authors (Hui-Jun Lin and Xiao-Lei Wang), and they would make a discussion when the results were inconsistent. If no agreement could be reached, a third author (Meng-Yuan Tian) would join to discuss and reach a consensus. The higher the score of both scales, the better the quality. And the score of < 6, 6-7, and > 7 indicated that the studies had low, medium and high quality, respectively. Only literatures of medium quality or above could be included in this meta-analysis.

Data analysis

The data were analyzed using Stata 12.0 (Statacorp, Cannon Creek, TX, United States). We measured the effect size with ORs and 95% CIs, and performed heterogeneity tests using I^2 statistics. If I^2 was more than 50%, indicating greater heterogeneity, a random-effects model was used to calculate pooled effects; otherwise, a fixed-effects model was used. Sensitivity analysis was performed using the trim-and-fill method, which investigated whether the pooled OR changed significantly when the studies were eliminated one by one. Publication bias was assessed using Egger's regression and funnel plots, lnOR in the X-axis, standard error of lnOR in the Y-axis. Asymmetric funnel plots suggested the existence of bias.

Subgroup analysis was conducted to analyze the association between SM or DK and OPMDs, and preliminarily explore the interaction effect of SM/DK and BQ using stratified analysis. Relative excess risk of interaction (RERI) was used to estimate the size of interaction. A and B are denoted two exposure factors. A and B stand for their absence, with formula (If $RERI = 0$, there was no interaction[21]).

$$RERI_{AB} = OR_{AB} - OR_{A\bar{B}} - OR_{\bar{A}B} + 1$$

RESULTS

A total of 987 studies were retrieved, of which 344 were duplicates. We excluded 499 articles when reading titles and abstracts including 441 completely unrelated articles and 48 reviews or meta-analysis. And an additional 145 articles were excluded after reading full text. Finally, nine articles were included in the meta-analysis. The article selection process is shown in [Figure 1](#).

Study characteristics

[Table 2](#) presents the characteristics of the final nine studies. Only one study contained four precancerous lesions[22], the other studies only included fewer condition[7,12,13,20,23-26]. One of the studies reported leukoplakia and OSF as two separate diseases, and discussed the harm of betel quid to these two respectively[13], so the effect sizes were extracted separately. Among the studies, 9 studies reported the effect size of BQ, 9 of SM, 6 of DK, 8 of BQ + SM, 9 of BQ + DK. All studies included both males and females.

Betel quid chewing exposure and oral potential malignant disorders

Nine studies were included in our meta-analysis of the relationships between BQ exposure and OPMDs. The analysis revealed great heterogeneity ($I^2 = 73.5\%$, $P < 0.001$) between studies, so a random-effects model was used to estimate the pooled OR and 95%CI. The results suggested BQ was associated with an increased risk of OPMDs (pooled $OR_{(BQ)}$: 8.70, 95%CI: 5.18-14.61). A forest plot of the results of the individual studies and the pooled result was shown in [Figure 2](#).

Sensitivity analysis and publication bias

Sensitivity analysis suggested that the combined $OR_{(BQ)}$ was stable and that none of the individual studies disproportionately influenced the results. The funnel plot and Egger's regression test showed no evidence of publication bias ([Figure 3](#)).

Subgroup analysis

Separate effects of smoking or alcohol drinking: In order to better explore the role of SM or DK on the association between BQ and OPMDs, we first analyzed the separate effects of SM or DK with the disease. Nine studies were included in meta-analysis of the relationships between SM exposure and OPMDs, six of DK. Heterogeneity tests showed a lack of heterogeneity in DK ($I^2 = 0.0\%$, $P = 0.889$), but significant heterogeneity in SM ($I^2 = 61.6\%$, $P = 0.008$). We found both SM and DK increased the risk of OPMDs (pooled $OR_{(SM)}$: 4.35, 95%CI: 3.06-6.2; pooled $OR_{(DK)}$: 1.95, 95%CI: 1.5-2.55). There was no evidence of publication bias and the sensitivity analysis showed that the results were stable.

Effects of combined exposure with oral potential malignant disorders: In the assessment of combined BQ and SM, or BQ and DK, there was no evidence of publication bias and the sensitivity analysis showed the results were stable.

Effects of smoking and betel quid chewing on oral potential malignant disorders: Eight studies were included in our meta-analysis of the relationships between BQ + SM exposure and OPMDs. After the heterogeneity test, a random-effects model was used to calculate the pooled OR. The results showed that smoking and betel quid exposure increased the incidence of OPMDs (pooled $OR_{(BQ+SM)}$: 14.38, 95%CI: 7.14-28.95), and a higher risk degree than BQ alone ([Figure 4](#)).

Effect of alcohol drinking and betel quid chewing on oral potential malignant disorders: Nine studies were included in our meta-analysis of the relationships between BQ + DK exposure and OPMDs. A fixed-effects model was used to calculate the pooled OR for the risk of both BQ and DK exposure. We discovered alcohol consumption reinforced the effect of betel nut on OPMDs (pooled $OR_{(BQ+DK)}$: 11.12, 95%CI: 8.00-15.45) ([Figure 5](#)).

Interaction between smoking or alcohol drinking and betel quid

As shown in [Table 3](#), pooled $OR_{(BQ)}$ plus pooled $OR_{(SM)}$ was smaller than pooled $OR_{(BQ+SM)}$, and pooled $OR_{(BQ)}$ plus pooled $OR_{(DK)}$ was smaller than pooled $OR_{(BQ+DK)}$. We speculated that both SM and DK had synergistic effects with BQ. After calculation, $RERI_{(BQ+SM)}$ was 2.33, and $RERI_{(BQ+DK)}$ was 1.47, providing

Table 2 Selected studies characteristics

Ref.	Publication year	Study diseases	Study area	Study design	Sample size	Age	ORs and 95% confidence intervals					NOS/AHRQ scale
							BQ	SM	DK	BQ + SM	BQ + DK	
Yang <i>et al</i> [12]	2001	Leukoplakia, OSF	Taiwan	Cross sectional	312	20 and above	4.51 (1.30-15.62)	1.05 (0.5-2.23)	1.8 (0.88-3.73)	3.75 (0.99-14.19)	7.49 (1.95-28.69)	8
Lee <i>et al</i> [13]	2003	Leukoplakia	Taiwan	1:4 case control	125:876	15 and above	22.3 (11.3-43.8)	6.1 (3.4-10.6)	1.8 (1.1-2.8)	40.2 (16.3-99.2)	16.8 (7.2-39.5)	8
		OSF		1:4 case control	94:876	15 and above	40.7 (16-103.7)	7 (3.5-14.3)	1.8 (1.1-3.1)	57.9 (16-209.6)	31.7 (10.1-99.3)	
Yang <i>et al</i> [26]	2005	OSF	Taiwan	1:1 case control	62:62	20 and above	4.51 (1.20-16.94)	— —	— —	8.68 (1.87-40.23)	— —	7
Chung <i>et al</i> [25]	2005	Leukoplakia	Taiwan	Cross sectional	525	15 and above	— —	3.69 (1.61-8.43)	2.36 (0.91-6.13)	8.54 (2.58-28.23)	5.82 (1.10-30.84)	7
Thomas <i>et al</i> [7]	2008	Leukoplakia	PNG	Case control	197:1282	18 and above	3.8 (1.7-8.4)	6.4 (4.1-9.9)	— —	24.3 (8.7-67.4)	— —	6
Mehrotra <i>et al</i> [27]	2013	OSF	India	Case control	448:2688	15 and above	6.38 (4.29-9.48)	4.31 (2.84-6.55)	3.19 (1.36-7.45)	— —	9.79 (6.51-14.75)	8
Chher <i>et al</i> [23]	2018	OPMDs	Cambodia	Cross sectional	1614	18 and above	6.75 (3.32-13.72)	3.74 (1.89-7.41)	1.88 (0.81-4.36)	— —	5.40 (0.64-45.36)	7
Ikeda <i>et al</i> [24]	1995	OSF, lichen planus	Cambodia	Cross sectional	1319	15 and above	21.66 (5.06-92.75)	8.89 (1.78-44.42)	— —	17.16 (3.09-95.48)	— —	7
Ray <i>et al</i> [21]	2013	Leukoplakia, OSF	India	Case control	698:766	10 and above	5.15 (2.63-10.09)	4.38 (2.14-8.96)	— —	5.75 (2.32-14.27)	— —	7

ORs: odds ratios; OSF: Oral submucous fibrosis; OPMDs: Oral potential malignant disorders; BQ: Betel quid chewing; SM: Smoking; DK: Alcohol drinking; BQ + SM: Betel quid chewing and smoking; BQ + DK: Betel quid chewing and alcohol drinking; NOS: Newcastle-Ottawa Scale; AHRQ: The Agency for Healthcare Research and Quality.

further evidence of an interaction.


DISCUSSION

Our study found that betel quid chewing was associated with OPMDs. There was an interaction between smoking and betel quid chewing, and alcohol drinking and betel quid chewing with respect to their association with OPMDs. The data extracted from the original study was adjusted for confounding factors such as gender, age, and education status, though varies from article to article. And the sensitivity analysis also showed that our study results were stable with no significant bias. This meta-analysis provides a new approach for the prevention of OPMDs. It revealed that individuals with multiple exposures had very high risk because of synergistic interaction, which remind us that we should pay more attention to these people.

Table 3 Pooled odds ratios and 95% confidence intervals in all exposure categories

Exposure category	Pooled OR	95%CI
BQ	8.70	5.18-14.61
SM	4.35	3.06-6.20
DK	1.95	1.50-2.55
BQ and SM	14.38	7.14-28.95
BQ and DK	11.12	8.00-15.45

OR: Odds ratio; BQ: Betel quid chewing; SM: Smoking; DK: Alcohol drinking; CI: Confidence interval.

**Figure 1 Literature screening flow chart.**

Oral potential malignant disorders are a kind of disease with the risk of malignant transformation. It is of great significance for the prevention of oral cancer to realize the related risk factors and avoid them in daily life so as to reduce the occurrence of the disease. Areca nut is the fourth largest psychoactive substance in the world. The International Agency for Research on Cancer recognized that betel quid and areca nut were carcinogens in 2004[27], and studies have found that betel quid chewing is a risk factor for many diseases, including oral cancer, oropharyngeal cancer[28] and esophageal cancer[29]. And a variety of oral diseases have also been linked to betel quid chewing[30,31]. Studies found that betel quid harms the cavity mainly in two aspects: first, arecoline, the active ingredient in betel quid, would produce a variety of nitrosamines in the acidic environment of the oral cavity and stomach. And nitrosamines could interact with DNA, protein or other biological macromolecules, and then induce oxidative stress and participate in the canceration of oral mucosa. Causes of damage to the oral mucosa include oxidative stress caused by nitrification of arecoline and mechanical stimulation of the mucosa [32]. In addition, hydrated lime in betel quid could alter intracellular calcium homeostasis, causing abnormal activation of calcium-mediated transduction cascades and leading to oral cancer. Second, the mechanical stimulation of betel quid to oral mucosa caused by repeated chewing. Moreover, the current studies found that chewing betel quid could cause the changes in oral microbiota which might be related to oral diseases[33,34].

We found betel quid chewing has significantly increased the risk of OPMDs which is consistent with the majority of studies[35-37]. But the effects of betel quid varied greatly across studies and there was a large variation of the ORs[36,38,39]. It might be related to the study area and population. The production methods and ingredients of betel quid vary in different countries and the degree of oral damage also differed. But all the countries in the literature we included chew the raw betel quid,




Figure 2 Forest plot of betel quid chewing exposure and oral potential malignant disorders.




Figure 3 Funnel plot (betel quid chewing).

supplemented with such auxiliary materials as betel nut leaves and flowers, and all of them do not contain tobacco. Therefore, the results of our meta-analysis were robust and reliable.

In the subgroup analysis, we found that smoking alone was also associated with an increased risk of OPMDs, which is consistent with other studies[9,40]. There was an interaction between smoking and betel quid chewing, and tobacco enhanced the toxicity of betel quid to the oral mucosa. And studies showed that the risk of OPMDs in betel quid with tobacco was greater than that in betel quid alone[37]. Several potential mechanisms can explain this synergistic effect. The site in the oral cavity where tobacco is placed becomes keratinized due to friction. With the prolongation of tobacco use, the degree of mucosal keratosis deepens and plaques are formed[41]. The mechanical damage caused by betel quid chewing could accelerate this process. Alternatively, both smoking and betel quid chewing could lead to the accumulation of nitrosamines in the cavity and increase oxidative stress[32], which might exert a synergistic effect. Additionally, nicotine has been reported to have a synergistic effect on betel quid cytotoxicity[42]. And the synergistic effect has been reported in other diseases. Liu *et al*[43] found the interactions between betel quid and cigarette or alcohol in malignant transformation of OSF. Similarly, the same synergistic effects in oral cancer were suggested in the study of Petti *et al*[44].




Figure 4 Forest plot of betel quid chewing and smoking exposure and oral potential malignant disorders.




Figure 5 Forest plot of alcohol drinking and betel quid chewing exposure and oral potential malignant disorders.

In the relationship between alcohol consumption and OPMDs, different studies had different results [10,40,45]. Our subgroup analysis found that alcohol consumption alone had an association with OPMDs. It was mainly due to the direct stimulation to oral mucosa by alcohol, which could be enhanced if there is damage in the mouth. The results of our meta-analysis showed that betel quid chewing interacted with alcohol consumption. Animal experiments have shown that chronic alcohol exposure can cause atrophy of the oral mucosa, making it more sensitive to stimulation[46], such as the mechanical stimulation of betel nut chewing. The increased permeability caused by alcohol could accelerate the absorption of arecoline by the mucosa. Additionally, both betel quid chewing and alcohol drinking could change the composition of the oral microbiome[47]. Betel quid chewing combined with heavy alcohol drinking has been reported to change the diversity of the oral microbiome[33]. Oral microorganisms normally maintain oral health through immune regulation[48] and microbial disorders are associated with various diseases including OPMDs[49,50].

There are still some limitations to our study. The selected studies had cross-sectional and case-control study design, which provide weak evidence of causality. The number of people in each subgroup may have been inadequate because of having five exposure categories, though the direct number were not reported. Further investigation should be conducted with a large cohort study to accurately assess the impact of smoking and alcohol drinking on the association between betel quid chewing and OPMDs.

CONCLUSION

The meta-analysis suggested that betel quid chewing is associated with an increased risk of oral potential malignant disorders. There was a synergistic effect between smoking or alcohol drinking and betel quid chewing. Therefore, we should focus on high-risk groups with multiple exposure, especially individuals with smoking and betel quid exposed, and provide health education about the harmful effects of unhealthy behavior. Governments should develop policies to quit betel quid, smoking and alcohol, especially in individuals with multiple exposure, in order to control and reduce the incidence of oral potential malignant disorders.

ARTICLE HIGHLIGHTS

Research background

Studies have found that betel quid chewing, smoking and alcohol drinking might be the risk factors of oral potential malignant disorders (OPMDs). But the relationships of them, especially the interaction are still inconclusive.

Research motivation

This meta-analysis was to determine the damage of betel quid on OPMDs and to explore the potential interaction between smoking, alcohol drinking and betel quid. It could provide new ideas for the prevention of OPMDs.

Research objectives

To evaluate the relationship between betel quid chewing and OPMDs and to explore the interaction of smoking and drinking on the relationship.

Research methods

We searched four database and finally selected 9 from 987 studies. The research data were extracted according to the inclusion criteria. The pooled odds ratios (ORs) and 95% confidence intervals (CIs) were used to evaluate the effect size and relative excess risk of interaction (RERI) was used to estimate the size of interaction.

Research results

The results showed that betel quid chewing (pooled OR: 8.70, 95%CI: 5.18-14.61) could significantly increase the risk of OPMDs. And smoking and drinking synergistically increased the association between betel quid chewing and OPMDs (pooled OR_(BQ+SM): 14.38, 95%CI: 7.14-28.95, RERI_(BQ+SM): 2.33; pooled OR_(BQ+DK): 11.12, 95%CI: 8.00-15.45, RERI_(BQ+DK): 1.47, respectively).

Research conclusions

Betel quid chewing is associated with an increased risk of oral potential malignant disorders. There was a synergistic effect between smoking or drinking and betel quid chewing.

Research perspectives

Governments should focus on individuals with multiple exposure.

ACKNOWLEDGEMENTS

We are thankful to Tan HZ for his support guidance and Wang XL, Tian MY and Li XL to contributions in data analysis.

FOOTNOTES

Author contributions: Lin HJ contributed to the literature search, data extraction, analysis and interpretation and manuscript drafting; Wang XL and Tian MY contributed to the literature search, data extraction and analysis; Li XL contributed to data interpretation and manuscript revision; Tan HZ contributed to data interpretation, manuscript drafting and revision; all authors read and approved the final manuscript.

Conflict-of-interest statement: The authors declare having no potential conflicts of interest.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Hui-Jun Lin 0000-0001-9284-6625; Xiao-Lei Wang 0000-0001-8049-2177; Meng-Yuan Tian 0000-0002-5416-3046; Xing-Li Li 0000-0001-8922-6708; Hong-Zhuan Tan 0000-0002-4292-5947.

S-Editor: Wu YXJ

L-Editor: Filipodia

P-Editor: Guo X

REFERENCES

- 1 **Warnakulasuriya S.** Living with oral cancer: epidemiology with particular reference to prevalence and life-style changes that influence survival. *Oral Oncol* 2010; **46**: 407-410 [PMID: 20403722 DOI: 10.1016/j.oraloncology.2010.02.015]
- 2 **Mello FW, Miguel AFP, Dutra KL, Porporatti AL, Warnakulasuriya S, Guerra ENS, Rivero ERC.** Prevalence of oral potentially malignant disorders: A systematic review and meta-analysis. *J Oral Pathol Med* 2018; **47**: 633-640 [PMID: 29738071 DOI: 10.1111/jop.12726]
- 3 **Jayasooriya PR, Dayaratne K, Dissanayake UB, Warnakulasuriya S.** Malignant transformation of oral leukoplakia: a follow-up study. *Clin Oral Investig* 2020; **24**: 4563-4569 [PMID: 32436158 DOI: 10.1007/s00784-020-03322-4]
- 4 **Fitzpatrick SG, Hirsch SA, Gordon SC.** The malignant transformation of oral lichen planus and oral lichenoid lesions: a systematic review. *J Am Dent Assoc* 2014; **145**: 45-56 [PMID: 24379329 DOI: 10.14219/jada.2013.10]
- 5 **Iocca O, Sollecito TP, Alawi F, Weinstein GS, Newman JG, De Virgilio A, Di Maio P, Spriano G, Pardiñas López S, Shanti RM.** Potentially malignant disorders of the oral cavity and oral dysplasia: A systematic review and meta-analysis of malignant transformation rate by subtype. *Head Neck* 2020; **42**: 539-555 [PMID: 31803979 DOI: 10.1002/hed.26006]
- 6 **Ray JG, Chatterjee R, Chaudhuri K.** Oral submucous fibrosis: A global challenge. Rising incidence, risk factors, management, and research priorities. *Periodontol* 2000 2019; **80**: 200-212 [PMID: 31090137 DOI: 10.1111/prd.12277]
- 7 **Thomas SJ, Harris R, Ness AR, Tauro J, MacLennan R, Howes N, Bain CJ.** Betel quid not containing tobacco and oral leukoplakia: a report on a cross-sectional study in Papua New Guinea and a meta-analysis of current evidence. *Int J Cancer* 2008; **123**: 1871-1876 [PMID: 18688850 DOI: 10.1002/ijc.23739]
- 8 **Paulino YC, Hurwitz EL, Warnakulasuriya S, Gatewood RR, Pierson KD, Tenorio LF, Novotny R, Palafox NA, Wilkens LR, Badowski G.** Screening for oral potentially malignant disorders among areca (betel) nut chewers in Guam and Saipan. *BMC Oral Health* 2014; **14**: 151 [PMID: 25495475 DOI: 10.1186/1472-6831-14-151]
- 9 **Kusiak A, Maj A, Cichońska D, Kochańska B, Cydejko A, Świetlik D.** The Analysis of the Frequency of Leukoplakia in Reference of Tobacco Smoking among Northern Polish Population. *Int J Environ Res Public Health* 2020; **17** [PMID: 32971842 DOI: 10.3390/ijerph17186919]
- 10 **Shiu MN, Chen TH, Chang SH, Hahn LJ.** Risk factors for leukoplakia and malignant transformation to oral carcinoma: a leukoplakia cohort in Taiwan. *Br J Cancer* 2000; **82**: 1871-1874 [PMID: 10839305 DOI: 10.1054/bjoc.2000.1208]
- 11 **Maserejian NN, Joshupura KJ, Rosner BA, Giovannucci E, Zavras AI.** Prospective study of alcohol consumption and risk of oral premalignant lesions in men. *Cancer Epidemiol Biomarkers Prev* 2006; **15**: 774-781 [PMID: 16614123 DOI: 10.1158/1055-9965.EPI-05-0842]
- 12 **Yang YH, Lee HY, Tung S, Shieh TY.** Epidemiological survey of oral submucous fibrosis and leukoplakia in aborigines of Taiwan. *J Oral Pathol Med* 2001; **30**: 213-219 [PMID: 11302240 DOI: 10.1034/j.1600-0714.2001.300404.x]
- 13 **Lee CH, Ko YC, Huang HL, Chao YY, Tsai CC, Shieh TY, Lin LM.** The precancer risk of betel quid chewing, tobacco use and alcohol consumption in oral leukoplakia and oral submucous fibrosis in southern Taiwan. *Br J Cancer* 2003; **88**: 366-372 [PMID: 12569378 DOI: 10.1038/sj.bjc.6600727]
- 14 **Herzog TA, Pokhrel P.** Introduction to the Special Issue: International Research on Areca Nut and Betel Quid Use. *Subst Use Misuse* 2020; **55**: 1383-1384 [PMID: 32569543 DOI: 10.1080/10826084.2020.1767345]
- 15 **Oakley E, Demaine L, Warnakulasuriya S.** Areca (betel) nut chewing habit among high-school children in the Commonwealth of the Northern Mariana Islands (Micronesia). *Bull World Health Organ* 2005; **83**: 656-660 [PMID: 16141233 DOI: 10.1186/1472-6831-83-656]

- 16211156]
- 16 **Wang SC**, Tsai CC, Huang ST, Hong YJ. Betel nut chewing and related factors in adolescent students in Taiwan. *Public Health* 2003; **339** [DOI: [10.1016/S0033-3506\(03\)00082-9](https://doi.org/10.1016/S0033-3506(03)00082-9)]
 - 17 **Lee CH**, Ko AM, Warnakulasuriya S, Yin BL, Sunarjo, Zain RB, Ibrahim SO, Liu ZW, Li WH, Zhang SS, Kuntoro, Utomo B, Rajapakse PS, Warusavithana SA, Razak IA, Abdullah N, Shrestha P, Kwan AL, Shieh TY, Chen MK, Ko YC. Intercountry prevalences and practices of betel-quid use in south, southeast and eastern Asia regions and associated oral preneoplastic disorders: an international collaborative study by Asian betel-quid consortium of south and east Asia. *Int J Cancer* 2011; **129**: 1741-1751 [PMID: [21128235](https://pubmed.ncbi.nlm.nih.gov/21128235/) DOI: [10.1002/ijc.25809](https://doi.org/10.1002/ijc.25809)]
 - 18 **Hsu KY**, Tsai YF, Huang CC, Yeh WL, Chang KP, Ling CC, Chen CY, Lee HL. Tobacco-Smoking, Alcohol-Drinking, and Betel-Quid-Chewing Behaviors: Development and Use of a Web-Based Survey System. *Jmir Mhealth Uhealth* 2018 [DOI: [10.2196/preprints.9783](https://doi.org/10.2196/preprints.9783)]
 - 19 **Wen CP**, Tsai SP, Cheng TY, Chen CJ, Levy DT, Yang HJ, Eriksen MP. Uncovering the relation between betel quid chewing and cigarette smoking in Taiwan. *Tob Control* 2005; **14** Suppl 1: i16-i22 [PMID: [15923442](https://pubmed.ncbi.nlm.nih.gov/15923442/) DOI: [10.1136/tc.2004.008003](https://doi.org/10.1136/tc.2004.008003)]
 - 20 **Ray JG**, Ganguly M, Rao BS, Mukherjee S, Mahato B, Chaudhuri K. Clinico-epidemiological profile of oral potentially malignant and malignant conditions among areca nut, tobacco and alcohol users in Eastern India: A hospital based study. *J Oral Maxillofac Pathol* 2013; **17**: 45-50 [PMID: [23798829](https://pubmed.ncbi.nlm.nih.gov/23798829/) DOI: [10.4103/0973-029X.110720](https://doi.org/10.4103/0973-029X.110720)]
 - 21 **Chu H**, Nie L, Cole SR. Estimating the relative excess risk due to interaction: a bayesian approach. *Epidemiology* 2011; **22**: 242-248 [PMID: [21228700](https://pubmed.ncbi.nlm.nih.gov/21228700/) DOI: [10.1097/EDE.0b013e318208750e](https://doi.org/10.1097/EDE.0b013e318208750e)]
 - 22 **Chher T**, Hak S, Kallarakkal TG, Durward C, Ramanathan A, Ghani WMN, Razak IA, Harun MH, Ashar NAM, Rajandram RK, Prak P, Hussaini HM, Zain RB. Prevalence of oral cancer, oral potentially malignant disorders and other oral mucosal lesions in Cambodia. *Ethn Health* 2018; **23**: 1-15 [PMID: [27781495](https://pubmed.ncbi.nlm.nih.gov/27781495/) DOI: [10.1080/13557858.2016.1246431](https://doi.org/10.1080/13557858.2016.1246431)]
 - 23 **Ikeda N**, Handa Y, Khim SP, Durward C, Axéll T, Mizuno T, Fukano H, Kawai T. Prevalence study of oral mucosal lesions in a selected Cambodian population. *Community Dent Oral Epidemiol* 1995; **23**: 49-54 [PMID: [7774177](https://pubmed.ncbi.nlm.nih.gov/7774177/) DOI: [10.1111/j.1600-0528.1995.tb00197.x](https://doi.org/10.1111/j.1600-0528.1995.tb00197.x)]
 - 24 **Chung CH**, Yang YH, Wang TY, Shieh TY, Warnakulasuriya S. Oral precancerous disorders associated with areca quid chewing, smoking, and alcohol drinking in southern Taiwan. *J Oral Pathol Med* 2005; **34**: 460-466 [PMID: [16091112](https://pubmed.ncbi.nlm.nih.gov/16091112/) DOI: [10.1111/j.1600-0714.2005.00332.x](https://doi.org/10.1111/j.1600-0714.2005.00332.x)]
 - 25 **Yang YH**, Lien YC, Ho PS, Chen CH, Chang JS, Cheng TC, Shieh TY. The effects of chewing areca/betel quid with and without cigarette smoking on oral submucous fibrosis and oral mucosal lesions. *Oral Dis* 2005; **11**: 88-94 [PMID: [15752081](https://pubmed.ncbi.nlm.nih.gov/15752081/) DOI: [10.1111/j.1601-0825.2004.01061.x](https://doi.org/10.1111/j.1601-0825.2004.01061.x)]
 - 26 **Mehrotra D**, Kumar S, Agarwal GG, Asthana A. Odds ratio of risk factors for oral submucous fibrosis in a case control model. *Br J Oral Maxillofac Surg* 2013; **51**: e169-e173 [PMID: [22951384](https://pubmed.ncbi.nlm.nih.gov/22951384/) DOI: [10.1016/j.bjoms.2012.08.001](https://doi.org/10.1016/j.bjoms.2012.08.001)]
 - 27 **Ko AM**, Lee CH, Ko YC. Betel quid-associated cancer: Prevention strategies and targeted treatment. *Cancer Lett* 2020; **477**: 60-69 [PMID: [32112902](https://pubmed.ncbi.nlm.nih.gov/32112902/) DOI: [10.1016/j.canlet.2020.02.030](https://doi.org/10.1016/j.canlet.2020.02.030)]
 - 28 **Lambert R**, Sauvaget C, de Camargo Cancela M, Sankaranarayanan R. Epidemiology of cancer from the oral cavity and oropharynx. *Eur J Gastroenterol Hepatol* 2011; **23**: 633-641 [PMID: [21654320](https://pubmed.ncbi.nlm.nih.gov/21654320/) DOI: [10.1097/MEG.0b013e3283484795](https://doi.org/10.1097/MEG.0b013e3283484795)]
 - 29 **Akhtar S**, Sheikh AA, Qureshi HU. Chewing areca nut, betel quid, oral snuff, cigarette smoking and the risk of oesophageal squamous-cell carcinoma in South Asians: a multicentre case-control study. *Eur J Cancer* 2012; **48**: 655-661 [PMID: [21733677](https://pubmed.ncbi.nlm.nih.gov/21733677/) DOI: [10.1016/j.ejca.2011.06.008](https://doi.org/10.1016/j.ejca.2011.06.008)]
 - 30 **Giovannoni ML**, Valdivia-Gandur I, Lozano de Luaces V, Varela Véliz H, Balasubbaiah Y, Chimenos-Küstner E. Betel and tobacco chewing habit and its relation to risk factors for periodontal disease. *Oral Dis* 2018; **24**: 829-839 [PMID: [29284200](https://pubmed.ncbi.nlm.nih.gov/29284200/) DOI: [10.1111/odi.12820](https://doi.org/10.1111/odi.12820)]
 - 31 **Parmar G**, Sangwan P, Vashi P, Kulkarni P, Kumar S. Effect of chewing a mixture of areca nut and tobacco on periodontal tissues and oral hygiene status. *J Oral Sci* 2008; **50**: 57-62 [PMID: [18403885](https://pubmed.ncbi.nlm.nih.gov/18403885/) DOI: [10.2334/josnusd.50.57](https://doi.org/10.2334/josnusd.50.57)]
 - 32 **Islam S**, Muthumala M, Matsuoka H, Uehara O, Kuramitsu Y, Chiba I, Abiko Y. How Each Component of Betel Quid Is Involved in Oral Carcinogenesis: Mutual Interactions and Synergistic Effects with Other Carcinogens-a Review Article. *Curr Oncol Rep* 2019; **21**: 53 [PMID: [31028548](https://pubmed.ncbi.nlm.nih.gov/31028548/) DOI: [10.1007/s11912-019-0800-8](https://doi.org/10.1007/s11912-019-0800-8)]
 - 33 **Hernandez BY**, Zhu X, Goodman MT, Gatewood R, Mendiola P, Quinata K, Paulino YC. Betel nut chewing, oral premalignant lesions, and the oral microbiome. *PLoS One* 2017; **12**: e0172196 [PMID: [28225785](https://pubmed.ncbi.nlm.nih.gov/28225785/) DOI: [10.1371/journal.pone.0172196](https://doi.org/10.1371/journal.pone.0172196)]
 - 34 **Chen MY**, Chen JW, Wu LW, Huang KC, Chen JY, Wu WS, Chiang WF, Shih CJ, Tsai KN, Hsieh WT, Ho YH, Wong TY, Wu JH, Chen YL. Carcinogenesis of Male Oral Submucous Fibrosis Alters Salivary Microbiomes. *J Dent Res* 2021; **100**: 397-405 [PMID: [33089709](https://pubmed.ncbi.nlm.nih.gov/33089709/) DOI: [10.1177/0022034520968750](https://doi.org/10.1177/0022034520968750)]
 - 35 **Amarasinghe HK**, Usgodaarachchi US, Johnson NW, Laloo R, Warnakulasuriya S. Betel-quid chewing with or without tobacco is a major risk factor for oral potentially malignant disorders in Sri Lanka: a case-control study. *Oral Oncol* 2010; **46**: 297-301 [PMID: [20189448](https://pubmed.ncbi.nlm.nih.gov/20189448/) DOI: [10.1016/j.oraloncology.2010.01.017](https://doi.org/10.1016/j.oraloncology.2010.01.017)]
 - 36 **Chen PC**, Pan CC, Kuo C, Lin CP. Risk of oral nonmalignant lesions associated with human papillomavirus infection, betel quid chewing, and cigarette smoking in Taiwan: an integrated molecular and epidemiologic study. *Arch Pathol Lab Med* 2006; **130**: 57-61 [PMID: [16390239](https://pubmed.ncbi.nlm.nih.gov/16390239/) DOI: [10.5858/2006-130-57-ROONLA](https://doi.org/10.5858/2006-130-57-ROONLA)]
 - 37 **Zaw KK**, Ohnmar M, Hlaing MM, Oo YT, Win SS, Htike MM, Aye PP, Shwe S, Htwe MT, Thein ZM. Betel Quid and Oral Potentially Malignant Disorders in a Periurban Township in Myanmar. *PLoS One* 2016; **11**: e0162081 [PMID: [27611195](https://pubmed.ncbi.nlm.nih.gov/27611195/) DOI: [10.1371/journal.pone.0162081](https://doi.org/10.1371/journal.pone.0162081)]
 - 38 **Song H**, Wan Y, Xu YY. Betel Quid Chewing Without Tobacco: A Meta-analysis of Carcinogenic and Precarcinogenic Effects. *Asia-Pac J Public He* 2015; **Np47** [DOI: [10.1177/1010539513486921](https://doi.org/10.1177/1010539513486921)]
 - 39 **Chen PH**, Mahmood Q, Mariottini GL, Chiang TA, Lee KW. Adverse Health Effects of Betel Quid and the Risk of Oral and Pharyngeal Cancers. *Biomed Res Int* 2017; **2017**: 3904098 [PMID: [29376073](https://pubmed.ncbi.nlm.nih.gov/29376073/) DOI: [10.1155/2017/3904098](https://doi.org/10.1155/2017/3904098)]
 - 40 **Thomas G**, Hashibe M, Jacob BJ, Ramadas K, Mathew B, Sankaranarayanan R, Zhang ZF. Risk factors for multiple oral premalignant lesions. *Int J Cancer* 2003; **107**: 285-291 [PMID: [12949809](https://pubmed.ncbi.nlm.nih.gov/12949809/) DOI: [10.1002/ijc.11383](https://doi.org/10.1002/ijc.11383)]

- 41 **Muthukrishnan A**, Warnakulasuriya S. Oral health consequences of smokeless tobacco use. *Indian J Med Res* 2018; **148**: 35-40 [PMID: [30264752](#) DOI: [10.4103/ijmr.IJMR_1793_17](#)]
- 42 **Chang YC**, Lii CK, Tai KW, Chou MY. Adverse effects of arecoline and nicotine on human periodontal ligament fibroblasts in vitro. *J Clin Periodontol* 2001; **28**: 277-282 [PMID: [11284543](#) DOI: [10.1034/j.1600-051x.2001.028003277.x](#)]
- 43 **Liu B**, Shen M, Xiong J, Yuan Y, Wu X, Gao X, Xu J, Guo F, Jian X. Synergistic effects of betel quid chewing, tobacco use (in the form of cigarette smoking), and alcohol consumption on the risk of malignant transformation of oral submucous fibrosis (OSF): a case-control study in Hunan Province, China. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2015; **120**: 337-345 [PMID: [26130067](#) DOI: [10.1016/j.oooo.2015.04.013](#)]
- 44 **Petti S**, Masood M, Scully C. The magnitude of tobacco smoking-betel quid chewing-alcohol drinking interaction effect on oral cancer in South-East Asia. A meta-analysis of observational studies. *PLoS One* 2013; **8**: e78999 [PMID: [24260143](#) DOI: [10.1371/journal.pone.0078999](#)]
- 45 **Varshney S**, Sandhir S, Mishra S. A study of oral pre-malignant lesions and related risk factors. *Indian J Community Health* 2015; **130**
- 46 **Wight AJ**, Ogden GR. Possible mechanisms by which alcohol may influence the development of oral cancer--a review. *Oral Oncol* 1998; **34**: 441-447 [PMID: [9930353](#) DOI: [10.1016/s1368-8375\(98\)00022-0](#)]
- 47 **Uehara O**, Hiraki D, Kuramitsu Y, Matsuoka H, Takai R, Fujita M, Harada F, Paudel D, Takahashi S, Yoshida K, Muthumala M, Nagayasu H, Chiba I, Abiko Y. Alteration of oral flora in betel quid chewers in Sri Lanka. *J Microbiol Immunol Infect* 2021; **54**: 1159-1166 [PMID: [32611503](#) DOI: [10.1016/j.jmii.2020.06.009](#)]
- 48 **Zhong XH**, Lu Q, Zhang Q, He Y, Wei WJ, Wang YM. Oral microbiota alteration associated with oral cancer and areca chewing. *Oral Dis* 2021; **226** [DOI: [10.1111/odi.13545](#)]
- 49 **Zhang Y**, Wang X, Li H, Ni C, Du Z, Yan F. Human oral microbiota and its modulation for oral health. *Biomed Pharmacother* 2018; **99**: 883-893 [PMID: [29710488](#) DOI: [10.1016/j.biopha.2018.01.146](#)]
- 50 **Deng S**, Xu Y, Wang X, Liu M, Li L, Yu X, Wang Y, Wu Y, Wang W, Gao M, Cong B. Study on the Role of Salivary Flora and NF-κB Inflammatory Signal Pathway in Oral Lichen Planus. *Inflammation* 2020; **43**: 994-1008 [PMID: [32016629](#) DOI: [10.1007/s10753-020-01185-1](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

