

World Journal of *Clinical Cases*

World J Clin Cases 2022 May 16; 10(14): 4327-4712



OPINION REVIEW

- 4327 Emerging role of biosimilars in the clinical care of inflammatory bowel disease patients
Najeeb H, Yasmin F, Surani S

MINIREVIEWS

- 4334 Practical insights into chronic management of hepatic Wilson's disease
Lynch EN, Campani C, Innocenti T, Dragoni G, Forte P, Galli A
- 4348 Adipose-derived stem cells in the treatment of hepatobiliary diseases and sepsis
Satilmis B, Cicek GS, Cicek E, Akbulut S, Sahin TT, Yilmaz S

ORIGINAL ARTICLE**Clinical and Translational Research**

- 4357 Learning curve for a surgeon in robotic pancreaticoduodenectomy through a "G"-shaped approach: A cumulative sum analysis
Wei ZG, Liang CJ, Du Y, Zhang YP, Liu Y
- 4368 Clinical and prognostic significance of expression of phosphoglycerate mutase family member 5 and Parkin in advanced colorectal cancer
Wu C, Feng ML, Jiao TW, Sun MJ

Case Control Study

- 4380 Significance of preoperative peripheral blood neutrophil-lymphocyte ratio in predicting postoperative survival in patients with multiple myeloma bone disease
Xu ZY, Yao XC, Shi XJ, Du XR

Retrospective Study

- 4395 Association between depression and malnutrition in pulmonary tuberculosis patients: A cross-sectional study
Fang XE, Chen DP, Tang LL, Mao YJ
- 4404 Pancreatic cancer incidence and mortality patterns in 2006-2015 and prediction of the epidemiological trend to 2025 in China
Yin MY, Xi LT, Liu L, Zhu JZ, Qian LJ, Xu CF
- 4414 Evaluation of short- and medium-term efficacy and complications of ultrasound-guided ablation for small liver cancer
Zhong H, Hu R, Jiang YS

- 4425** Hematopoiesis reconstitution and anti-tumor effectiveness of Pai-Neng-Da capsule in acute leukemia patients with haploidentical hematopoietic stem cell transplantation

Yuan JJ, Lu Y, Cao JJ, Pei RZ, Gao RL

- 4436** Oral and maxillofacial pain as the first sign of metastasis of an occult primary tumour: A fifteen-year retrospective study

Shan S, Liu S, Yang ZY, Wang TM, Lin ZT, Feng YL, Pakezhati S, Huang XF, Zhang L, Sun GW

- 4446** Reduced serum high-density lipoprotein cholesterol levels and aberrantly expressed cholesterol metabolism genes in colorectal cancer

Tao JH, Wang XT, Yuan W, Chen JN, Wang ZJ, Ma YB, Zhao FQ, Zhang LY, Ma J, Liu Q

Observational Study

- 4460** Correlation of pressure gradient in three hepatic veins with portal pressure gradient

Wang HY, Song QK, Yue ZD, Wang L, Fan ZH, Wu YF, Dong CB, Zhang Y, Meng MM, Zhang K, Jiang L, Ding HG, Zhang YN, Yang YP, Liu FQ

- 4470** Multi-slice spiral computed tomography in diagnosing unstable pelvic fractures in elderly and effect of less invasive stabilization

Huang JG, Zhang ZY, Li L, Liu GB, Li X

SYSTEMATIC REVIEWS

- 4480** Distribution and changes in hepatitis C virus genotype in China from 2010 to 2020

Yang J, Liu HX, Su YY, Liang ZS, Rao HY

CASE REPORT

- 4494** Bow hunter's syndrome successfully treated with a posterior surgical decompression approach: A case report and review of literature

Orlandi N, Cavallieri F, Grisendi I, Romano A, Ghadirpour R, Napoli M, Moratti C, Zanichelli M, Pascarella R, Valzania F, Zedde M

- 4502** Histological remission of eosinophilic esophagitis under asthma therapy with IL-5 receptor monoclonal antibody: A case report

Huguenot M, Bruhm AC, Essig M

- 4509** Cutaneous mucosa-associated lymphoid tissue lymphoma complicating Sjögren's syndrome: A case report and review of literature

Liu Y, Zhu J, Huang YH, Zhang QR, Zhao LL, Yu RH

- 4519** Plexiform neurofibroma of the cauda equina with follow-up of 10 years: A case report

Chomanskis Z, Juskys R, Cepkus S, Dulko J, Hendrixson V, Ruksenas O, Rocka S

- 4528** Mixed porokeratosis with a novel mevalonate kinase gene mutation: A case report

Xu HJ, Wen GD

- 4535** Isolated pancreatic injury caused by abdominal massage: A case report

Sun BL, Zhang LL, Yu WM, Tuo HF

- 4541** Bronchiolar adenoma with unusual presentation: Two case reports
Du Y, Wang ZY, Zheng Z, Li YX, Wang XY, Du R
- 4550** Periodontal-orthodontic interdisciplinary management of a “periodontally hopeless” maxillary central incisor with severe mobility: A case report and review of literature
Jiang K, Jiang LS, Li HX, Lei L
- 4563** Anesthesia management for cesarean section in a pregnant woman with odontogenic infection: A case report
Ren YL, Ma YS
- 4569** Convulsive-like movements as the first symptom of basilar artery occlusive brainstem infarction: A case report
Wang TL, Wu G, Liu SZ
- 4574** Globe luxation may prevent myopia in a child: A case report
Li Q, Xu YX
- 4580** Computer tomography-guided negative pressure drainage treatment of intrathoracic esophagojejunal anastomotic leakage: A case report
Jiang ZY, Tao GQ, Zhu YF
- 4586** Primary or metastatic lung cancer? Sebaceous carcinoma of the thigh: A case report
Wei XL, Liu Q, Zeng QL, Zhou H
- 4594** Perianesthesia emergency repair of a cut endotracheal tube’s inflatable tube: A case report
Wang TT, Wang J, Sun TT, Hou YT, Lu Y, Chen SG
- 4601** Diagnosis of cytomegalovirus encephalitis using metagenomic next-generation sequencing of blood and cerebrospinal fluid: A case report
Xu CQ, Chen XL, Zhang DS, Wang JW, Yuan H, Chen WF, Xia H, Zhang ZY, Peng FH
- 4608** Primary sigmoid squamous cell carcinoma with liver metastasis: A case report
Li XY, Teng G, Zhao X, Zhu CM
- 4617** Acute recurrent cerebral infarction caused by moyamoya disease complicated with adenomyosis: A case report
Zhang S, Zhao LM, Xue BQ, Liang H, Guo GC, Liu Y, Wu RY, Li CY
- 4625** Serum-negative Sjogren's syndrome with minimal lesion nephropathy as the initial presentation: A case report
Li CY, Li YM, Tian M
- 4632** Successful individualized endodontic treatment of severely curved root canals in a mandibular second molar: A case report
Xu LJ, Zhang JY, Huang ZH, Wang XZ

- 4640** Successful treatment in one myelodysplastic syndrome patient with primary thrombocytopenia and secondary deep vein thrombosis: A case report
Liu WB, Ma JX, Tong HX
- 4648** Diagnosis of an extremely rare case of malignant adenomyoepithelioma in pleomorphic adenoma: A case report
Zhang WT, Wang YB, Ang Y, Wang HZ, Li YX
- 4654** Management about intravesical histological transformation of prostatic mucinous carcinoma after radical prostatectomy: A case report
Bai SJ, Ma L, Luo M, Xu H, Yang L
- 4661** Hepatopulmonary metastases from papillary thyroid microcarcinoma: A case report
Yang CY, Chen XW, Tang D, Yang WJ, Mi XX, Shi JP, Du WD
- 4669** PD-1 inhibitor in combination with fruquintinib therapy for initial unresectable colorectal cancer: A case report
Zhang HQ, Huang CZ, Wu JY, Wang ZL, Shao Y, Fu Z
- 4676** Cutaneous metastasis from esophageal squamous cell carcinoma: A case report
Zhang RY, Zhu SJ, Xue P, He SQ
- 4684** Rare pattern of Maisonneuve fracture: A case report
Zhao B, Li N, Cao HB, Wang GX, He JQ
- 4691** Suprasellar cistern tuberculoma presenting as unilateral ocular motility disorder and ptosis: A case report
Zhao BB, Tian C, Fu LJ, Zhang XB
- 4698** Development of plasma cell dyscrasias in a patient with chronic myeloid leukemia: A case report
Zhang N, Jiang TD, Yi SH
- 4704** Ovarian growing teratoma syndrome with multiple metastases in the abdominal cavity and liver: A case report
Hu X, Jia Z, Zhou LX, Kakongoma N

LETTER TO THE EDITOR

- 4709** Perfectionism and mental health problems: Limitations and directions for future research
Nazari N

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Jamir Pitton Rissardo, MD, Academic Research, Adjunct Associate Professor, Research Associate, Department of Medicine, Federal University of Santa Maria, Santa Maria 97105110, Brazil. jamirrissardo@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin, Production Department Director: Xu Guo, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

May 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Association between depression and malnutrition in pulmonary tuberculosis patients: A cross-sectional study

Xue-E Fang, Dan-Ping Chen, Ling-Ling Tang, Yan-Jun Mao

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Abdelbasset WK, Saudi Arabia

Received: July 16, 2021

Peer-review started: July 16, 2021

First decision: October 18, 2021

Revised: October 26, 2021

Accepted: March 25, 2022

Article in press: March 25, 2022

Published online: May 16, 2022



Xue-E Fang, Dan-Ping Chen, Ling-Ling Tang, Department of Tuberculosis, Shanghai Pulmonary Hospital Affiliated to Tongji University, Shanghai 200000, Shanghai Province, China

Yan-Jun Mao, Department of Nursing, Shanghai Pulmonary Hospital Affiliated to Tongji University, Shanghai 200000, Shanghai Province, China

Corresponding author: Yan-Jun Mao, PhD, Academic Fellow, Department of Nursing, Shanghai Pulmonary Hospital Affiliated to Tongji University, No. 507, Zhengmin Road, Yangpu District, Shanghai 200000, Shanghai Province, China. maoyanjuan_fk@tongji.edu.cn

Abstract

BACKGROUND

Depression has been reported to be prevalent in patients with pulmonary tuberculosis (PTB). Moreover, several clinical symptoms of PTB and depression overlap, such as loss of appetite and malnutrition. However, the association between depression and malnutrition in TB patients has not been fully elucidated.

AIM

To explore the association between depression and malnutrition in patients with PTB.

METHODS

This hospital-based cross-sectional study included patients with PTB in Shanghai Pulmonary Hospital Affiliated to Tongji University from April 2019 to July 2019. The Patient Health Questionnaire-9 (PHQ-9) scale was used to evaluate depression. The cut-off value was set at 10, and the nutritional state was determined by the body mass index (BMI). In addition, the Quality of Life Instruments for Chronic Diseases was employed to establish the quality of life (QOL). Univariable analysis and multivariable analysis (forward mode) were implemented to identify the independent factors associated with depression.

RESULTS

A total of 328 PTB patients were screened for analysis. Eight were excluded for missing demographic data, four excluded for missing nutrition status, and sixteen for missing QOL data. Finally, 300 PTB patients were subjected to analysis. We found that depressive state was present in 225 PTB patients (75%). The ratio of malnutrition in the depressive PTB patients was 45.33%. Our results revealed significantly lower BMI, hemoglobin, and prealbumin in the depression group

than in the control group ($P < 0.05$). Moreover, the social status differed significantly ($P < 0.05$) between the groups. In addition, glutamic pyruvic transaminase and glutamic oxaloacetic transaminase in the depression group were significantly higher than those in the control group ($P < 0.05$). Multivariable logistic regression analysis showed that BMI [odds ratio (OR) = 1.21, 95% confidence interval (CI): 1.163-1.257, $P < 0.001$] and poor social function (OR = 0.95, 95% CI: 0.926-0.974, $P = 0.038$) were independently associated with depression.

CONCLUSION

Malnutrition and poor social function are significantly associated with depressive symptoms in PTB patients. A prospective large-scale study is needed to confirm these findings.

Key Words: Depression; Pulmonary tuberculosis; Malnutrition; Quality of life; Comorbidity; China

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In this hospital-based cross-sectional study, the data of 300 pulmonary tuberculosis (PTB) patients were subjected to analysis. The ratio of malnutrition in tuberculosis-depression syndemic patients was 45.33%. We found that body mass index and poor social function were independently associated with depression. Our present findings suggest that the early diagnosis and management of depression in patients with PTB can decrease the burden of malnutrition and improve outcomes.

Citation: Fang XE, Chen DP, Tang LL, Mao YJ. Association between depression and malnutrition in pulmonary tuberculosis patients: A cross-sectional study. *World J Clin Cases* 2022; 10(14): 4395-4403

URL: <https://www.wjgnet.com/2307-8960/full/v10/i14/4395.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i14.4395>

INTRODUCTION

Pulmonary tuberculosis (PTB), a chronic wasting disease, is a chronic pulmonary infection which is caused by *Mycobacterium tuberculosis*. An estimated 9.0-11.1 million PTB cases were diagnosed in 2018 worldwide, 1.0 million of which were children[1]. PTB presents as a global public health problem, and the situation in developing countries, including China, is even worse. Despite the decreasing trend of PTB prevalence in China, PTB remains a considerable threat to public health due to the high number of PTB patients and the multidrug-resistant PTB burden[2]. Cumulative evidence revealed that depression was prevalent in people with chronic diseases[3]. In addition, the ratio of PTB patients with depression was higher than that in healthy populations[4]. In a hospital-based cross-sectional study conducted in Cameroon, Kehbila *et al*[5] found that more than 50% of PTB patients were affected by depression. In Manila, the Philippines, approximately 16.8% of the PTB patients reportedly had depression[6]. However, no hospital-based study has been published on the prevalence of this state in patients with PTB in China. Previous reports have evidenced that human immune deficiency virus infection, poor social support, and perceived stigma are risk factors for the development of depression in PTB patients [7-9]. Moreover, the depression in patients with PTB is associated with insufficient health care and poor treatment compliance, which has led to drug resistance, morbidity, and mortality[10], negatively affecting the health-related quality of life (QOL) of PTB patients[11,12]. Additionally, PTB patients were susceptible to malnutrition, with a ratio of malnutrition from 38.3% to 75.0%[13]. Furthermore, malnutrition also triggered PTB relapse and increased mortality[14,15]. Appropriate and timely intervention for malnutritional and/or depressed PTB patients is a medical need. We hypothesized that depression may be prevalent in malnutritional PTB patients in China. Therefore, in this study, we aimed to evaluate the association between depression and malnutrition in PTB patients in China.

MATERIALS AND METHODS

Study design

This is a hospital-based cross-sectional study, which was conducted from April to July 2019 in Shanghai Pulmonary Hospital Affiliated to Tongji University, China. Patients with PTB were consecutively recruited for analysis. The inclusion criteria were as follows: (1) Clear consciousness; (2) Ability to communicate; (3) Patients who have provided informed consent and voluntarily participated in this study; and (4) Age above 18 years. The following exclusion criteria were applied: (1) A history of mental




Figure 1 Flow chart. PTB: Pulmonary tuberculosis; PLICD-PT: Quality of Life Instruments for Chronic Diseases-Pulmonary Tuberculosis.

illness; (2) Complications, such as disturbance of consciousness, chronic respiratory failure, and pulmonary encephalopathy; (3) Metabolic-related diseases such as thyroid disease; (4) Requirements for continuous non-invasive or invasive ventilation; (5) Unstable hemodynamics; (6) Cardiac or renal insufficiency; and (7) Extrapulmonary tuberculosis. All subjects provided written informed consent. The study protocol was approved by the Ethics Committee (No. K19-146).

Data collection

Data including age, educational level, occupation, marital status, body mass index (BMI), income, comorbidity, treatment duration, hemoglobin (Hb), albumin, liver function [alanine transaminase (ALT) and aspartate aminotransferase (AST)], and medical cost origin were collected by nurses (Fang XE, Chen DP, and Tang LL) who received uniform training by face-to-face interviews. On post-admission day, the patient's height and weight were measured. The height was measured using a calibrated ruler (± 0.5 cm); the actual body weight was measured using a corrected scale (± 0.2 kg). BMI was calculated as [weight (kg)/height (m^2)]. Next, BMI was used to assess the nutritional status[16] (Supplementary Table 1), and BMI less than $18.5 \text{ kg}/\text{m}^2$ was considered to represent malnutrition[17].

Patient Health Questionnaire-9

Depression was evaluated by the Patient Health Questionnaire-9 (PHQ-9)[18], which consists of nine questions and has been validated in China with a Cronbach's alpha value higher than 0.8. Each item was scored as 0 (not at all), 1 (several days), 2 (more than half of the days), or 3 (nearly every day); the total score ranged from 0 to 27. A PHQ-9 value higher than 10 showed a higher susceptibility to depression[19]. Hence, the included PTB patients were divided into two groups: Depression and control, based on a PHQ-9 threshold of 10.

QOL assessment

The level of QOL was assessed by the Quality of Life Instruments for Chronic Diseases-Pulmonary Tuberculosis (QLICD-PT)[20], which has been validated in China with a Cronbach's alpha value higher than 0.7. The QLICD-PT includes three domains and a specific model: Physiological function (basic physiological function, independence, energy, and discomfort), psychological function (cognition, emotion, will, and personality), social function (interpersonal interaction, social support, and social role), and specific module (respiratory symptoms, systemic symptoms, drug side effects, and special psychology).

Sample size

The sample size for this study was calculated using the formula: $n = (z)^2 p(1-p)/e^2$, where z is 1.96 [the value at 95% confidence interval (CI)], e is the standard error (estimated at $1/8$), and p is the ratio of depression. We estimated that 50% of the PTB patients would develop depression. Considering a potential 20% loss, we established that at least 300 PTB patients for inclusion were required.

Statistical analysis

SPSS software (version 20.0 Chicago, IL, United States) was used to analyze the data. Continuous data

Table 1 Demographic characteristics

Variable	
Age, yr	35.96 ± 13.17
Gender	
Male	189 (63.00%)
Female	111 (37.00%)
Marital status	
Married	180 (60.00%)
Unmarried	115 (38.30%)
Divorced/separated	5 (1.70%)
Education	
Junior primary	153 (51.00%)
Senior primary secondary and above	147 (49.00%)
Occupation	
Unemployed	170 (56.67%)
Employed	130 (43.33%)
Monthly income, Yuan	
< 5000	159 (53.00%)
5000-10000	121 (40.30%)
> 10000	20 (6.70%)
TB treatment duration	
< 6 mo	144 (48.00%)
6-12 mo	49 (16.30%)
> 12 mo	107 (35.70%)
Comorbidity	
Diabetes mellitus	11 (3.70%)
Source of medical costs	
Medical insurance	203 (67.60%)
Self-supporting	66 (22.00%)
Publicly-funded	31 (10.30%)
BMI	19.71 ± 2.94
Hb (g/L)	122.21 ± 22.88
ALB (g/L)	40.25 ± 11.61
Prealbumin	200.13 ± 75.46
PHQ-9	16.45 ± 5.35
AST (U/L)	25.66 ± 52.64
ALT (U/L)	26.12 ± 52.16
QLICD-PT	
Physical domain	22.46 ± 3.90
Psychological domain	22.89 ± 8.13
Social domain	23.21 ± 6.50
Specific domain	39.38 ± 8.00

BMI: Body mass index; Hb: Hemoglobin; ALB: Albumin; ALT: Alanine transaminase; AST: Aspartate aminotransferase; PHQ-9: Patient Health Questionnaire-9; TB: Tuberculosis.

are presented as the mean \pm SD. Normality distribution was determined by the Shapiro-Wilk test. The Student's independent *t*-test or Mann-Whitney test was used depending on the normality. Categorical data are expressed as numbers (percentages) and were analyzed using the χ^2 test. Univariable analysis was applied to identify the independent factors which are associated with depression. To identify potential confounders, factors with $P < 0.1$ in the univariable analysis were entered into the multivariable logistic regression model and were assessed using the forward mode. $P < 0.05$ was considered to indicate statistically significant differences.

RESULTS

Demographic data

A total of 328 PTB patients were recruited. Of them, we excluded eight for missing data, four for missing nutrition indicators of blood, and sixteen for missing QLICD-PT scale data. Finally, 300 PTB patients (91.46%) were subjected to analysis (Figure 1). The mean age of the respondents was 35.96 (\pm 13.17; range 21-40, median 30) years. Of the patients included, 189 (63%) were men, 180 (60%) were married, 93 (31%) had undergraduate education, and 170 (56.67%) were unemployed (Table 1).

Factors associated with depression in PTB patients

Based on the PHQ-9 score at 10, the PTB patients were divided to depression ($n = 225$, 75%) and control ($n = 75$, 25%) groups. The ratio of malnutrition among depressive status with PTB patients was 45.33% (Table 2). No statistically significant differences were detected between the groups in age, gender, marital status, education level, occupation, monthly income, TB treatment duration, comorbidity, or origin of medical costs ($P > 0.05$). The values of BMI ($P < 0.001$), Hb ($P < 0.05$), prealbumin ($P < 0.05$), and social function of QLICD-PT ($P < 0.05$) in the depression group were significantly lower than those in the control group. In addition, AST and ALT in the depression group were significantly higher than those in the control group (Table 2). Finally, logistic regression analysis was used to evaluate the possible factors that influence depression. As can be seen in Table 3, BMI [odds ratio (OR) = 1.21, 95% confidence interval (CI): 1.163-1.257, $P < 0.001$] and poor social function (OR = 0.95, 95% CI: 0.926-0.974, $P = 0.038$) were independently associated with depression.

DISCUSSION

The assessment of the depressive state in patients with PTB using the PHQ-9 scale showed that 75% of the study subjects developed depression. In addition, the results of the present study also suggest that nutritional status and social function were independent risk factors for depression. In clinical practice, nutrition management and psychological counseling for PTB patients are highly necessary.

The prevalence of depression in the PTB patients included in the present study was estimated to be higher than that determined in other studies; for example, it was 41.1% in Nigeria[21], 61.1% in Cameroon[5], 56% in Pakistan[22], 54% in Ethiopia[23], and 69.6% in Liaoning Province of China[3]. In other investigations, the comorbidity of mental disorders in hospitalized patients ranged from 19%[24] to 80%[25]. In a study performed in the Philippines, the depressive state among PTB patients was 16.8% [6]. These findings suggest that the depressive state in PTB patients varies and is country-specific. In the present study, the depression ratio was 75%, which was higher than those in most of the published reports. This discrepancy may be due to differences in the sample size, race, country-specific features, patient populations (hospitalized or not), and the specific depression assessment tool implemented.

Here, we found that the ratio of malnutrition among depressive PTB patients was 45.33%. Patients with non-depression status had higher levels of BMI, Hb, and prealbumin than patients with depression. Furthermore, the ratio of anemia among depressive PTB patients was 86.32%, which may be due to the effect of TB on red blood cell production, such as decreased erythrocyte lifespan, poor erythrocyte iron incorporation, and decreased sensitivity to erythropoietin[16]. Masumoto *et al*[6] also recommended that additional attention should be paid to malnourished PTB patients and those with poor social support to identify depression. Therefore, nutritional support for PTB patients may be necessary.

Nutrition problems may be caused by mental health issues, and thus the symptoms of malnutrition and psychological distress may overlap[26]. In this study, we found an association between malnutrition and depression, in which the following factors might be involved or causative: (1) Depression may lead to loss of appetite and digestive dysfunction; (2) Continuous mental stimulation

Table 2 Factors associated with depression in pulmonary tuberculosis patients

	Depression status		P value
	PHQ 10 (n = 225)	PHQ < 10 (n = 75)	
Age, yr	36.20 ± 13.32	35.27 ± 12.27	0.598
Gender			0.371
Male	146 (64.40%)	44 (58.70%)	
Female	36 (80.00%)	31 (41.30%)	
Marital status			0.555
Married	134 (59.60%)	46 (61.30%)	
Single co-habiting	88 (39.10%)	27 (36.00%)	
Divorced/separated	3 (1.30%)	2 (2.70%)	
Educational level			0.809
Junior primary	114 (51.10%)	38 (50.67%)	
Senior primary, secondary, and above	111 (48.90%)	37 (49.33%)	
Occupation			0.998
Unemployed	123 (54.67%)	47 (61.84%)	
Employed	102 (45.33%)	28 (37.33%)	
Monthly income, Yuan			0.302
< 5000	123 (54.70%)	36 (48.00%)	
5000-10000	88 (39.10%)	33 (44.00%)	
> 10000	14 (6.20%)	6 (8.00%)	
TB treatment duration			0.559
< 6 mo	105 (46.70%)	39 (52.00%)	
6-12 mo	39 (17.30%)	10 (13.30%)	
> 12 mo	81 (36.00%)	26 (34.70%)	
Comorbidity			
Diabetes mellitus	7 (3.10%)	4 (5.33%)	0.376
Hemoptysis	29 (12.90%)	6 (7.80%)	0.254
Source of medical costs			0.503
Medical insurance	153 (68.00%)	50 (66.67%)	
Self-supporting	53 (23.60%)	13 (17.33%)	
Publicly-funded	19 (8.40%)	12 (16.00%)	
Nutrition related indicators			
BMI	19.26 ± 2.63	21.22 ± 3.45	< 0.001 ^a
Hb (g/L)	120.21 ± 24.37	128.79 ± 16.12	0.004 ^a
ALB (g/L)	40.81 ± 13.10	38.59 ± 4.60	0.254
Prealbumin	176.10 ± 2.68	210.96 ± 63.17	0.001 ^a
QLICD-PT			
Physical function	27.27 ± 53.00	8.04 ± 1.28	0.331
Psychological function	30.18 ± 8.55	23.01 ± 4.83	0.173
Social function	23.21 ± 6.66	29.01 ± 6.68	0.004 ^a
Specific function	40.03 ± 7.93	23.21 ± 6.00	0.562
Liver function			

ALT (U/L)	37.77 ± 40.92	34.04 ± 90.36	0.036 ^a
AST (U/L)	37.84 ± 38.41	33.25 ± 94.75	0.014 ^a

^a*P* < 0.05.

BMI: Body mass index; Hb: Hemoglobin; ALB: Albumin; ALT: Alanine transaminase; AST: Aspartate aminotransferase; PHQ-9: Patient Health Questionnaire-9; TB: Tuberculosis.

Table 3 Multivariable logistic regression

Variable	OR	95%CI	P value
BMI	1.21	1.163-1.257	< 0.001
Social function	0.95	0.926-0.974	0.038
Hb	0.99	0.982-1.012	0.67
Prealbumin	1.00	0.998-1.002	0.65
Psychological function	0.99	0.958-1.043	0.97
ALT	1.00	0.978-1.025	0.93
AST	0.99	0.967-1.025	0.75

OR: Odds ratio; CI: Confidence interval; Hb: Hemoglobin; ALT: Alanine transaminase; AST: Aspartate aminotransferase.

leads to serious vegetative nerve dysfunction and endocrine imbalance, which affects the body's absorption of nutrients; and (3) The disease itself can increase catabolism, promoting protein decomposition and reducing protein synthesis. In addition, a negative association between depression and poor social function may exist. There may be a vicious circle, including malnutrition, QOL, and depression. Malnutrition may aggravate depression and seriously affect the QOL, while the loss of appetite in depressed patients can lead to malnutrition.

This study is not without limitations. As it was hospital-based cross-sectional, the risk factors for depression in different treatment periods in patients with PTB could not be identified. Additionally, no additional validation of the depression and QOL scales was performed. Moreover, the nutritional status was evaluated by BMI, while many other indicators could also reflect the nutritional status. The energy intake was not assessed, which could have introduced bias. Furthermore, data of the severity of PTB were not collected. Socio-economic status was reported to be a confounding factor between nutritional status and depression[27]. However, we did not explore that association.

CONCLUSION

In conclusion, the findings of the present study suggest that depression is common in hospitalized PTB patients, and psychological counseling or management and nourishment adjustments may be needed. To confirm the findings of the present study, a well-designed prospective large-scale study is needed.

ARTICLE HIGHLIGHTS

Research background

It has been reported that depression is prevalent in patients with pulmonary tuberculosis (PTB). Moreover, several clinical symptoms of PTB and depression overlap, such as loss of appetite and malnutrition. However, the association between depression and malnutrition in TB patients has not been fully understood.

Research motivation

The present study aimed to explore the association between depression and malnutrition in patients with PTB.

Research objectives

The present study aimed to explore the association between depression and malnutrition in patients

with PTB.

Research methods

This hospital-based cross-sectional study included patients with PTB in Shanghai Pulmonary Hospital Affiliated to Tongji University from April 2019 to July 2019. The Patient Health Questionnaire-9 (PHQ-9) scale was used to evaluate depression and the cut-off value was set at 10, and the nutritional state was determined by the body mass index (BMI). In addition, the Quality of Life Instruments for Chronic Diseases was employed to quantify the quality of life (QOL). Univariable analysis and multivariable analysis (forward mode) were used to identify the independent factors associated with depression.

Research results

A total of 328 PTB patients were screened for analysis. Eight were excluded for missing demographic data, four excluded for missing nutrition status, and sixteen for missing QOL data. Finally, 300 PTB patients were subjected to analysis. It was found that depressive state was present in 225 PTB patients (75%). The ratio of malnutrition in the depressive PTB patients was 45.33%. It was found that BMI, hemoglobin, and prealbumin in the depression group were significantly lower than those in the control group ($P < 0.05$). Moreover, the social status ($P < 0.05$) significantly differed between the groups. In addition, glutamic pyruvic transaminase and glutamic oxaloacetic transaminase in the depression group were significantly higher than those in the control group ($P < 0.05$). Multivariable logistic regression analysis showed that BMI [odds ratio (OR) = 1.21, 95% confidence interval (CI): 1.163-1.257, $P < 0.001$] and poor social function (OR = 0.95, 95% CI: 0.926-0.974, $P = 0.038$) were independently associated with depression.

Research conclusions

Malnutrition and poor social function are significantly associated with depressive symptoms in PTB patients. A prospective large-scale study is needed to confirm these findings.

Research perspectives

Malnutrition and poor social function are significantly associated with depressive symptoms in PTB patients. A prospective large-scale study is needed to confirm these findings.

ACKNOWLEDGEMENTS

The authors appreciate the respective study institution for their help and the study participants for their cooperation in providing all necessary information.

FOOTNOTES

Author contributions: YJ Mao was the guarantor and designed the study; Fang XE, Chen DP, and Tang LL participated in the acquisition, analysis, and interpretation of the data, and drafted the initial manuscript; Fang XE, Chen DP, Tang LL, and Mao YJ revised the article critically for important intellectual content.

Institutional review board statement: The study protocol was approved by the Ethics Committee of Shanghai Pulmonary Hospital Affiliated to Tongji University (No. K19-146).

Informed consent statement: All study participants or their legal guardian provided informed written consent prior to study enrollment.

Conflict-of-interest statement: There are no conflicts of interest to report.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Xue-E Fang 0000-0003-0457-3554; Dan-Ping Chen 0000-0002-6265-7409; Ling-Ling Tang 0000-0003-2186-0627; Yan-Jun Mao 0000-0003-1272-4033.

S-Editor: Wang JJ

L-Editor: Wang TQ

P-Editor: Wang JJ

REFERENCES

- 1 **World Health Organization.** Global tuberculosis report 2018. [cited 15 June 2021]. Available from: <https://www.who.int/publications/i/item/9789241565646>
- 2 **Huang L, Li XX, Abe EM, Xu L, Ruan Y, Cao CL, Li SZ.** Spatial-temporal analysis of pulmonary tuberculosis in the northeast of the Yunnan province, People's Republic of China. *Infect Dis Poverty* 2017; **6**: 53 [PMID: 28335803 DOI: 10.1186/s40249-017-0268-4]
- 3 **Contini B.** Threats and organizational design. *Behav Sci* 1967; **12**: 453-462 [PMID: 6060437 DOI: 10.3389/fpsy.2018.00308]
- 4 **Shen TC, Wang CY, Lin CL, Liao WC, Chen CH, Tu CY, Hsia TC, Shih CM, Hsu WH, Chung CJ.** People with tuberculosis are associated with a subsequent risk of depression. *Eur J Intern Med* 2014; **25**: 936-940 [PMID: 25459212 DOI: 10.1016/j.ejim.2014.10.006]
- 5 **Kehbila J, Ekabe CJ, Aminde LN, Noubiap JJ, Fon PN, Monekosso GL.** Prevalence and correlates of depressive symptoms in adult patients with pulmonary tuberculosis in the Southwest Region of Cameroon. *Infect Dis Poverty* 2016; **5**: 51 [PMID: 27268138 DOI: 10.1186/s40249-016-0145-6]
- 6 **Masumoto S, Yamamoto T, Ohkado A, Yoshimatsu S, Querri AG, Kamiya Y.** Prevalence and associated factors of depressive state among pulmonary tuberculosis patients in Manila, The Philippines. *Int J Tuberc Lung Dis* 2014; **18**: 174-179 [PMID: 24429309 DOI: 10.5588/ijtld.13.0335]
- 7 **Olden K, Hempfling WP.** The 503-nm pigment of *Escherichia coli* B: characterization and nutritional conditions affecting its accumulation. *J Bacteriol* 1973; **113**: 914-921 [PMID: 4570612 DOI: 10.1186/s12888-015-0598-3]
- 8 **Naidoo P, Mwaba K.** Helplessness, depression, and social support among people being treated for tuberculosis in south africa. *Soc Behav Personal* 2010; **38**: 1323-1333 [DOI: 10.2224/sbp.2010.38.10.1323]
- 9 **Lee LY, Tung HH, Chen SC, Fu CH.** Perceived stigma and depression in initially diagnosed pulmonary tuberculosis patients. *J Clin Nurs* 2017; **26**: 4813-4821 [PMID: 28370819 DOI: 10.1111/jocn.13837]
- 10 **Gaze H.** Ethics: a question of morality. *Nurs Times* 1987; **83**: 18-19 [PMID: 3649695 DOI: 10.1155/2013/489865]
- 11 **Darby DW.** The dentist and continuing education--attitudes and motivations. *J Am Coll Dent* 1969; **36**: 165-170 [PMID: 5256348 DOI: 10.4046/trd.2017.80.1.69]
- 12 **Oppo GT, Segre A, Morelli L.** [Changes in serum leucine aminopeptidase during pregnancy, delivery and puerperium]. *Minerva Ginecol* 1969; **21**: 881-887 [PMID: 5398494 DOI: 10.1371/journal.pone.0174605]
- 13 **Cammer L.** Antidepressants as a prophylaxis against depression in the obsessive compulsive person. *Psychosomatics* 1973; **14**: 201-206 [PMID: 4794866 DOI: 10.4103/0366-6999.170255]
- 14 **Kant S, Gupta H, Ahluwalia S.** Significance of nutrition in pulmonary tuberculosis. *Crit Rev Food Sci Nutr* 2015; **55**: 955-963 [PMID: 24915351 DOI: 10.1080/10408398.2012.679500]
- 15 **Kumar A, Kakkar R, Kandpal S, Sindhwani G.** Nutritional status in multi-drug resistance-pulmonary tuberculosis patients. *IJCH* 2015; **26**: 204-208
- 16 **Portal RW.** Elective surgery after myocardial infarction. *Br Med J (Clin Res Ed)* 1982; **284**: 843-844 [PMID: 6802320 DOI: 10.1186/s12890-019-0953-0]
- 17 **World Health Organization.** Nutrition landscape information system (nlis) country profile indicators: Interpretation guide. [cited 16 June 2021]. Available from: <https://iris.who.int/handle/10665/44397>
- 18 **Menzel G.** [Equipment for recording isotonic muscle contractions with a mechanico-electric transducer]. *Z Med Labortech* 1971; **12**: 108-114 [PMID: 5573982 DOI: 10.1002/jclp.22390]
- 19 **Wignyosumarto S, Mukhlis M, Shirataki S.** Epidemiological and clinical study of autistic children in Yogyakarta, Indonesia. *Kobe J Med Sci* 1992; **38**: 1-19 [PMID: 1495268 DOI: 10.1046/j.1525-1497.2001.016009606.x]
- 20 **Cliffe P.** Measurement and recording during intensive patient care. *Postgrad Med J* 1967; **43**: 195-201 [PMID: 6042382 DOI: 10.1186/s12955-018-0960-5]
- 21 **Nishikawa S, Ueno A, Ishida H, Nagata T, Hamasaki A, Koda N, Kido J, Wakano Y.** [Gingival hyperplasia induced by nifedipine. Effect of nifedipine and EGF on cell proliferation in human gingival fibroblasts]. *Nihon Shishubyo Gakkai Kaishi* 1986; **28**: 168-175 [PMID: 3487604 DOI: 10.2329/perio.28.168]
- 22 **Amreen, Rizvi N.** Frequency of depression and anxiety among tuberculosis patients. *JTR* 2016; **4**: 183-190 [DOI: 10.4236/jtr.2016.44021]
- 23 **Ruel-Kellermann M.** [Women dentists]. *Inf Dent* 1984; **66**: 3265-3274 [PMID: 6397440 DOI: 10.1186/s12888-019-2042-6]
- 24 **Aydin IO, Uluşahin A.** Depression, anxiety comorbidity, and disability in tuberculosis and chronic obstructive pulmonary disease patients: applicability of GHQ-12. *Gen Hosp Psychiatry* 2001; **23**: 77-83 [PMID: 11313075 DOI: 10.1016/s0163-8343(01)00116-5]
- 25 **Sulehri MA, Dogar IA, Sohail H, Mehdi Z, Azam M, Niaz O, Javed SM, Sajjad IA, Iqbal Z.** Prevalence of depression among tuberculosis patients. *APMC* 2010; **4**: 133-137
- 26 **Punton S.** Burford Nursing Development Unit. Self-medication. *Nurs Times* 1985; **80**: 45 [PMID: 3851352 DOI: 10.3747/co.20.1651]
- 27 **Khalid S, Williams CM, Reynolds SA.** Is there an association between diet and depression in children and adolescents? *Br J Nutr* 2016; **116**: 2097-2108 [PMID: 28093091 DOI: 10.1017/S0007114516004359]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

