

13<sup>th</sup> September, 2021

To whom it may concern,

**Re: "Higher infliximab and adalimumab trough levels are associated with fistula healing in patients with fistulising perianal Crohn's Disease"**

This letter is to confirm that the statistical methods of this study were reviewed by Dr Wei Xuan, a biostatistician from the Ingham Institute for Applied Medical Research.

Yours sincerely,



Dr Bonita Gu

On behalf of **A/Professor Susan Connor**

Conjoint A/Professor UNSW/ Senior Staff Specialist

Department of Gastroenterology & Hepatology

Clinic 123 Level 1 Liverpool Hospital Liverpool NSW 2170

Locked Bag 7103. Liverpool BC NSW 1871

Tel +61-2- 8738 4085 | Fax +61-2- 8738 3094 |

**Working together to provide expert individualised care for Crohn's and Colitis**

[www.health.nsw.gov.au](http://www.health.nsw.gov.au)

**A/Prof Susan Connor**  
Senior Staff Specialist  
MBBS (Hons1 UNSW)  
BMed Sci (UNSW)  
FRACP  
PhD (UNSW)  
Provider No 2019835Y

**Dr Watson Ng**  
Staff Specialist  
MBBS (Hons USYD)  
FRACP  
PhD (UNSW)  
Provider No 2170784T

**Dr Astrid-Jane Williams**  
Staff Specialist  
BSc (UQ)  
MBBS (UQ)  
FRACP  
MHSc (UBC)  
Provider No 4117236F

**Alexandra Sechi**  
IBD Clinical Nurse Consultant  
B Nursing (UOW), Grad Dip Nursing  
(Gastroenterology) (UQ)

**Gordana Kuzet**  
IBD Clinical Nurse Specialist  
B Nursing (WSU)

**Aura Cabaero**  
IBD Registered Nurse  
B Nursing (USyd)